

NET CONTENTS

WEED KILLER

No. 125

10% SOLUTION

A Water-soluble Herbicide containing 2, 4-D

Active Ingredient:	
Mixed Isopropanolamine Salts of	16%
2, 4-Dichlorophenoxyacetic Acid	84%
Inert Ingredients:	
	Total 100%

Equivalent to 2, 4-Dichlorophenoxyacetic acid - 9.8 %

DIRECTIONS

Use at the rate of one gallon with water to make 125 gallons of spray, equivalent to one teaspoonful to 1 1/2 gallons water. Apply with hand or power sprayer in sufficient quantity to give thorough coverage of the weeds without excessive runoff or wetting of the soil. For turf, approximately 2 1/2 gallons per acre (one gallon to 200 square feet) will be required. More luxuriant foliage will require more spray. If extremely efficient spray equipment is available, it may be possible to obtain thorough coverage with a smaller volume of spray mixture, in which case the concentration of Weed Killer in the spray should be proportionately increased.

CAUTION

- Do not graze treated areas by dairy animals within 7 days after treatment.
- Do not use on newly seeded turf.
- Do not contaminate irrigation ditches or water used for domestic purposes.
- Do not apply to clover as injury may result.

The recommended concentration should be doubled for the difficult-to-control weeds, such as ~~vine~~ and ~~goldenrod~~. Double the concentration also for ~~poison ivy~~ growing in sunny locations. Control of this species in shady locations is doubtful. Deep-rooted ~~perennial weeds~~ such as ~~lamb's ears~~, ~~thistles~~, and ~~leafy spurge~~ are more difficult to kill and may require repeated applications.

KILLS
~~DANDELION - POISON IVY - POISON OAK - RAGWEED - PLAIN~~
~~AIN - (BROAD AND NARROW LEAF) - THISTLES - DOCKS~~
~~WILD MORNING GLORY - SHEPHERD'S PURSE - WILD CARROT~~
~~MALLOW~~

For lawns and ornamental turf in golf courses, parks, playgrounds, and recreation areas, non-crop areas such as along highways, railroad rights-of-way, ~~airfields~~ and around farm buildings.

Licensed under U.S. Pat. 2,322,761 and 2,326,471

CAUTION

KEEP OUT OF REACH OF CHILDREN

CAUTION - May Cause Skin Irritation. Avoid contact with eyes, skin and clothing. Avoid spray drift to susceptible plants as this product may injure cotton, beans, peas, ornamentals, etc. (Coarse sprays are less likely to drift.) Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes. Do not store near fertilizers, seeds, insecticides or fungicides.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.



MANUFACTURED BY
BAIRD & MCGUIRE, INC.
 HOLBROOK, MASS.

USDA REG.
 NO. 551-4

NET CONTENTS

WEED KILLER

No. 125

10% SOLUTION

(Water-soluble Herbicide containing 2, 4-D)

Active Ingredient:	
1. Mixed Isopropanolamine Salts of	16%
2. 4-Dichlorophenoxyacetic Acid	84%
REGISTERED INACTIVE INGREDIENTS:	
	Total 100%

Equivalent to 2, 4-Dichlorophenoxyacetic acid - 9.8%

KILLS

~~DANDELION - POISON IVY - POISON OAK - RAGWEED - PLAIN
LAIN - (BROAD AND NARROW LEAF) - THISTLES - DOCKS
WILD MORNING GLORY - SHEPHERD'S PURSE - WILD CARROT
MALLOW~~

~~For lawns and ornamental turf in golf courses, parks, playgrounds, and recrea-
tional areas, non-crop areas such as along highways, railroad rights-of-way, air-
fields, and around farm buildings.~~

Licensed under U.S. Pat. 2,322,761 and 2,326,471

CAUTION

KEEP OUT OF REACH OF CHILDREN

CAUTION - May Cause Skin Irritation. Avoid contact with eyes, skin and clothing. Avoid spray drift to susceptible plants as this product may injure cotton, beans, peas, ornamentals, etc. (Coarse sprays are less likely to drift.) Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes. Do not store near fertilizers, seeds, insecticides or fungicides.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.



MANUFACTURED BY
BAIRD & MCGUIRE, INC.
HOLBROOK, MASS.

USDA REG.
NO. 551-A

Apply with pressures of 75 to 250 pounds per square inch where dense foliage occurs. Higher pressures may cause matting of vegetation and make thorough coverage impossible. Apply during warm weather when temperature is at least 60° F. and when weeds are growing actively. Apply to lawns between mowings so that a maximum of foliage is present.

IMPORTANT

Avoid application to vegetables, flowering plants, shrubbery, trees or other desirable type of plants. This is particularly necessary when treating weeds in orchards, vineyards and nurseries.

Do not use on bent grass lawns, except for spot spraying.

Always clean spraying equipment thoroughly before using for any other purpose. Rinse several times with hot water containing soap or trisodium phosphate (one pound in twenty-five gallons of water). Allowing the tank to stand filled with water overnight is recommended. Be sure to clean out hose and spray nozzle.

WARNING: Do not dispose of poison ivy by burning after it has been killed. Smoke from the burning vine is very poisonous on contact with the skin.

54A

54

WEED KILLER

No.
10% SOL

Water-soluble Herbicide

Active Ingredient:
Mixed Isopropanolamine
2, 4-Dichlorophenoxy
Acetic Acid

Equivalent to 2, 4-Dichlorophenoxy


DIRECTIONS

Use at the rate of one gallon with water to make 125 gallons of spray, equivalent to eight teaspoons to 1 1/4 gallons water. Apply with hand or power sprayer in sufficient quantity to give thorough coverage of the weeds without excessive runoff or wetting of the soil. For turf, approximately 218 gallons per acre (one gallon to 200 square feet) will be required. More luxuriant foliage will require more spray. If extremely efficient spray equipment is available, it may be possible to obtain thorough coverage with a smaller volume of spray mixture, in which case the concentration of Weed Killer in the spray should be proportionately increased.

CAUTION

- Do not graze treated areas by dairy animals within 7 days after treatment.
- Do not use on newly seeded turf.
- Do not contaminate irrigation ditches or water used for domestic purposes.
- Do not apply to clover as injury may result.

The recommended concentration should be doubled for the difficult-to-control weeds, such as yarrow and goldenrod. Double the concentration also for poison ivy growing in sunny locations. Control of this species in shady locations is doubtful. Deep-rooted perennial weeds such as bindweed, Canada thistle, and leafy spurge are more difficult to kill and may require repeated applications.



KILL
DANDELION - POISON IVY - POISON
TAIN - (BROAD AND NARROW LEAF)
WILD MORNING GLORY - SHEPHERD'S
BELL

For lawns and ornamental turf in golf courses, residential areas, non-crop areas such as along roads, lanes, airfields and around farm buildings.

Licensed under U.S. Pat. 2,422,000

CAUTION

KEEP OUT OF REACH OF CHILDREN

CAUTION — May Cause Skin Irritation. Avoid contact with clothing. Avoid spray drift to susceptible plants, such as beans, peas, ornamentals, etc. (Coarsely clean spray equipment with a suitable detergent for other purposes. Do not store near fertilizers.)

Rinse equipment and containers and dump contents away from water supplies. Contain and dispose of holes in them and burying with wastes.



MANUFACTURED BY
BAIRD & MCGEE
HOLBROOK, MASS.