

1 GALLON

NO-ROMA DISINFECTANT SANITIZER and DEODORANT

Active Ingredient:
n-Alkyl (50% C14, 40% C12, 10% C16) *dimethyl benzyl ammonium chlorides* 10%
Inert Ingredient: Water not over 90%
Minimum Phenol Coefficients (A. O. A. C. —20°C)
S. Aureus 100
S. Typhosa 70

Directions for Dairy, Restaurant, Industrial and Home Use.

For sanitation of dishes, glassware, cooking utensils, silverware, bottles: Clean with suitable detergent, rinse thoroughly, then dip in a sanitizing rinse made with 1 ounce of disinfectant to 4 gallons of water.

Give final rinse in clean, potable water.
AIR DRY — DO NOT USE A TOWEL. —

For sanitation of milk cans and miscellaneous dairy equipment: Clean and rinse thoroughly. Wash in disinfectant solution using 1 ounce disinfectant to 4 gallons of water; or spray, using 1 ounce disinfectant to 2 gallons of water. Give final rinse in clean, potable water.

NO-ROMA fulfills the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinance 1965 Recommendations of the U.S. Public Health Service in waters up to 550 p.p.m. of hardness calculated as CaCO₃ when tested by the A.O.A.C. Germicidal and Detergent Sanitizers-Official Method.

For general use in disinfecting bath tubs, toilet seats, floors, walls, banisters, showers, telephones: Apply freely after cleaning using 1 ounce disinfectant to 2 gallons of water.

Unexcelled as a deodorant. Use 1 oz. disinfectant per 1 gallon of water.

ODORLESS

DANGER

KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician immediately.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock as well as oxygen and measures to support breathing manually or mechanically may be needed. If persistent, convulsions may be controlled by the cautious intravenous injection of a short-acting barbiturate drug.

EPA REG. NO. 551-17

BAIRD & MCGUIRE, INC.
HOLBROOK, MASS.

Directions for the Barber or Beauty Shop Operator

To sanitize brushes, combs, scissors, razors, curlers, manicure bowls, and other equipment:

(1) Wash instruments being careful that they are free from hair and other accumulations.

(2) Rinse thoroughly in clear water

(3) Dip in solution using 1 tablespoonful disinfectant to 1 quart of water for at least one minute, being careful to immerse instruments completely.

(4) Remove from solution, air-dry and store in a clean dry place.
Solution should be changed twice a week, or sooner if it clouds.

Rinse empty container thoroughly with water and discard it.



ACCEPTED

NOV 06 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
EPA UNDER NO. 551-17