

DIRECTIONS FOR USE

For use on certain agricultural crops and ornamentals and for fly and mosquito control.

APPLES. For control of apple maggot, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of European spruce sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

PEARS. For control of pear sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of pear codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

PEACHES. For control of peach codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of peach sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

PLUMS. For control of plum codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of plum sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

CHERRIES. For control of cherry codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of cherry sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ALMONDS. For control of almond sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of almond codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

ORCHARD FRUIT TREES. For control of orchard fruit tree sawfly, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water. For control of orchard fruit tree codling moth, apply 1 to 2 quarts of concentrate per acre, diluted in 100 to 200 gallons of water.

MALATHION 5 LB. EMULSIFIABLE CONCENTRATE

For use on certain agricultural crops and ornamentals and for fly and MOSQUITO CONTROL

ACTIVE INGREDIENTS:

Malathion*	57%
Heavy Aromatic Naptha	34%

INERT INGREDIENTS: 9%

TOTAL 100%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate
Contains 5 pounds of Malathion per U. S. gallon at 70°F.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED

Avoid breathing of spray mist
Avoid contact with skin. Wash thoroughly
Avoid contamination of feed and foodstuffs.

Do not use in commercial food preparation areas or in edible products areas of food processing plants.

NOTE TO PHYSICIANS: Malathion may cause cholinesterase inhibition. Atropine is antidotal.

ACCEPTED

AUG 1 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this pesticide registered under EPA Reg. No. 551-131

ACTIVE INGREDIENTS:

Malathion*

57%

Heavy Aromatic Naptha

34%

INERT INGREDIENTS:

9%

TOTAL 100%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate
Contains 5 pounds of Malathion per U. S. gallon at 70 °F.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED

Avoid breathing of spray mist.

Avoid contact with skin. Wash thoroughly

Avoid contamination of feed and foodstuffs.

Do not use in commercial food preparation areas or in edible products areas of food processing plants.

NOTE TO PHYSICIANS: Malathion may cause cholinesterase inhibition.
Atropine is antidotal.

SEE OTHER CAUTIONS ON SIDE PANEL

APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL

Do not use or store near heat or open flame

U.S. STD. MEASURE GALLONS

EPA Reg. No. 551-131/EPA Est 551-MA-1

ACCEPTED

AUG 1 1975

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 551-131

MALATHION 5 LB. EMULSIFIABLE CONCENTRATE

For use on certain agricultural crops and ornamentals,
and for fly and MOSQUITO CONTROL

ACTIVE INGREDIENTS:

Malathion*	57%
Heavy Aromatic Naptha	34%

INERT INGREDIENTS:	9%
TOTAL	100%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate
Contains 5 pounds of Malathion per U. S. gallon at 70°F.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED

Avoid breathing of spray mist.

Avoid contact with skin. Wash thoroughly.

Avoid contamination of feed and foodstuffs.

Do not use in commercial food preparation areas or in edible products areas of food processing plants.

NOTE TO PHYSICIANS: Malathion may cause cholinesterase inhibition

DIRECTIONS FOR USE (Cont.)

BEETS To control or exclude apply 1 to 2 pints (10 to 20 ounces actual Malathion) per 100 gallons of water. Make full-coverage applications, repeated as necessary.

Do not apply within 10 days of harvest if tops are to be used as feed.

COTTON To control aphids and spider mites, make two or more applications, as needed, of a spray containing 1 to 1½ pints (10 to 15 ounces actual Malathion) per acre.

Do not apply within 7 days of picking.

SEED ALFALFA To control Pacific spruce saw spotted mite and aphids, use 1 to 2 pints (15 to 20 ounces actual Malathion) per 50 to 100 gallons of water, per acre, by ground sprayer, or 10 gallons of water per acre by aerial.

Do not apply within 21 days of harvest.

ORNAMENTALS Make thorough, full-coverage sprays with repeated applications, as necessary. For scales, apply when crawlers have settled in foliage injury may occur on certain trees, including Boston, Mazzoni, and Pines, as well as some species of Crataegus. Apply 1 pint (16 ounces actual Malathion) per 100 gallons of water to control lacebugs and cysteids, etc.

Apply 1 pint (16 ounces actual Malathion) per 100 gallons of water to control aphids, 4 to 6 pints (64 to 96 ounces actual Malathion) per 100 gallons of water to control scales, and 1 to 2 pints (16 to 32 ounces actual Malathion) per 100 gallons of water to control mites.

For control of scales on trees and shrubs, apply 1 to 2 pints (16 to 32 ounces actual Malathion) per 100 gallons of water. For control of scales on fruit trees, apply 1 to 2 pints (16 to 32 ounces actual Malathion) per 100 gallons of water.

For control of scales on fruit trees, apply 1 to 2 pints (16 to 32 ounces actual Malathion) per 100 gallons of water. For control of scales on fruit trees, apply 1 to 2 pints (16 to 32 ounces actual Malathion) per 100 gallons of water.

For control of scales on fruit trees, apply 1 to 2 pints (16 to 32 ounces actual Malathion) per 100 gallons of water. For control of scales on fruit trees, apply 1 to 2 pints (16 to 32 ounces actual Malathion) per 100 gallons of water.

For control of scales on fruit trees, apply 1 to 2 pints (16 to 32 ounces actual Malathion) per 100 gallons of water. For control of scales on fruit trees, apply 1 to 2 pints (16 to 32 ounces actual Malathion) per 100 gallons of water.

For control of scales on fruit trees, apply 1 to 2 pints (16 to 32 ounces actual Malathion) per 100 gallons of water. For control of scales on fruit trees, apply 1 to 2 pints (16 to 32 ounces actual Malathion) per 100 gallons of water.

ACCEPTED

AUG 1 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 551-131

ACTIVE INGREDIENTS:

Malathion*	57%
Heavy Aromatic Naptha	34%

INERT INGREDIENTS: 9%

TOTAL 100%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate
Contains 5 pounds of Malathion per U. S. gallon at 70°F.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED

Avoid breathing of spray mist.

Avoid contact with skin. Wash thoroughly.

Avoid contamination of feed and foodstuffs.

Do not use in commercial food preparation areas or in edible products areas of food processing plants.

NOTE TO PHYSICIANS: Malathion may cause cholinesterase inhibition.
Atropine is antidotal.

SEE OTHER CAUTIONS ON SIDE PANEL

APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

Do not use or store near heat or open flame.

U.S. STD. MEASURE GALLONS

EPA Reg. No. 551-131/EPA Est 551-MA-1

ACCEPTED

AUG 1 1975

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. **551-131**

MOSQUITO CONTROL

MOSQUITO LARVAE IN STANDING WATER - INTERMITTENTLY FLOODED AREAS, STAGNANT WATER, TEMPORARY RAINPOOLS, ETC., MAY BE FOUND IN THE AREA
of 100 ft. x 100 ft. around the house. Mosquito larvae are found in the area by an

OUTDOORS MOSQUITOES FEELS THE SWARM. IT YOUNG THIS OF IS READY TO BE
 THE FIRST OF THE SEASON TO BE SEEN IN THE SWAMP. IT IS THE FIRST OF THE SEASON
 TO BE SEEN IN THE SWAMP. IT IS THE FIRST OF THE SEASON TO BE SEEN IN THE SWAMP.

[illegible]

CAUTION

1. 1995年5月15日，在《中国日报》发表题为“中国：从‘韬光养晦’到‘有所作为’”的文章，指出中国外交政策的重大转变。

$$1.000 \pm 0.009 \quad \text{and} \quad 1.000 \pm 0.009$$

$$+2.37 \times 10^{-6} \pm 3.21 \times 10^{-6}$$

Do not apply when air is below 50°F.
Do not apply when wind is above 15 mph.
Do not apply when rain is expected within 2 hours.
Do not apply when temperature is below 50°F.

[illegible]