

NET CONTENTS:

# METHOXYCHLOR 25

## { 25% EMULSIFIABLE SOLUTION }

CONTAINS 2 LBS. METHOXYCHLOR PER GALLON

FOR CONTROL OF INSECT PESTS OF  
CROPS AND STORED GRAIN

ACCEPTED  
OCT 2 1974  
UNDER THE FEDERAL INSECTICIDE ACT  
FUNGICIDE ACT, AND OTHER ACTS  
FOR ECONOMIC PEST CONTROL  
REGISTERED UNDER NO. 551-139


### ACTIVE INGREDIENTS:

Methoxychlor* technical .....	25%
Xylene .....	63%
Aromatic Petroleum Derivative Solvent .....	7%
INERT INGREDIENTS: .....	5%
Total .....	100%

\*Equivalent to 22% of 2,2-bis(4-methoxyphenyl)-1,1,1-trichloroethane and 3% of other isomers and related compounds.

## WARNING

### KEEP OUT OF REACH OF CHILDREN



This product is toxic to honey bees when bees are actively visiting the area.

EPA Reg. No. 551-139 EPA Est. 551-MA-1

### DIRECTIONS FOR USE

**HOUSEFLIES AND STABLE FLIES (In Farm Buildings other than Dairy and Dairy Barns):** Use 1 gal. per 10 gals. of water and apply at the rate of 100 to 200 gal. per 1000 sq. ft. of surface. Spray particularly those areas where flies crawl or rest and congregate. Repeat application as infestations warrant.

**BEANS (Mexican Bean Beetles, Leafhoppers):** Apply 2 qts. per acre in sufficient water to give good coverage of both upper and lower leaf surfaces. Repeat in 2 weeks if infestations warrant.

**SOYBEANS, PEANUTS (Velvetbean Caterpillars):** Apply 2 qts. per acre in sufficient water to give coverage. Treat at first sign of infestation and repeat in 10 days if infestation warrants.

**SEEDS**  
**ALFALFA and CLOVER—Seed or Hay—(Alfalfa Weevils, Alfalfa Caterpillars, Lygus Bugs, Spittle Bugs, Leafhoppers):** Apply 2-3 qts. per acre in sufficient water to cover. Begin application at first sign of infestation and repeat as infestations warrant. Do not treat during bloom.

**IMPORTANT: DO NOT APPLY TO CROPS WITH 30 DAYS BEFORE HARVEST.**

**MOSQUITO CONTROL (Non-Agricultural Land Parks, Beaches, Public & Residential Areas, and Dump, Marsh, and Sloughs and Low Lying Areas):** Apply 1/2 to 1 gal. per 1000 sq. ft. of surface. Spray particularly those areas where mosquitoes breed and rest. Repeat application as infestations warrant.

### DIRECTIONS FOR USE

**MIST BLOWERS:** 4 gallons emulsifiable concentrate to 96 gallons water at 5 gal. nozzle and 200 ft. nozzle; 1 1/2 pints to 39 pints water at 5 gal. nozzle and 200 ft. nozzle.

For all types of application thorough even coverage is essential for effective control.

**ELM BARK BEETLES** **ELM TREES**

**How to Use:**  
**Hydraulic Sprayers:** Mix 2 1/2 gal. of concentrate with 100 gal. of spray. Apply enough spray to thoroughly wet all bark on trunk of tree. Use 200 to 250 gal. of spray (about 200 to 250 ft. of trunk).  
**Mist Blowers:** Mix 1 1/2 gal. of concentrate with 39 gal. of spray. Thoroughly cover all bark on trunk. 100 gal. of spray should cover 200 to 250 ft. of trunk.

**When to use:**  
Apply as a dormant treatment in late fall or early spring when leaves appear, when temperature is 40 to 60 F. and there is no danger of freezing temperatures.

## WARNING

DO NOT USE IN FOOD AREAS

**Fish and Wildlife WARNING —** This product is toxic to Fish. Keep out of lakes and streams and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.