

NET CONTENTS GALLONS

**DIRECTIONS FOR USE**

LV-6T BRUSH KILLER is designed for use in control of brush and woody plants. Its principal use is on rangeland clearance, power lines, rights-of-way, stump land, and certain waste areas where it is desired to kill or repress growth of woody plants. This product is not intended for weed control in growing crops.

**FOLIAGE TREATMENT:** Foliage or leaf stem sprays are used during the growing season on dense brush or where vine growths such as Smilax, Greenbrier, and other common woody vines are a problem. Apply LV-6T after foliage is well developed, using a drenching spray containing 2 to 2-2/3 quarts in 100 gallons of water. Coverage should be complete, wetting all parts of the plants, including leaves and stems to the point of run-off. The best time for spraying is when soil moisture levels are favorable for growth. DO NOT treat during periods of severe drought or in early fall when leaves have lost their healthy green color.

**BASAL BARK TREATMENT:** Brush and small trees can be controlled anytime of the year by spraying the basal parts of stems and tree trunks, less than 6 inches in diameter, from the ground line up to 12 or 16 inches in height. Apply a solution of 2-2/3 gallons of LV-6T in 97-1/3 gallons (2-1/5 quarts in 25 gallons, or 3/4 pint in 3 gallons) of diesel oil, fuel oil, or kerosene, completely wetting the indicated area until run-off is noticed at the ground line. Frequently a delayed response and killing can be expected.

**STUMP TREATMENT.** In areas where brush and trees have been cut, all stumps and stubs should be treated to prevent regrowth. Apply LV-6T at the rate of 2-2/3 gallons in 97-1/3 gallons (2-1/5 quarts in 25 gallons, or 3/4 pint in 3 gallons) of fuel oil, diesel oil or kerosene. Wet thoroughly all exposed areas such as cut surfaces, surface roots, and body of stump to ground line. Apply anytime, including winter, preferably to freshly cut stumps.

**AIRPLANE APPLICATION:** To control mesquite use 1 pint of LV-6T in 1 gallon of diesel oil and 3 gallons of water per acre. Best results are obtained when treatments are made 50 to 80 days after mesquite first begins to leaf out in the spring. DO NOT treat if drought prevents full foliage. For control of moderate to dense stands of Alder, Bur, Post, Blackjack, Northern Red and Northern Pin Oaks in conifer plantings, apply 1 to 2 quarts in a solution of 3 to 4 gallons of water and 1 gallon of diesel oil, or in a solution of 2 to 5 gallons of diesel oil per acre. To avoid injury to tender new growth, spray conifer stands immediately after annual growth is completed.

**Brush Killer**  
(Low Volatile Emulsifiable Concentrate)

**LV-6T**

Contains 6.0 Pounds  
2, 4, 5-Trichlorophenoxyacetic Acid\*

**ACTIVE INGREDIENT:**

Isooctyl Ester of 2, 4,  
5-Trichlorophenoxyacetic Acid\* .. 86.4%

**INERT INGREDIENTS:** ..... 13.6%


**TOTAL** ..... 100.0%

\* Equivalent to 2,4,5-Trichlorophenoxyacetic Acid\*

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**  
**SEE SIDE PANEL FOR ADDITIONAL CAUTIONS**

Manufactured by  
**BAIRD & MCGUIRE, INC.**  
HOLBROOK, MASSACHUSETTS



USDA  
Reg. No. 551-

**GENERAL INFORMATION**

LV-6T BRUSH KILLER is an emulsifiable concentrate which forms an emulsion in water and will mix readily with water-free diesel oil, fuel oil, or kerosene

**PREPARING THE SPRAY:** Fill the spray tank with half the required amount of water or water-free diesel oil, fuel oil, or kerosene, add the recommended amount of LV-6T emulsifiable concentrate with agitation, and continue filling the spray tank with water or petroleum oil. In straight oil mixtures do not let water get into the LV-6T itself or in the finished product.

**CAUTION:** Avoid spray drift to susceptible plants such as cotton, tomatoes, flowers, grapes, fruit trees, and ornamentals. Spray drift may cause severe injury to both growing and dormant plants. Coarse sprays are less likely to drift. Although this ester is much less volatile than butyl or isopropyl esters, high temperatures may produce vapors after application which might injure 2,4D and 2,4,5T-sensitive plants growing nearby. Contamination of shallow fish-bearing waters at application rates recommended on this label may kill fish.

DO NOT graze treated areas by dairy animals within seven days after treatment.

DO NOT contaminate irrigation ditches or water used for domestic purposes.

DO NOT use the same spray equipment for applying other materials to plants, as injury will result.

DO NOT store near fertilizers, seeds, insecticides, or fungicides.

DO NOT take internally. Avoid contact with skin, eyes, or clothing.

We guarantee the contents of this package to conform to the labeled composition. The use of this material being beyond our knowledge and control, we do not make any warranty, express or implied, as to the effects of such use whether or not in accordance with the directions.