




NET CONTENTS:

BAIRD'S WEED KILLER

A DINITRO WEED KILLER

FOR CONTROLLING MOST ANNUAL WEEDS AND GRASSES AND KILLING TOP
GROWTH OF HERBACEOUS PERENNIAL WEEDS AND GRASSES; ALSO FOR
PREHARVEST SPRAYING OF POTATOES AND OF SEED CROPS OF
LEGUMES AND SOYBEANS TO FACILITATE HARVEST.

Use on: Airfields, Driveways, Walks, Dry Yards, Fence Lines, Railroads, Roadsides,
Waste Places, and around Industrial Establishments, Pole Yards and
Oil Tank Areas for Controlling Vegetation.

DINITROPHENOL SOLUTION

Active Ingredient:	
DINOSEB (2-sec-butyl-4, 6-dinitrophenol)	55%
Inert Ingredients:	45%
Total	100%

Contains 5 pounds of 4, 6 Dinitro-o-sec-butylphenol per gallon.

NOT FOR USE OR STORAGE IN OR AROUND HOME!



**DANGER
POISON**



KEEP OUT OF REACH OF CHILDREN

POISONOUS IF SWALLOWED OR ABSORBED THROUGH SKIN
DO NOT BREATHE VAPORS OR SPRAY MIST

KEEP AWAY FROM HEAT AND OPEN FLAME
DO NOT WEAR CONTAMINATED CLOTHING OR SHOES

COMBUSTIBLE

EPA Reg. No. 551-195
EPA EST. 551-MA-1

BAIRD & McGUIRE, INC.
Manufacturing Chemists
HOLBROOK, MASSACHUSETTS, U. S. A.

DIRECTIONS

For control of weeds and grasses, mix with the proper amount of water and spray thoroughly on the foliage of the weeds and grasses. The low toxicity of the weed killer gives below the satisfactory for control of weeds and grasses. For control of weeds and grasses, mix with the proper amount of water and spray thoroughly on the foliage of the weeds and grasses. For control of weeds and grasses, mix with the proper amount of water and spray thoroughly on the foliage of the weeds and grasses.

MIXING INSTRUCTIONS: For oil-water sprays, add approximately 1/2 gallon of water per gallon of the spray tank in a separate container. Thoroughly premix the proper amounts of Baird's Weed Killer and oil. With vigorous agitation add this premix to the water. Add the remaining required amount of water. Agitate just before use. For tank sprays, mix with from 1/2 to 1 gallon of water per gallon of the spray tank. Add the remaining required amount of water. Agitate just before use. For oil sprays, mix with from 1/2 to 1 gallon of water per gallon of the spray tank. Add the remaining required amount of water. Agitate just before use.

GENERAL WEED CONTROL: For control of weeds and grasses, mix with the proper amount of water and spray thoroughly on the foliage of the weeds and grasses.

CONTACT PRE-EMERGENCE WEED CONTROL: For control of weeds and grasses, mix with the proper amount of water and spray thoroughly on the foliage of the weeds and grasses.

WEED CONTROL IN ALLIALLY: For control of weeds and grasses, mix with the proper amount of water and spray thoroughly on the foliage of the weeds and grasses.

PREHARVEST SPRAYING OF ANNUAL CROPS: For control of weeds and grasses, mix with the proper amount of water and spray thoroughly on the foliage of the weeds and grasses.

PREHARVEST SPRAYING OF PERENNIAL CROPS: For control of weeds and grasses, mix with the proper amount of water and spray thoroughly on the foliage of the weeds and grasses.

PREHARVEST SPRAYING OF SEED CROPS: For control of weeds and grasses, mix with the proper amount of water and spray thoroughly on the foliage of the weeds and grasses.

PREHARVEST SPRAYING OF LEGUMES AND SOYBEANS: For control of weeds and grasses, mix with the proper amount of water and spray thoroughly on the foliage of the weeds and grasses.

PREHARVEST SPRAYING OF POTATOES: For control of weeds and grasses, mix with the proper amount of water and spray thoroughly on the foliage of the weeds and grasses.

PREHARVEST SPRAYING OF OTHER CROPS: For control of weeds and grasses, mix with the proper amount of water and spray thoroughly on the foliage of the weeds and grasses.

DIRECTED SPRAY TO GROUND COVER: For Citrus, Pine and Nut trees including almonds, filberts, pecans and walnuts, Dutch Banks along drainage and irrigation canals. Mix 2 to 3 pints of Baird's Weed Killer in 10 to 25 gallons of water or any weed oil and enough water to make 100 gallons of spray. Apply as coarse spray and thoroughly cover all weed foliage. On the crop areas do not apply more than 100 gallons of spray per acre. Avoid getting any of the spray on the fruits. Destroy any such fruit accidentally sprayed. Avoid spraying the base of young trees as girdling may occur. Do not allow livestock to graze on treated ground cover.

CONTROL OF THE BASAL SPIKE PHASE OF DOWNY MILDEW ON HOPS: Control is obtained by killing the basal parts of the hop plants. Mix 1 part of Baird's Weed Killer with 5 gallons of water or any weed oil and enough water to make 100 gallons of spray. Provide sufficient agitation when mixing and apply spraying is finished. Make the first application to the plants at the basal two feet of the hop bines when they have reached eight feet of height. Repeat the rate of 50 gallons per acre. Repeat as necessary, but use no more than four applications per growing season. When the plants have reached 4 of the distance to the top of the trees, the basal 4 feet of the bines should be sprayed. The gallonage required for the treatment of the basal four feet of the bines will need to be increased proportionately. The treatment will water and soil for the control of weeds.

USE PRECAUTIONS:

Do not spray entire vines. Do not apply to young replants in mature plantings or to plants in early yards. Do not apply within 14 days of harvest.

IMPORTANT: Do not use at temperatures below 20 degrees Fahrenheit. Do not use on plants that are stressed or injured by frost, drought or other factors.

First Aid: In case of contact with skin, wash immediately with plenty of water. In case of contact with eyes, flush immediately with plenty of water. In case of contact with clothing, remove clothing immediately. In case of contact with shoes, remove shoes immediately.

PREVENTIVE TREATMENT: ANTIDOTE

SYMPTOMS OF POISONING: Nausea, vomiting, diarrhea, dizziness, headache, and other symptoms.

FIRST AID: In case of contact with skin, wash immediately with plenty of water. In case of contact with eyes, flush immediately with plenty of water. In case of contact with clothing, remove clothing immediately. In case of contact with shoes, remove shoes immediately.

IF SWALLOWED, SEND FOR A PHYSICIAN: Do not induce vomiting. Give the patient plenty of water to drink. If the patient is unconscious, do not give anything by mouth. If the patient is conscious, give the patient plenty of water to drink. If the patient is unconscious, do not give anything by mouth. If the patient is conscious, give the patient plenty of water to drink.

NOTE TO VETERINARIAN:

NOTICE: Baird's Weed Killer is a potent herbicide and should be used with care. It is not to be used on food crops. It is not to be used on ornamental plants. It is not to be used on lawns. It is not to be used on golf courses. It is not to be used on tennis courts. It is not to be used on swimming pools. It is not to be used on playgrounds. It is not to be used on school grounds. It is not to be used on church grounds. It is not to be used on cemetery grounds. It is not to be used on any other area where it may be harmful to humans or animals.