

NET CONTENTS

MIXING INSTRUCTIONS

This concentrate will not form emulsions that are stable indefinitely. To insure good performance:

1. Prepare the emulsion as soon before actual use as possible.
2. Pour the proper amount of concentrate slowly into the water with constant agitation. The agitation shall be as vigorous as possible and shall be continued for at least 2 minutes after the last of the concentrate has been added.
3. The emulsion shall again be thoroughly agitated for at least 1 minute immediately before use.

DIRECTIONS FOR USE

RESIDUAL SPRAY IN BUILDING

Mosquitoes (Around doors, screens and wall surfaces wherever the insects light or congregate - For Spot Application Only.)

Add 1 part of emulsifiable concentrate to 38 parts of water (3 1/2 ounces of emulsifiable concentrate per gallon of water) and mix thoroughly. This makes a 0.5% Lindane Spray. Apply just short of run-off to areas where these insects congregate or gather.

Roaches and Ants

Mix as above and apply the 0.5% Lindane Emulsion as a surface spray to cracks, crevices, and other places where the roaches and ants crawl or hide. Treat ant trails and areas around door sills and window frames where ants enter building. Repeat application as often as necessary for both ants and roaches.

20% Lindane is to be used in such buildings as kennel buildings and barns, therefore the animal quarters intended are kennels and barnyards.

DO NOT USE IN POULTRY HOUSES. DO NOT USE IN DAIRY BARNS AND MILK ROOMS.

OUTDOOR SPRAY (Ground Equipment)

Mosquitoes, Flies and Sand Flies (Culicoides) When Lindane is applied by fogging, rather frequent and regular repeated applications are usually necessary.

For temporary control of these insects around camp sites and bivouac areas, add 1 part of emulsifiable concentrate to 49 parts of water (2 6/8 ounces of emulsifiable concentrate per gallon of water) and mix thoroughly. This makes a 0.4 percent lindane spray. Apply liberally to vegetation, tree trunks, and surfaces other than food growing areas, at the rate of 5 to 10 gallons per acre depending on the amount of vegetation and other surfaces to be treated. To control mosquito larvae apply the spray as a fine mist at the rate of 5 gallons per acre of water surface.

Mites (Chiggers), Ticks and Fleas - In swamps and marshes. When Lindane is applied by fogging, rather frequent and regular repeated applications are usually necessary.

5/75

# 20% LINDANE

## EMULSIFIABLE CONCENTRATE

<p><b>Active Ingredients:</b>  Gamma Isomer of Benzene Hexachlorocyclopentadiene (from Lindane)  Aromatic Petroleum Distillate</p> <p><b>Inert Ingredients:</b></p>	<p><b>ACCEPTED</b></p> <p>MAY 23 1975</p> <p>Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 551-221</p>
---	---

### CAUTION

551-221

KEEP OUT OF REACH OF CHILDREN

Concentrated solution must be diluted with water as directed before use. Avoid contact with skin.

May be fatal if swallowed. Toxic when taken internally or when absorbed through the skin. In case of spillage on the skin, wash immediately with soap and water. Do not use on humans or household pets.

Avoid contaminating food or food utensils while applying sprays. Avoid breathing spray mist.

"See other cautions on side panel"

"Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed."



Manufactured by:  
**BAIRD & MCGUIRE, INC.**  
Holbrook, Massachusetts

EPA Reg. No. 551-221  
EPA Est. 551-MA-1

Add 1 part of emulsion concentrate to 49 parts of water as above. For control of mites and ticks apply as a spray to ground litter and low vegetation other than food growing areas at the rate of 10 to 20 gallons per acre depending on the amount of vegetation and litter. For flea control apply at the same rate to infested grounds and under buildings.

For Control of Birch Leaf Miners, and Holly Leaf Miners, Apply when leaves are well extended using 1 teaspoon of Lindane to 1 gallon of water.

OUTDOOR SPRAY (Airplane Application)

Mosquitoes - In swamps and marshes.

Do not treat Estuaries, Bayous, etc., valuable shrimp resources may be damaged.

Add 1 part of the emulsifiable concentrate to 3 parts of water (2 pounds emulsifiable concentrate per gallon of water). This makes a 5 percent lindane spray. Apply to areas where adult mosquitoes occur, at the rate of 1 quart per acre in wooded and heavily vegetated areas and 1 pint per acre in open or sparsely vegetated areas. For mosquito larvae, apply 1 quart per acre of water surface.

Remark Care of Equipment After use with lindane emulsion, all equipment should be carefully cleaned, dried, treated with an anti-rust agent, and stored so as to avoid corrosion.

CAUTION

Toxic to fish and wild life.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Birds feeding on treated areas may be killed.

Keep out of lakes, ponds, and streams.

Do not apply when weather conditions favor drift from areas treated.

Do not apply where runoff is likely to occur.

Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

DRUM DISPOSAL:

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

SMALLER CONTAINER DISPOSAL:

Do not reuse container, destroy by perforating or crushing and burying or discarding in a safe place.