

[illegible]

The first of these is the *Journal of the American Medical Association* (JAMA), which has been the most influential of the medical journals in the United States. It was founded in 1883 and has since then published a wide range of medical research, including clinical trials, laboratory studies, and reviews of the literature. The JAMA has been a leading voice in the medical community, and its publications have been widely cited in the medical literature.

1. *Abstracts of the 1998 Annual Meeting of the American Society of Human Genetics*. *Am J Hum Genet* 63:1-100 (1998).

actively growing. The spray is often applied when 50 percent of the tops have fallen but all tops are still green. This condition usually occurs about 10 days to two weeks before harvest time. Green tops are essential for the absorption of BAIRD'S MALEIC HYDRAZIDE. CAUTION: Avoid too early spraying of edible products. Spraying too early may result in spraying is done earlier than two weeks before maturity.

In control sprouting of potatoes in storage: Apply 1 gallon BAIRD'S MALEIC HYDRAZIDE in 30-150 gallons of water per acre for ground equipment. Airplane application can be made in irrigated sections of the country. Some soil is said to be if you have a good set of tubers. BAIRD'S MALEIC HYDRAZIDE is applied when most of the tubers set are approximately 1 inch in diameter. This occurs just as the lower leaves show a yellow color. When the potatoes flower, usually, this is 4 weeks after tubers fall off the vine, usually in May, June or July 6 weeks before harvest (see page 13, 14, 15).

For inducing dormancy in young non-bearing citrus trees in Florida: By inducing dormancy in young MALE CYPRIPED HYDRAZIDE offers a safe and effective way to damage (down to 21 percent) the recent experiments. Add 2 quarts MALE CYPRIPED HYDRAZIDE to 100 gallons water and spray with coarse coarse nozzle between September 1-15. The spray should thoroughly wet the leaf and lower leaf surfaces, branches and stems of trees. A second treatment may be necessary in October to November. Thus after dormancy is induced in young trees during the summer months, the trees will be dormant and will not produce fruit. For trees with a heavy fruit load, which are being sprayed with a coarse and mist spray, the spray should be applied to the leaves of the trees. Do not spray trees in the winter. The fruit to be harvested in the spring should be at least 24 hours before the spray is applied. The spray should be applied to the leaves of the trees 24 hours after application and the fruit should be harvested. The trees sprayed with MALE CYPRIPED HYDRAZIDE may show a slight increase in growth. The first growth of the trees may be slightly delayed. The growth of the trees should be continued growth will be delayed.

To control wild onions and wild garlic in lawns and grass along highways, 100 gallons of 2,4-D, 40% Aqueous Suspension, 20 to 100 gallons of 2,4-D, 20% Suspension, 20 to 100 gallons of water per acre, or 100 to 200 gallons of 2,4-D, 90% Concentrate, per acre, applied in late spring or early summer, will control growth of these weeds.

use of this material in a manner or at any time other than in accordance with the directions may result in plant injury or other undesirable results. Borden's McGraw-Hill makes no express or implied warranties, other than specified on this label.

ACTIVE INGREDIENTS:	
Diethanolamine salt of 1,2 Dihydro 3,6 pyridazin-1-one *	58.0 %
INERT INGREDIENTS:	42.0 %
	100.0 %

BAIRD'S Maleic Hydrazide

© 2000 Blackwell Science Ltd, *Journal of Internal Medicine* 247: 395–401

Avoid inhalation of mist. Avoid contact with skin, eyes, or clothes. May be harmful if swallowed.
Container Disposal: Dig 6 inches in top and bottom. Bury 18" below soil surface at safe disposal site. Never reuse for food or water.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of waste.

USDA Reg No 551 194

* To control tobacco sucking, apply THALIC HYDRAZIDE at the same time. Spraying too early may prevent normal development of top leaf. Spraying too late will stunt the plant growth. For further information, call 1-800-4-A-Plant or your local nursery for a list of dealers.

Revised by the 76th Congress, 2d

5. Foreign Disarmament Commission (1978-1980). "Toward a Comprehensive and Total Disarmament: The Role of the International Community," *Disarmament Yearbook*, Vol. 13, pp. 1-10.

4. MALTOL HYDRAZIDE (88-14-1) is a white crystalline solid, soluble in water, with a melting point of 150-155°C. It is used as a reagent in the synthesis of various heterocyclic compounds.

[illegible][illegible][illegible]



• **Journal of the American Medical Association**

TECHNICAL DNB
For Manufacturing Purposes Only

CONTENTS

Active Ingredient
DIFLUSEB 2,4-Dichloro-4-bromophenyl
Inert Ingredients

Net Weight 4.00
Gross Weight 4.00



Keep out of Reach of Children

ABSORBED THROUGH THE SKIN MAY BE FATAL IF SWALLOWED
Do Not Get in Eyes on Skin on Clothing Avoid Breathing Spray Mist
Do Not Wear Contaminated Clothing or Shoes
Keep Away from Heat and Open Flame
Do not store near fertilizers, seeds, insecticides or fungicides

FIRST AID TREATMENT POISON ANTIDOTE

Symptoms of Poisoning: Excessive Fatigue, Sweating, Thirst and Fever. If symptoms of poisoning develop from any type of exposure, Send For A Physician

FIRST AID: Have patient lie flat in coolest spot available. If feverish, cool with cold compresses or by immersion in cool water.

If Swallowed: SEND FOR A PHYSICIAN. Induce vomiting by giving an emetic such as 2 tablespoonful of table salt in a glass of warm water. Repeat until vomit fluid is clear, then give two teaspoonful of baking soda in a glass of warm water. Treat as in FIRST AID above.

If Splashed in Eyes: Immediately flush eyes with plenty of water for at least 15 minutes and get medical attention. If Spilled on Skin, immediately remove contaminated clothing, including shoes, and wash skin with soap and plenty of water. If symptoms of poisoning develop, send for a physician and treat as in FIRST AID above. Discard contaminated clothing and shoes or clean them thoroughly before reuse. NOTE TO ATTENDING PHYSICIAN: Active ingredient is a metabolic stimulant. Treat Symptomatically.

USE PRECAUTIONS

To avoid staining of carpets, upholstery, etc., keep contaminated clothing and shoes outdoors until cleaned. Do not contaminate water used for irrigation or domestic purposes. Dispose of empty containers by puncturing holes in them and burying with wastes.

For manufacturing purposes only.

This product is toxic to fish, birds, and other wildlife. Rinse equipment and containers with water and dispose of wastes by burying in non-crop lands away from water supplies. Keep away from farm animals.

Manufactured for:
Baird & McGuire, Inc.
South St.
Holbrook, Mass.

EPA Reg. No. 551-229
EPA EST. 551-MA-1