GENERAL DIRECTIONS

Baird's N-D Herbicide is a selective herbicide combination for the control of the following annual grasses and broadleaf weeds

ANNUAL GRASSES [illegible] Barnyardgrass, Crabgrass, Sandbur, [illegible] Wild Proso, Cupgrass and Johnsongrass [illegible]

BROADLEAF WEEDS [illegible] Purslane, Pigweed, Quickweed, Carpetweed, Morningglory, Beggarweed (Beggar Lice) and Jewelweed (Touch-Me-Not)

DO NOT USE BAIRD'S N-D HERBICIDE ON SILT LOAM SOILS OF EXTREMELY FINE TEXTURE

PREPARATION OF SOIL The area to be treated should be well prepared and as free as possible from stubble and litter. If the land has been plowed, the soil clods should be broken up by harrow or roller wheel. Plant on a flat or slightly raised bed. If weather factors make replanting necessary after Baird's N-D Herbicide has been applied, thoroughly disk the area prior to replanting.

METHOD OF APPLICATION Thorough and uniform coverage of the soil surface is essential for good weed control. Apply Baird's N-D Herbicide as a spray **at planting time, using either a planter-mounted sprayer or a separate sprayer after** the planting operation. The chemical may be applied up to the time of crop emergence or "cracking time" on peanuts but before the crop plants emerge on soybeans.

DIRECTIONS FOR USE

Put one-half the recommended amount of water in the spray tank, add the required amount of Baird's N-D Herbicide, then add the remainder of the water to the tank while mixing. Stir or use by-pass agitation until completely mixed. It is essential that the sprayer be cleaned accurately prior to use on corn.

SOYBEANS Use Baird's N-D Herbicide [illegible] overall spray (broadcast) application just after the soybeans are planted but before the soybean plants emerge from the soil.

RECOMMENDED DOSAGE RATE FOR SOYBEANS Use Baird's N-D Herbicide [illegible] 40 ounces of water per acre is an overall spray [illegible] Treatment may be applied over the entire area [illegible]

Amount of Baird's N-D Herbicide Per Acre to Use for Band Treatment of Soybeans

[illegible]	Row Spacing	[illegible]
[illegible]	[illegible]	[illegible]
[illegible]	[illegible]	[illegible]
[illegible]	[illegible]	[illegible]
[illegible]	[illegible]	[illegible]
[illegible]	[illegible]	[illegible]
[illegible]	[illegible]	[illegible]

NOTE 1 [illegible]

NOTE 2 [illegible]

NOTE 3 [illegible] pattern.

BAIRD'S N-D HERBICIDE

Controls Seedling Grasses and Broadleaf Weeds in Soybeans and Peanuts

ACTIVE INGREDIENTS	
Sodium N-1-Naphthylphthalamate*	22.3
Sodium salt of dinoseb** (2-sec-butyl-4,6-dinitrophenol)	11.5%
INERT INGREDIENTS	66.2%
Total	100.0%

*Equivalent to 20.8% N-1-naphthylphthalamic acid

**Equivalent to 10.5% 4,6-Dinitro-o-sec-butylphenol

Each gallon contains two pounds of active NPA and one pound of Active DNBP

POISON **DANGER POISON** POISON

KEEP OUT OF REACH OF CHILDREN AND FARM ANIMALS

May be fatal if swallowed or absorbed through skin
[illegible] skin or on clothing
[illegible] contaminated clothing
[illegible] the spray mist
[illegible] internally
[illegible] home

SYMPTOMS OF POISONING [illegible] symptoms [illegible] SEND FOR A PHYSICIAN.

FIRST AID [illegible] cold compresses [illegible] IF SPLASHED IN EYES, immediately flush [illegible] minutes and get medical attention. IF SPLASHED ON SKIN [illegible] and clothing, INCLUDING SHOES, [illegible] and plenty of water. If symptoms of poisoning develop, see [illegible] a physician and follow FIRST AID above. Discard contaminated clothing and shoes or clean them thoroughly before reuse. Staining may result if [illegible] articles are allowed to contact carpets, upholstery, etc.

ANTIDOTE [illegible]

[illegible] waste. [illegible] instructions.

Manufactured by
[illegible] McGUIRE INC.
Holbrook, Mass.

EPA REG. NO. [illegible]

NET CONTENTS [illegible]

ACCEPTED
Nov. 16, 1973
UNDER THE FEDERAL INSECTICIDE, FUNGICIDE, AND RODENTICIDE ACT, AS AMENDED, FOR THE PRODUCT NAMED, AND UNDER THE ESTABLISHMENT NUMBER [illegible]
551-237

RECEIVED
OCT 29 1973
Pesticides Regulation Division, EPA

NOTE 4. Baird's N-D Herbicide [illegible] best results are obtained when [illegible] mixture.

NOTE 5 May be used in the [illegible] incorporated herbicide.

NOTE 6 [illegible]

PEANUTS [illegible] This is normally from 5 to 7 da[illegible] made from planting time up to th[illegible] the "cracking stage."

RECOMMENDED DOSAGE RAT[illegible] [illegible] at the rate of 6 quarts in 20 to [illegible] (broadcast application). Treatmen[illegible] desired. The soil should not be di[illegible] treatment is used.

Amount of Ba[illegible] to Use for B[illegible]

Width of Band	[illegible]
12"	6
14"	7
Width of Band	32
12"	4
14"	5

Use in 12-20 gallons of water per [illegible]

NOTE 1 Plant on a flat or slightl[illegible]

NOTE 2 [illegible] and plant at least 2 inches deep.

NOTE 3. May be used in the [illegible] incorporated herbicide.

Do not allow spray[illegible]
Plant not recomme[illegible]

CONTAINER STORAGE AND H[illegible] [illegible] but it may crystallize [illegible] 75-125 F. [illegible] solubilizing [illegible] storing before use is recommend[illegible] fertilizers, seeds, plants and pestic[illegible]

EQUIPMENT CLEANING Thor[illegible] after application [illegible] by washing thoroughly with [illegible] water containing [illegible] detergent.

Container disposal *[illegible] and 5 gallon* [illegible] for supplies.
30 and 55 gallon containers [illegible] reconditioning [illegible] by perf[illegible]

NOTICE [illegible] may produce [illegible] on the label.