

DIRECTIONS FOR USE

HOUSEFLIES AND STABLE FLIES (In Farm Buildings other than Dairy and Dairy Barns): Use 1 gal. per 10 gal. of water and apply at the rate of 1 qt. per 500 sq. ft. of surface. Spray particularly those areas where flies crawl or rest and congregate. Repeat application as infestations warrant.

BEANS (Mexican Bean Beetles, Leafhoppers): Apply 2 qts. per acre in sufficient water to give good coverage of both upper and lower leaf surfaces. Repeat in 2 weeks if infestations warrant.

SOYBEANS, PEANUTS (Velvetbean Caterpillars): Apply 2 qts. per acre in sufficient water to give coverage. Treat at first sign of infestation and repeat in 10 days if infestation warrants.

ALFALFA and CLOVER Seed or Hay—(Alfalfa Weevils, Larvae, Alfalfa Caterpillars, Lygus Bugs, Spittle Bugs, Leafhoppers): Apply 2-3 qts. per acre in sufficient water to cover. Begin application at first sign of infestation and repeat as infestations warrant. Do not treat during bloom.

IMPORTANT DO NOT APPLY TO CROPS WITHIN 7 DAYS BEFORE HARVEST

MOSQUITO CONTROL (Non Agricultural Land Parks, Beaches, Public & Residential Areas, and dumps, Marshlands, Swamps and low lying areas) By Thermal Fogging and Mist Blowers. For Adults and Larvae: Apply 2-3 qts. per acre in sufficient water. Mix 1 part to 10 parts of water. Apply only to known mosquito breeding areas. For Thermal Fogging: 2-3 gallons per acre. Apply only to known mosquito breeding areas. For Mist Blowing: 1 part to 37 parts of water. Apply 2-3 qts. per acre. Avoid fogging when wind is from the north.

9/75

NET CONTENTS

1 5 30 55

METHOXYCHLOR 25 HAN

{25% EMULSIFIABLE SOLUTION}
CONTAINS 2 LBS. METHOXYCHLOR PER GALLON}

**FOR CONTROL OF INSECT PESTS OF
CROPS AND STORED GRAIN**

ACTIVE INGREDIENTS:

Methoxychlor*, technical 25%
Aromatic Petroleum Derivative Solvent 70%

INERT INGREDIENTS: 5%
Total 100%

*Equivalent to 22% of 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 3% of other isomers and related compounds.

CAUTION
KEEP OUT OF REACH OF CHILDREN

Avoid inhalation and contact with the skin.

Wash with soap and water.

Harmful if swallowed.

FLAMMABLE MIXTURE

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

ACCEPTED

001 11 1975

USE OF THIS PRODUCT IS LIMITED TO THE PURPOSES SPECIFIED ON THE LABEL. IT IS NOT TO BE USED FOR ANY OTHER PURPOSES. THE USER SHALL BE RESPONSIBLE FOR THE PROPER USE OF THIS PRODUCT.

DIRECTIONS FOR USE

MIST BLOWERS (For use in stored grain): Apply 1 qt. per 500 sq. ft. of surface. Spray particularly those areas where flies crawl or rest and congregate. Repeat application as infestations warrant.

For all types of application, thoroughness is essential for effective control.

ELM BARK BEETLES

How to Use

Hydraulic Sprayers:

Apply 2-3 qts. per acre in sufficient water to cover.

Mist Blowers:

Apply 1 qt. per 500 sq. ft. of surface.

When to Use

Apply at first sign of infestation and repeat as infestations warrant.

Do not treat during bloom.

Do not treat when wind is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

Do not treat when temperature is below 50°F.

Do not treat when humidity is below 50%.

Do not treat when precipitation is expected.

Do not treat when wind speed is above 10 mph.

Do not treat when wind direction is from the north.

NET CONTENT

5

30

55

DIRECTIONS FOR USE

For use on all crops and stored grain. Use as directed on label. Do not use on crops or stored grain which are to be used for human consumption.

For use on all crops and stored grain.

For use on all crops and stored grain. Use as directed on label. Do not use on crops or stored grain which are to be used for human consumption.

For use on all crops and stored grain. Use as directed on label. Do not use on crops or stored grain which are to be used for human consumption.

For use on all crops and stored grain. Use as directed on label. Do not use on crops or stored grain which are to be used for human consumption.

METHOXYCHLOR 25 HAN

{ 25% EMULSIFIABLE SOLUTION }
{ CONTAINS 2 LBS. METHOXYCHLOR PER GALLON }

FOR CONTROL OF INSECT PESTS OF
CROPS AND STORED GRAIN

ACTIVE INGREDIENTS

Methoxychlor, technical

Aromatic Petroleum Derivative

INERT INGREDIENTS

Total

20

70

5

100

CAUTION

KEEP OUT OF REACH OF CHILDREN

DO NOT USE IN FOOD