

PRECAUTIONARY STATEMENTS

Hazards to humans and domestic animals

CAUTION: May be irritating to eyes, nose, throat and skin. Flush area of irritation with plenty of water. May be harmful if swallowed. Do not eat, drink, or smoke while using this product. Wash hands thoroughly after use.

Environmental Hazards

Keep out of water. Do not discharge into water. Do not contaminate water with wash water or empty containers.

DIRECTIONS FOR USE

This material is a Federal Law Pesticide. It is to be used in a manner consistent with the label.

To Control Sprouting of Onions in Storage: Apply 1 gallon Baird's Super Maleic Hydrazide in 100-150 gallons of water per acre with a ground sprayer. Airplane application is also effective. The spray should be applied when the bulbs are fully mature, have five to six green leaves, and are soft enough to be topped to fall if they have not already done so. The spray is effective when 50 percent of the tops are still green. The condition usually occurs about 10 weeks to 10 days before harvest time. Super Maleic Hydrazide is effective for the absorption of Baird's Super Maleic Hydrazide.

CAUTION: Avoid too early spraying of onions. Sprouting bulbs may be delayed in time earlier than they were before spraying.

To Control Sprouting Potatoes in Storage: Apply 1 gallon Baird's Super Maleic Hydrazide in 100-150 gallons of water per acre with a ground sprayer. Airplane application is also effective. Check your lot to see if you have a lot of potatoes. Baird's Super Maleic Hydrazide is applied when the potatoes are set and are approximately 1 1/2 inches in diameter. This is usually the time when leaves show a yellowish color and potatoes have begun to grow. This is usually after blossoming in the field. August about 4-6 weeks before harvest. See notes, label, 20.

To Control Wild Onions and Wild Garlic in Lawns and Grass Areas Along Highways (not pasture or cropland): Apply 1.1 to 2.3 gallons Baird's Super Maleic Hydrazide in 100 gallons of water per acre in spray or fall when 80-90 percent of the foliage has started to grow with new leaves. When grass is green and dormant. Do not use when leaves are dying or turning brown.

To Inhibit Grass Growth: Apply 1 gallon Baird's Super Maleic Hydrazide in 100 gallons of water per acre to green grass. Since brown grass will not respond to spray. One application per year is either in the fall or the spring followed by correct maintenance. To reduce mowing, use 1/2 gallon per acre.

Spring Application (About April 10-June 1): Use 1.1 to 4 gallons Baird's Super Maleic Hydrazide in 50 gallons of water per acre to reduce or eliminate mowing.

Fall Application (About Oct 1 - Nov 15): Use 2.1 to 4 gallons Baird's Super Maleic Hydrazide in 50 gallons of water per acre to reduce growth next season. The treatment is applied to green grass. It becomes dormant. Do not apply during the summer (June 1-Sept 1) since most permanent grasses become partially dormant in summer.

Use of 2,4-D

2,4-D is a herbicide that is used to control broadleaf weeds in corn, soybeans, and other crops. It is also used to control weeds in pastures and lawns. 2,4-D is a systemic herbicide, meaning it is absorbed by the plant and moves to the site of action. It is effective against a wide range of weeds, including clover, alfalfa, and other broadleaf species. 2,4-D is also used to control weeds in ornamental plants and trees.

For Inducing Dormancy in Young Non-Bearing Citrus Trees in Florida

Apply 1 gallon Baird's Super Maleic Hydrazide in 100 gallons of water per acre to young non-bearing citrus trees in Florida. This treatment is applied to the foliage of the trees and is effective in inducing dormancy, which helps to reduce the risk of frost damage and other environmental stresses.

Apply 1 gallon Baird's Super Maleic Hydrazide in 100 gallons of water per acre to young non-bearing citrus trees in Florida. This treatment is applied to the foliage of the trees and is effective in inducing dormancy, which helps to reduce the risk of frost damage and other environmental stresses.

NOTES

1. For best results, apply this product in the early morning or late afternoon when temperatures are between 60°F and 80°F. Do not apply when temperatures are above 90°F or below 50°F. 2. Do not apply this product to wet foliage. 3. Do not apply this product to plants that are stressed or diseased. 4. Baird's Super Maleic Hydrazide is a systemic herbicide. It is absorbed by the plant and moves to the site of action. 5. Do not use this product on crops that are sensitive to herbicides. 6. Do not use this product on crops that are growing in water. 7. Do not use this product on crops that are growing in soil that is contaminated with herbicides. 8. Do not use this product on crops that are growing in soil that is contaminated with other pesticides. 9. Do not use this product on crops that are growing in soil that is contaminated with other chemicals. 10. Do not use this product on crops that are growing in soil that is contaminated with other substances.

WARRANTY AND LIMITATION OF DAMAGES

Baird & McGuire, Inc. warrants that this product is free from defects in material and workmanship. It also warrants that this product is effective against the weeds and grasses listed on the label. However, Baird & McGuire, Inc. does not warrant that this product will be effective against all weeds and grasses. The user assumes all responsibility for the results of the use of this product. Baird & McGuire, Inc. is not liable for any damages, including consequential damages, arising from the use of this product. The user agrees to hold Baird & McGuire, Inc. harmless from and against all claims, damages, and expenses, including attorney's fees, arising from the use of this product.

Baird's Super Maleic Hydrazide

Liquid Growth Retardant for Preventing the Sprouting of Potatoes & Onions.


ACTIVE INGREDIENT

Potassium salt of 6-hydroxy-2-naphthyl pyridazinone

INERT INGREDIENTS

Tar

See label for full list of ingredients. The total weight of the active and inert ingredients is 100 percent.



KEEP OUT OF THE REACH OF CHILDREN

CAUTION

Statement of Practical Treatment

May be irritating to eyes, nose, throat and skin. Flush area of irritation with plenty of water.

See Side Panel For Additional Precautionary Statements

EPA Reg. No. 551-276
EPA Est. 551-MA-1

NET CONTENTS

STORAGE AND DISPOSAL

Store in a cool, dry place. Do not store in a metal container.

PESTICIDE DISPOSAL

Do not use this product in a way that will result in the release of hazardous substances into the environment. This product must be disposed of in accordance with Federal, State, and local laws and regulations. Do not use this product in a way that will result in the release of hazardous substances into the environment.

CONTAINER DISPOSAL

Metal containers must be recycled or disposed of in accordance with local laws and regulations. Do not use this product in a way that will result in the release of hazardous substances into the environment.

Baird & McGuire, Inc.
Holbrook, MA 02343