DIRECTIONS

This is a selective herbicide used for the control of susceptible weeds in small grains and corn, and for control of certain deep rooted perennials and woody plants in non-crap lands. For best results, apply when soil and climatic conditions are conducive for rapid wend growth.

WEEDS CONTROLLED: Mustard, Frenchweed, Arrowhead, Coddebur, Dandellon, Gumweed, Marsh Elder, Hedge Bindweed, Plantain, Ragweed, Field Horsetail, Plantain and other broadled weeds.

WHEAT, BARLEY, RYE - Use 1/2 to 1 pint of this herbicide per acre in sufficient water to obtain good coverage. It is recommended that the lawer doeage be used for more susceptible weeds such as common mustard, and the higher doeage level may be necessary for the control of more resistant weeds such as Carada thistle. Make application only between the stooling and joint stages.

NOTICE: Do not forage or graze on treated grain fields within two (2) weeks after treatment. Do not feed treated straw to livestock.

CORN - Use ½ to 1 pint of this herbicide per acre in sufficient water to obtain good coverage. Make application following emergence of majority of weeds but before weeds have damaged the corn. Post emergence application of this herbicide should be made after emergence but before tasseling. Do not apply from tasseling to dough stage. Use drop nazzles when corn is over 8 inches high. Strains and varieties of corn vary in their tolerance to 2,4-D and particular care should be exercised in using either amine or ester formulations.

DEEP ROOTED PERENNIALS AND WOODY PLANTS - In areas such as ditch banks, highway right of ways and fence rows, use 1½ to 3 pints of this herbicide per acre in sufficient water to insure good coverage. In the control of plants such as willows and poison oak, it is necessary to have thorough coverage. Certain weeds and woody plants will form new growth after treatment and repeat applications may be necessary. For best results apply in spring after new leaves appear and plants are growing rapidly.

NET CONTENTS 5 GALLONS

AGSCO 2,4-D AMINE WEED KILLER

ACTIVE INGREDIENTS

* Dimethylamine salt of 2,4 Dichterophenoxyacetic acid INERT INGREDIENTS:

50.6 **%** 100.0 **%**

* 2,4-Dichlorophenoxyacetic acid equivalent 41.0%
Contains 4 Pounds 2,4-D Acid Per Gallon

CAUTION KEEP OUT OF REACH OF CHILDREN

(Additional precautionary statements on side panel)

Harmful if swallowed. Do not get in eyes. Avoid contact with skin or clothing. In case of contact with weed killer, flush with plenty of water for fifteen (15) minutes. For eyes, get medical attention. Remove and wash contaminated clothing and shoes before re-use.

Protect from temperatures below 0°F.

Mary factured By

AGSCO CHEMICALS, INC.

Grand Forks, North Dakota

WARNING

Avoid inhaling spray,

Applications by air, ground or hand equipment should be made only when there is no hazard from drift. Spray drift may cause injury to susceptible plants such as vegetables, flowers, grupes or other plants sensitive to minute quantities of herbicide. To avoid injury to desirable plants, do not apply other agricultural chemicals with the same equipment used for this herbicide. Do not apply by air in the vicinity of sugar beets.

Do not store near fertilizer, seeds, insecticides, fungicides, feed or foodstuffs.

Excess amounts of this product in soil may temporarily inhibit seed germination or plant growth.

Do not contaminate any body of water by direct application, cleaning of equipment or disposal of waste and containers.

Destroy empty container by perforating and bury in a safe place. Do not burn.

Avoid excess use in environment of desirable animals and birds. Toxic to fish

Local conditions may affect the use of herbicides. State agricultural authorities in many states issue recommendations to fit local conditions. Be sure that the use of this product conforms to all applicable conditions.

WARRANIY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under renormal conditions, or under conditions not reasonably for eseeable to seller, and buyer assumes the risk of any such use.

11/17/67

USDA Registration Number 554-54