

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**WARNING**

Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with skin, eyes or clothing. Causes skin irritation. Avoid breathing spray mist. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

**ENVIRONMENTAL HAZARDS**

Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply directly to water. Do not apply when weather conditions favor drift from target area, as this product may injure cotton, beans, other vegetables, certain legumes and ornamentals. Toxic to fish. Excessive amount of this product in soil may temporarily inhibit seed germination or plant growth.

**STORAGE AND DISPOSAL**

**STORAGE:**

DO NOT contaminate water, food or feed by storage or disposal, or cleaning of equipment. DO NOT store near fertilizer, other pesticides or seeds. Store at a temperature above 0 degrees Fahrenheit. If allowed to freeze, remix before use. **FLAMMABLE**—Keep away from heat or open flame.

**PESTICIDE DISPOSAL:**

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:**

TRIPLE RINSE (or equivalent) or dispose in approved landfill. DO NOT BURN. Consult Federal, State, or Local Disposal Authorities for approved alternate procedures. Do not reuse empty container.

**AGSCO  
400  
HERBICIDE**

**A LOW VOLATILE FORMULATION**

ACTIVE INGREDIENT:

\*Isooctyl ester of 2-4-Dichlorophenoxyacetic acid ..... 71.26%

INERT INGREDIENTS ..... 28.74%

\*2-4-Dichlorophenoxyacetic acid equivalent 40.6%

Contains 4 Pounds Acid Equivalent Per Gallon

ACCEPTED  
NOV 24 1987  
554-83

**KEEP OUT OF REACH OF CHILDREN**

**WARNING**

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Do not induce vomiting. If vomiting occurs spontaneously, keep airway clear. Do not induce vomiting or give anything to an unconscious. Consult a physician.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**IF ON EYES:** Flush with plenty of water. Get medical attention if irritation persists.

EPA EST. NO. 554-ND-1

EPA REG. NO. 554-83

NET CONTENTS: 2.5 GALLONS

Manufactured By: AGSCO, INC. Grand Forks, ND

copy  
made  
1/2

554-83

11/24/1987

1/2

### ENDANGERED SPECIES RESTRICTIONS

The following restrictions apply to use of this product after February 1, 1988.

"Before using this pesticide on [corn, wheat, barley, rye] in the counties listed below, you must obtain the PESTICIDE USE BULLETIN FOR PROTECTION OF ENDANGERED SPECIES FOR THE COUNTY IN WHICH THE PRODUCT IS TO BE USED. The bulletin is available from your County Extension Agent, State Fish and Game Office, or your pesticide dealer. **Use of this product in a manner inconsistent with the PESTICIDE USE BULLETIN FOR PROTECTION OF ENDANGERED SPECIES is a violation of Federal laws.**"

ALABAMA: Colbert, Greene, Jackson, Lamar, Lauderdale, Limestone, Madison, Marshall, Morgan, Pickens and Sumter

ARKANSAS: Clay, Clark, Cross, Lawrence, Lee, Poinsette, Randolph, Sharp and St. Francis

CALIFORNIA: Solano

FLORIDA: Broward, Dade, Glades and Palm Beach

KENTUCKY: Ballard, Butler, Emundson, Green, Hart, Jackson, Laurel, Livingston, Marshall, McCracken, McCreary, Pulaski, Rockcastle, Taylor, Warren, and Wayne

MISSISSIPPI: Itawamba, Lowndes, Monroe and Noxubee

NORTH CAROLINA: Edgecombe, Nash and Pitt

TENNESSEE: Bedford, Blount, Claiborne, Decatur, Franklin, Hancock, Hardin, Hickman, Knox, Lincoln, Loudon, Marshall, Maury, Meigs, Monroe, Rhea, Roane, Scott, Sequatchie, Smith, Sullivan and Wayne

VIRGINIA: Lee, Russell, Scott, Smyth, Tazewell, Washington and Wise

### WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**WEEDS CONTROLLED:** Mustard, Frenchweed, Arrowhead, Cocklebur, Dandelion, Gumweed, Marsh Elder, Hedge Bindweed, Pigweed, Ragweed, Wild Buckwheat, Plantain, and other broadleaf weeds.

**WHEAT, BARLEY, RYE:** Use 1/2 to 1 pint of this herbicide per acre in 3-5 gallons of water by airplane application or 5 gallons of water by ground applicator. Make application only between stouling and joint stages. Do not spray from boot to dough stage.

**NOTICE:** Do not forage or graze on treated fields within two (2) weeks after treatment. Do not feed treated straw to livestock.

**CORN:** Use 1/2 to 1 pint of herbicide per acre in 5 gallons of water. Make application following emergence of majority of weeds but before weeds have damaged the corn. Post emergence application of this herbicide should be made after emergence but before tasseling. Do not apply from tasseling to dough stage. Use drop nozzles when corn is over 8 inches high. Strains and varieties of corn vary in their tolerance to 2, 4-D and particular care should be exercised in using ester formulations.

**DEEP ROOTED PERENNIALS AND WOODY PLANTS:** in areas such as highway right of ways and fence rows, use 1 1/2 to 3 pints of this herbicide per acre in 5 gallons of water by ground applicator. In the control of plants such as Willows and Poison Oak, it is necessary to have thorough coverage. Certain weeds and woody plants will form new growth after treatment and repeat applications may be necessary.

**RED POTATOES:** Use 1/8 pint of this herbicide per acre in 5-25 gallons of water by ground applicators. To intensify red color and improve skin appearance apply when potatoes are in pre-bud stage (7-10 inches high); second application 10-14 days later.

Avoid use of small-diameter spray nozzles. Avoid spray drifts onto susceptible plants such as vegetables, flowers, tomatoes, beans, sugarbeets, sunflowers and other legumes. Coarse sprays are less likely to drift. **DO NOT** spray at all in the vicinity near susceptible plants to prevent any vapor drift. Local conditions may affect the use of herbicides. State agricultural authorities in many states issue recommendations to fit local conditions. Be sure that the use of this product conforms to all applicable conditions.

LABEL CODE 1019 87