DIRECTIONS

In preparing spray mixtures add at least half of the water to the spray tank before adding this herbicide. Apply at moderate pressure by air or ground equipment during warm weather when plants are growing actively.

Ground application: Apply at the rate of 5-10 gallons of diluted herbicide per acre.

Aircraft application: Apply at the rate of 3-5 gallons of dituted herbicide per acre.

Barley) — For control of mustard, lambsquarter and other susceptible weeds use 1/3 pint of this herbicide in sufficient water for coverage. Apply at any time from emergence to the boot stage of the grain. With certain weed species such as Canada thistle that are more difficult to control and under conditions where easily controlled weeds are more resistant as much as 1 pint of herbicide may be needed for effective control. If mustard is the only problem and if it is in a susceptible stage of growth, as little as 1/4 pint may be effective. Do not use on small grains interplanted with sweet clover unless some injury to the legume can be tolerated. Use on grain underseeded with alfalfa only when specifically recommended by local agricultural experiment station or extension service weed specialists.

WEED CONTROL IN FLAX: For control of mustard and other susceptible weeds use $\frac{1}{3}$ pint per acre in the necessary amount of water required for coverage. Treat flax when 2 to 6 inches tall and before the bud begins to form.

NOTICE: Do not forage or graze on treated grain fields within two (2) weeks after treatment. Do not feed treated straw to livestock.

NET CONTENTS

GALLONS



AGSCO MA HERBICIDE

ACTIVE INGREDIENT:

*Diethanolamine Salt of 2-Methyl-4-chlorophenoxyacetic acid 65.5% INERT INGREDIENTS: 34.5%

100.0%

*Acid equivalent 42.75%

Contains 4 Pounds Acid per Gallon

WARNING KEEP OUT OF REACH OF CHILDREN

(Additional precautionary statements on side panel)

Harmful if swallowed. Severe eye irritant. Do not get in eyes. Avoid contact with skin or clothing. In case of contact with weed killer, flush with plenty of water for fifteen (15) minutes. For eyes, get medical attention. Remove and wash contaminated clothing and shoes before re-use.

Protect from temperatures below 0° F.

Manufactured By

AGSCO CHEMICALS, INC. Grand Forks, North Dakota

WARNING

Avoid inhaling spray.

Coarse sprays are less likely to drift.

Applications by air, ground or hand equipment should be made only when there is no hazard from drift. Spray drift may cause injury to susceptible plants such as vegetables, flowers, grapes or other plants sensitive to minute quantities of herbicide. To avoid injury to desirable plants do not apply other agricultural chemicals with the same equipment used for this herbicide. Do not apply by air in the vicinity of sugar beets.

Do not store near fertilizer, seeds, insecticides, fungicides, feed or foodstuffs.

Local conditions may affect the use of herbicides. State agricultural authorities in many states issue recommendations to fit local conditions. Be sure that the use of this product conforms to all applicable conditions.

Excess amounts of this product in soil may temporarily inhibit seed germination or plant growth.

ENVIRONMENTAL AND WILDLIFE WARNING: Do not contaminate any body of water by direct application, cleaning of equipment or disposal of waste and containers.

Destroy empty container by perforating and bury in a safe place.

Toxic to fish.

WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability of fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

51771

USDA Registration Number 554-123

