

Reg # 554-125

PM-23

1 9 2

AGSCO MXL HERBICIDE

A LOW VOLATILE FORMULATION

INGREDIENTS

ACTIVE INGREDIENTS:

*2-Methyl-4-Chlorophenoxyacetic acid, Isooctyl ester 74.93%

INERT INGREDIENTS: 25.07%

TOTAL 100.00%

*Equivalent to 47.83% 2-Methyl-4-Chlorophenoxyacetic acid.
Contains 4 pounds Acid Equivalent per gallon.

KEEP OUT OF REACH OF CHILDREN WARNING STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything to an unconscious person.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

See Side Panel for Additional Precautionary Statements.

EPA EST. NO. 554-ND-1
EPA REG. NO. 554-125

NET CONTENTS: 2.5 GALLONS

Manufactured By: AGSCO, INC., Grand Forks, ND

DATE: NOV 7 1992
LABEL IN PER PR NOTICE 6-3

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with skin, eyes or clothing. Causes skin irritation. Avoid breathing spray mist. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

PROTECTIVE CLOTHING: Mixers/loaders or applicators are required to use face shields or goggles.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters or rinsate.

GROUND WATER CONTAMINATION: Most cases of groundwater contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCPA pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

CLEANING OF EQUIPMENT: When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: DO NOT contaminate water, food or feed by storage or disposal, or cleaning of equipment. Open dumping is prohibited. DO NOT reuse empty container. DO NOT store near fertilizer, other pesticides or seeds. Store at a temperature above 0°F. If allowed to freeze, remix before use.

FLAMMABLE — Keep away from heat or open flame.

PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Or if this is an "OUT-N-BACK" returnable container, it must be returned to AGSCO, INC.



WARRANTY LIMITATIONS, DISCLAIMER AND LIMITATION OF REMEDY

AGSCO, INC. warrants that this product (a) conforms to the chemical description on the label and (b) is reasonably fit for the purpose stated on the label when used in strict accordance with the directions therein under normal growing conditions. This is the only warranty made on this product. The manufacturer neither makes nor intends, nor does it authorize any agent or representative to make any other warranties, express or implied, and expressly includes all implied warranties of merchantability or fitness for a particular purpose.

Buyer or user's exclusive remedy and manufacturer's or seller's exclusive liability for any and all claims, losses, damage, or injuries resulting from the use or handling of this product, whether or not based on contract, negligence, strict liability in tort or otherwise shall be limited to one of the following, at the election of the seller:

- (1) Refund of the purchase price paid by the buyer or user for the product bought and used with respect to the claim for which damages are sought, or
- (2) Replacement of the amount of product used incident to the crop for which damages are sought.

The seller will not be liable for consequential or incidental damages or losses resulting from the use or handling of this product.

The terms of this limited warranty and remedy cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the seller is authorized to vary or exceed the terms of the limitations stated above.

DIRECTIONS FOR USE

WHEAT, OATS, BARLEY: Use 1 pint per acre in 5 gallons of water by ground application or in 3-5 gallons of water by airplane application. Apply anytime from emergence to boot stage of grain. DO NOT spray from boot to dough stage. DO NOT use on grains interplanted with sweet clover unless injury to the legume can be tolerated. Use on grain underseeded with Alfalfa where specifically recommended by local agricultural extension station and extension service weed specialists. For Canada Thistle, use 1 1/3 pint per acre rate.

DO NOT forage or graze meat animals on treated areas for seven days of slaughter.

FLAX: Use 1/2 pint per acre in 5 gallons of water by ground application or in 3-5 gallons of water by airplane application. Apply when weeds are up and when Flax is 2 to 6 inches high and before it comes into bud. Treatment after early bud stage may result in severe damage. If Canada Thistle is present it may be necessary to go as high as 3/4 pint per acre. Some injury to Flax may result.

Avoid use of small-diameter spray nozzles. Avoid spray drift to susceptible plants such as, vegetables, flowers, tomatoes, lettuce, sugarbeets, sunflowers and other legumes. Coarse sprays are more likely to drift. DO NOT spray at all in the vicinity near susceptible plants to prevent any vapor drift.

Susceptible weeds controlled are: Lambsquarters, Marshmallow, Annual Mustards, Ragweed, Field Pennycress, Yellow Rattle, Cocklebur, Shepherdspurse and many others. Less susceptible weeds controlled (may have more erratic results) are: Canada Thistle, Dandelions, Kochia, Plantains, Sunflowers, Pigweed, Purslane, Wild Buckwheat, Hempnettle, Stinging Nettle and others.