

ACCEPTED

7-26-71
557-1445

UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE AND FUNGICIDE MIXTURES
FOR ECONOMIC PEST CONTROL
REGISTERED UNDER NO. 557-1445
TO AGRICULTURAL CHEMICALS

VIGORO.

SNAIL AND SLUG KILLER

LIQUID FORM FOR EASY APPLICATION

LONG
LASTING
NOT
AFFECTED
BY WATER

ACTIVE INGREDIENT
Metaldehyde 12.5%
INERT INGREDIENTS 87.5%
Total 100.0%

WARNING: Keep Out of Reach of Children

NET 16 FL OZ (1 PT)

VIGORO, SNAIL AND SLUG KILLER

DIRECTIONS FOR USE: Lawns (including Dichondra), Ornamentals, (Roses, Dahlias, Gladiolus, Iris, Begonias, Pansies, Petunias, Primulas, Salvia, Strips and Ground Covers)

Area or areas to be treated should be watered thoroughly just prior to application of Vigoro Snail & Slug Killer.

SHAKE WELL BEFORE USING! Mix two tablespoonfuls per gallon of water and apply to 100 sq. ft. If Snails are an extreme problem, the amount used may be increased up to four tablespoonfuls per gallon of water. Vigoro Snail & Slug Killer will not burn foliage regardless of amount applied.

For ease of application, use hose-on type applicator. Use two tablespoonfuls for each gallon of mixture delivered by your applicator and apply at the rate of one gallon to each 100 sq. ft. of area to be treated.

GENERAL INFORMATION: Vigoro Snail & Slug Killer is especially designed for use in areas difficult to treat with dry baits or where snails and slugs can remain well hidden. It kills snails, slugs, and eggs by contact. It also kills snails and slugs that travel over treated areas.

Area or areas to be treated should be watered thoroughly just prior to application. **DO NOT APPLY ON FOOD OR FEED CROPS.**

WARNING: May be fatal if swallowed. Avoid contact with eyes and clothing. Avoid breathing spray mist. Avoid skin contact. In case of contact, wash immediately with soap and water. Avoid contamination of food and feedstuffs. Do not allow children or pets to play on treated area until residues have been washed off and into the ground. Protect contents from freezing. Do not store in direct sunlight. **DO NOT APPLY ON FOOD OR FEED CROPS.**

ANTIDOTE: Give a tablespoonful of salt in glass of warm water and repeat until vomit fluid is clear. Then give two tablespoonfuls of Epsom salts in water. Have patient lie down and keep warm. Give strong tea, coffee, or one teaspoonful of aromatic spirits of ammonia in water. **CALL A PHYSICIAN IMMEDIATELY!**

This product is toxic to wildlife. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

MANUFACTURED BY
SWIFT AGRICULTURAL CHEMICALS CORPORATION, CHICAGO, ILLINOIS 60606
U.S. P.A. REG. NO. 557-1445