

VIGORO

VIGORO CHICKWEED, CLOVER, AND POA ANNUA KILLER is an aqueous endothal solution for use as a selective spray for lawns to control Veronica (speedwell), knotweed, bur clover, black medic, chickweed (in the South), cranesbill and filaree, dichondra (pony foot), goat head, henbit, lespedeza, oxalis, vetch, little barley, rescue grass, cheat and other annual bromes, ryegrass, and for suppression of Poa annua in bent and blue grass turf. For control of certain other broadleaf weeds, 2,4 D in accordance with manufacturer's directions may be added to the sprayer with Vigoro Chickweed, Clover, & Poa annua Killer.

CONCENTRATED--1 PINT COVERS 10,000 Sq. Ft.

HOW TO MIX AND APPLY: Mix 1½ ounces (3 tablespoonfuls) of Vigoro Chickweed, Clover, & Poa annua Killer in 1 to 2 gallons of water and apply evenly to 1,000 square feet. For best results, make 2 applications at 2-3 week intervals. Apply only when moisture and temperature conditions are favorable for good turf growth. For Poa annua control, use 3 early spring applications at 10 to 14 day intervals at ¼ ounce (1½ tablespoonfuls) in 1 to 2 gallons of water and apply to 1,000 square feet. Apply evenly as a fine mist during the first 30 to 40 days of the spring growth period. Apply first application after appearance of Poa annua and before it forms seed heads. Do not use more than recommended amounts.

WARNING: May be fatal if swallowed. Irritating to skin, nose, eyes, and throat. May be absorbed through skin. Avoid breathing spray mist. Do not get in eyes, on skin, or on clothing. In case of contact, immediately flush skin or eyes with plenty of water for at least 15 minutes. For eyes, get medical attention. If swallowed, give cold milk or cold water. Call a doctor immediately. Store out of the reach of children.

CAUTION: Under certain conditions, some temporary turf browning may occur to St. Augustine, Bermuda, and other turf grasses, especially when temperatures are over 80° F. and the weather is dry. Apply only on well established vigorously growing turf. Do not use on putting greens or red fescue turfs. Do not store this material where it might contaminate seeds, feed, or foodstuffs. Avoid contact with or drift to trees or other plants as injury may occur. To avoid drift, do not apply in windy weather. Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

This product is toxic to wildlife. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Distributed by SWIFT AGRICULTURAL CHEMICALS CORP., Chicago, Ill.

©Reg. Trademark of Pennsalt Chemicals Corporation

U.S.D.A. Reg. No. 557-1611

WARNING:

CHICKWEED CLOVER & POA ANNUA KILLER

ACTIVE INGREDIENT

Disodium salt of endothal* 19%

INERT INGREDIENT 81%

Total 100%

*7-Oxabicyclo (2.2.1) heptane-2, 3-dicarboxylic acid U. S. Patents 2,550,494; 2,576,080; 2,576,081, 2,576,083; and other pat. pending.

ENDOTHAL®

AC
7-13-71
557-1611

NET CONTENTS
1 FLUID PINT

CHICKWEED CLOVER & POA ANNUA KILLER

Contains
ENDOTHAL®

ACTIVE INGREDIENT
Dipotassium salt of endothal * 10%
INERT INGREDIENT 81%
Total 100%
*7 Oxabicyclo (2,2,1) heptane-2,3-
dicarboxylic acid U. S. Patents
2,550,494; 2,576,080; 2,576,081,
2,576,083; and other pat. pending.

VIGORO CHICKWEED, CLOVER, AND POA ANNUA KILLER is an aqueous endothal solution for use as a selective spray for lawns to control Veronica (speedwell), knotweed, bur clover, black medic, chickweed (in the South), cranesbill and filoree, dichondra (pony foot), goat head, henbit, lespedeza, oxalis, vetch, little barley, rescue grass, cheat and other annual bromes, ryegrass, and for suppression of Poa annua in bent and blue grass turf. For control of certain other broad-leaf weeds, 2,4-D in accordance with manufacturer's directions may be added to the sprayer with Vigoro Chickweed, Clover, & Poa annua Killer.

CONCENTRATED 1 PINT COVERS 10,000 Sq. Ft.

HOW TO MIX AND APPLY: Mix 1 ounce, 3 tablespoons of Vigoro Chickweed, Clover & Poa annua Killer in 1 to 2 gallons of water and apply evenly to 1,000 square feet. For best results, make 2 applications at 2-3 week intervals. Apply only when moisture and temperature conditions are favorable for good turf growth. For Poa annua control, use 3 early spring applications at 10 to 14 day intervals at a rate of 1 1/2 tablespoonfuls in 1 to 2 gallons of water and apply to 1,000 square feet. Apply evenly as a fine mist during the first 30 to 40 days of the spring growth period. Apply first application after appearance of Poa annua and before it forms seed heads. Do not use more than recommended amounts.

May be fatal if swallowed or falling on skin, nose, eyes, and throat. May be absorbed through skin. Avoid breathing any mist. Do not get in eyes, on skin, or on clothing. In case of contact, immediately flush skin or eyes with plenty of water for at least 15 minutes. For eyes, get medical attention immediately. Wash with soap and milk or cold water. Call a doctor immediately. Show label of this material to children.

Under certain conditions, some temporary turf browning may occur to St. Augustine, Bermuda, and other turf grasses, especially when temperatures are over 80° F. and the weather is dry. Apply only on well established vigorously growing turf. Do not use on putting greens or red fescue turfs. Do not store this material where it might contaminate seeds, feed, or foodstuffs. Avoid contact with or drift to trees or other plants as injury may occur. To avoid drift, do not apply in windy weather. Do not reuse empty container. Destroy by perforating or crushing. Bury or discard in a safe place.

This product is toxic to wildlife. Do not contaminate water by leaking or equipment or disposal of wastes.

Distributed by SWIFT AGRICULTURAL CHEMICALS CORP., Chicago, Ill.

®Reg. Trademark of Pennsalt Chemicals Corporation

U.S.D.A. Reg. No. 557-1611

7-13-71
557-164

VIGORO

CHICKWEED CLOVER & POA ANNUA KILLER

WARNING: Keep out of reach of children. See panel for additional warnings and cautions.

ACTIVE INGREDIENT	
Disodium salt of endothall*	19%
INERT INGREDIENT	81%
Total	100%

*7-Oxabicyclo (2.2.1) heptane-2, 3-dicarboxylic acid U. S. Patents 2,550,494; 2,576,080; 2,576,081; 2,576,083; and other pat. pending.

VIGORO CHICKWEED, CLOVER, AND POA ANNUA KILLER is an aqueous endothal solution for use as a selective spray for lawns to control Veronica (speedwell), knotweed, bur clover, black medic, chickweed (in the South), cranesbill and filaree, dichondra (pony foot), goat head, henbit, lespedeza, oxalis, vetch, little barley, rescue grass, cheat and other annual bromes, ryegrass, and for suppression of Poa annua in bent and blue grass turf. For control of certain other broadleaf weeds, 2,4-D in accordance with manufacturer's directions may be added to the sprayer with Vigoro Chickweed, Clover, & Poa annua Killer.

CONCENTRATED—1 PINT COVERS 10,000 Sq. Ft.

HOW TO MIX AND APPLY: Mix 1½ ounces (3 tablespoonfuls) of Vigoro Chickweed, Clover, & Poa annua Killer in 1 to 2 gallons of water and apply evenly to 1,000 square feet. For best results, make 2 applications at 2-3 week intervals. Apply only when moisture and temperature conditions are favorable for good turf growth. For Poa annua control, use 3 early spring applications at 10 to 14-day intervals at ¾ ounce (1½ tablespoonfuls) in 1 to 2 gallons of water and apply to 1,000 square feet. Apply evenly as a fine mist during the first 30 to 40 days of the spring growth period. Apply first application after appearance of Poa annua and before it forms seed heads. Do not use more than recommended amounts.

WARNING: May be fatal if swallowed—irritating to skin, nose, eyes, and throat. May be absorbed through skin. Avoid breathing spray mist. Do not get in eyes, on skin, or on clothing. In case of contact, immediately flush skin or eyes with plenty of water for at least 15 minutes. For eyes, get medical attention. If swallowed, give cold milk or cold water. Call a doctor immediately. Store out of the reach of children.

CAUTION: Under certain conditions, some temporary turf browning may occur to St. Augustine, Bermuda, and other turf grasses, especially when temperatures are over 80° F. and the weather is dry. Apply only on well established vigorously growing turf. Do not use on putting greens or red fescue turfs. Do not store this material where it might contaminate seeds, feed, or foodstuffs. Avoid contact with or drift to trees or other plants as injury may occur. To avoid drift, do not apply in windy weather. Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

This product is toxic to wildlife. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Distributed by SWIFT AGRICULTURAL CHEMICALS CORP., Chicago, Ill.

©Reg. Trademark of Pennsalt Chemicals Corporation

U.S.D.A. Reg. No. 557-1611

