

VIGORO

5% SEVIN DUST

FOR USE ON
**VEGETABLES
 LAWNS·SHRUBS
 DOGS·CATS**

NET WT. 16 OZ. (1 LB.)

ACTIVE INGREDIENTS:
 Carbaryl (1-Naphthyl
 N-methylcarbamate) 5%
INERT INGREDIENTS 95%
 Total..... 100%

*Registered Trademark of
 Union Carbide Corporation
 for Carbaryl insecticide.

U.S.D.A. REG. NO. 557-1613
 Form No. 000

CAUTION: Keep out of reach of children.
 See other panel for additional caution.

VIGORO® 5% SEVIN® DUST

DIRECTIONS FOR USE. VIGORO 5% SEVIN® DUST is to be used without dilution. Apply uniformly to foliage covering all exposed parts (both upper and lower leaf surfaces) with a thin, even film of dust using any garden type duster. Apply preferably in

early morning or late evening when the air is quiet. Apply when insects or their damage appear. Best results are secured by dusting regularly at 7-10 day intervals throughout the growing season. Foliage should be dry before dust is applied.

CROPS	INSECTS	INTERVAL Last Day to Harvest
Beans	Lygus Bugs, Stink Bugs, Corn Earworm, Leafhoppers, Flea Beetles, Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetle, Western Bean Cutworm, Japanese Beetle, Tarnished Plant Bug, Velvetbean Caterpillar, Armyworms.	0 day
Sweet Corn	Corn Earworm Apply when first silks appear and repeat at 2 to 3 day intervals during silking period.	
Cucumber, Cantaloupes, Pumpkins, Squash, Watermelons, Eggplant, Okra, Peppers, Tomatoes, Carrots	Striped Cucumber Beetle, Spotted Cucumber Beetle, Melon worm, Pickleworm, Squash Bug, Leafhoppers, Cucumber Beetles, Flea Beetles, Armyworms, Corn Earworm, Tomato Fruitworm, Tomato Hornworms, Fall Armyworms, Tobacco Hornworms, European Corn Borrer, Stink Bugs, Japanese Beetle, Colorado Potato Beetle, Lace Bugs, Lygus Bugs	0 day
Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kohlrabi, Head Lettuce, Parsnips, Radishes, Rutabagas, Beets, Turnips, Salsify, Horseradish	Flea Beetles, Harlequin Bug, Corn Earworm, Armyworms, Cabbage Caterpillars, Imported Cabbage Worm, Cabbage Looper, Diamondback Moth, Stink Bugs, Leafhoppers, Lygus Bugs, Spittlebugs Do not use Beets, Salsify or Turnip tops for food.	3 days
Collards, Endive (Escarole), Kale, Leaf Lettuce, Mustard Greens, Parsley, Spinach, Swiss Chard, Chinese Cabbage	Flea Beetles, Stink Bugs, Corn Earworm, Leafhoppers, Armyworms	14 days
Grapes	Grape Leafhoppers, Grape Leaf Folder, Grape Berry Moth, European Fruit Locanium, Grape Leaf Skeletonizer, Japanese Beetle, June Beetles, Redbanded Leaf Roller	0 day
Strawberries	Flea Beetles, Lygus Bugs, Spittlebugs, Leafhoppers, Strawberry Leaf Roller	1 day
ORNAMENTALS: Arbor Vitae, Azalea, Aster, Birch, Boxwood, Carnation, Camellia, Chrysanthemums, Dahlia, Dogwood, Elm, Gladiolus, Holly, Juniper, Lilac, Marigold, Petunia, Rhododendron, Rose of Sharon, Yew, Zinnia	Bagworm, Blister Beetle, Boxelder Bugs, Leaf Miner (Dipterous), Leaf Beetles, Flea Beetle, Japanese Beetle, Leafhoppers, Leaf Rollers, Orange Tortrix, Periodical Cicada, Plant Bug, Tent Caterpillar, Thrips (Exposed)	
LAWNS & ORNAMENTAL TURF	Leafhoppers, Fall Armyworms, Sod Webworms (Lawn Moths), Ants, Fleas, Earwigs	Apply at the rate of 4 lbs. per 1000 sq. ft. For best results apply after rain or watering and do not water for at least 2 days after application. Repeat in 2 to 3 weeks if necessary.
Dogs and Cats	Fleas and Ticks	Dust liberally over the entire animal, avoiding the eye, and rub thoroughly into skin. Apply to feet between toes. Remove dead ticks by combing a few hours after treatment. Repeat at weekly intervals if needed. Do not treat kittens under 4 weeks of age.
Dog and Cat Quarters	Fleas and Ticks	Dust liberally in and around sleeping quarters, kennels, and dog houses by means of a shaker can, hand or dust gun. Repeat as needed. Clean quarters before treatment.

CAUTION: Harmful if inhaled or swallowed. Do not take internally. Avoid repeated breathing of dust. Skin contact may be harmful. Avoid contact. Wash hands before eating. Wash thoroughly after using. Do not feed treated forage to livestock. Keep out of reach of children and animals. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative State Agricultural Extension Service. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do not reuse container. Destroy when empty.

MANUFACTURED BY
SWIFT AGRICULTURAL CHEMICALS CORPORATION
 CHICAGO, ILLINOIS 60606

4-Ⓢ MV-69-4711

557-1613

7-13-1971

1/5

557-1613

ACCEPTED
JUL 13 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

557-1613

2/2