

# VIGORO®

## VIGORO® 5% SEVIN® DUST

**DIRECTIONS FOR USE, VIGORO 5% SEVIN® DUST** is to be used without dilution. Apply uniformly to foliage covering all exposed parts (both upper and lower leaf surfaces) with a thin, even film of dust using any garden type duster. Apply preferably in

early morning or late evening when the air is quiet. Apply when insects or their damage appear. Best results are secured by dusting regularly at 7-10 day intervals throughout the growing season. Foliage should be dry before dust is applied.

# 5% SEVIN® DUST

FOR USE ON  
**VEGETABLES  
LAWNS·SHRUBS  
DOGS·CATS**

**ACTIVE INGREDIENTS:**  
Carbaryl (1-Naphthyl  
N-methylcarbamate) 5%  
**INERT INGREDIENTS 95%**  
Total..... 100%

\*Registered Trademark of  
Union Carbide Corporation  
for Carbaryl insecticide.

U.S.D.A. REG. NO. 557-1613  
Form No. 000

NET WT. 16 OZ. (1 LB.)

**ACCEPTED**

OCT 19 1971

REGISTERED UNDER NO. 557-1613  
FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO.

| CROPS   | INSECTS   | INTERVAL<br>Last Day<br>to Harvest   |
|---|---|--|
| Beans   | Lygus Bugs, Stink Bugs, Corn Earworm, Leafhoppers, Flea Beetles, Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetle, Western Bean Cutworm, Japanese Beetle, Tarnished Plant Bug, Velvetbean Caterpillar, Armyworms.   | 0 day  |
| Sweet Corn  | Corn Earworm<br>Apply when first silks appear and repeat at 2 to 3 day intervals during silking period.   |  |
| Cucumber, Cantaloupes, Pumpkins, Squash, Watermelons, Eggplant, Okra, Peppers, Tomatoes, Carrots  | Striped Cucumber Beetle, Spotted Cucumber Beetle, Melon worm, Pickleworm, Squash Bug, Leafhoppers, Cucumber Beetles, Flea Beetles, Armyworms, Corn Earworm, Tomato Fruitworm, Tomato Hornworms, Fall Armyworms, Tobacco Hornworms, European Corn Borner, Stink Bugs, Japanese Beetle, Colorado Potato Beetle, Lace Bugs, Lygus Bugs | 0 day  |
| Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kohlrabi, Head Lettuce, Parsnips, Radishes, Rutabagas, Beets, Turnips, Salsify, Horseradish   | Flea Beetles, Harlequin Bug, Corn Earworm, Armyworms, Cabbage Caterpillars, Imported Cabbage Worm, Cabbage Looper, Diamondback Moth, Stink Bugs, Leafhoppers, Lygus Bugs, Spittlebugs<br>Do not use Beets, Salsify or Turnip tops for food.   | 3 days   |
| Collards, Endive (Escarole), Kale, Leaf Lettuce, Mustard Greens, Parsley, Spinach, Swiss Chard, Chinese Cabbage   | Flea Beetles, Stink Bugs, Corn Earworm, Leafhoppers, Armyworms  | 14 days  |
| Grapes  | Grape Leafhoppers, Grape Leaf Folder, Grape Berry Moth, European Fruit Locanium, Grape Leaf Skeletonizer, Japanese Beetle, June Beetles, Redbanded Leaf Roller  | 0 day  |
| Strawberries  | Flea Beetles, Lygus Bugs, Spittlebugs, Leafhoppers, Strawberry Leaf Roller  | 1 day  |
| <b>ORNAMENTALS:</b> Arbor Vitae, Azalea, Aster, Birch, Boxwood, Carnation, Camellia, Chrysanthemums, Dahlia, Dogwood, Elm, Gladiolus, Holly, Juniper, Lilac, Marigold, Petunia, Rhododendron, Rose of Sharon, Yew, Zinnia |   | Bagworm, Blister Beetle, Boxelder Bugs, Leaf Miner (Dipterous), Leaf Beetles, Flea Beetle, Japanese Beetle, Leafhoppers, Leaf Rollers, Orange Tortrix, Periodical Cicada, Plant Bug, Tent Caterpillar, Thrips (Exposed)  |
| <b>LAWNS &amp; ORNAMENTAL TURF</b>  | Leafhoppers, Fall Armyworms, Sod Webworms (Lawn Moths), Ants, Fleas, Earwigs  | Apply at the rate of 4 lbs. per 1000 sq. ft. For best results apply after rain or watering and do not water for at least 2 days after application. Repeat in 2 to 3 weeks if necessary.  |
| Dogs and Cats   | Fleas and Ticks   | Dust liberally over the entire animal, avoiding the eye, and rub thoroughly into skin. Apply to feet between toes. Remove dead ticks by combing a few hours after treatment. Repeat at weekly intervals if needed. Do not treat kittens under 4 weeks of age. Do not treat puppies under 4 weeks of age. |
| Dog and Cat Quarters  | Fleas and Ticks   | Dust liberally in and around sleeping quarters, kennels, and dog houses by means of a shaker can, hand or dust gun. Repeat as needed. Clean quarters before treatment.   |

**CAUTION:** Harmful if inhaled or swallowed. Do not take internally. Avoid breathing of dust. Skin contact may be harmful. Avoid contact. Wash hands before eating. Wash thoroughly after using. Do not feed treated forage to livestock. Keep out of reach of children and animals. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative State Agricultural Extension Service. **NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do not reuse container. Destroy when empty.

557-1613

10-19-1971

1/1