

Reg # 557-1706

PM-23

195



CERTIFICATION OF COMPLIANCE

I, Judith K. Lucash, being an authorized representative of Vigoro Industries, Inc., certify that all containers of

K Mart Weed And Feed

EPA Reg. No. 557-1706

produced by June 15, 1994 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing use products. I further certify that all containers of said product sold or distributed by this company by January 1, 1995 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products.


Judith K. Lucash
Manager of Regulatory Affairs

December 21, 1992

225

FRONT PANEL

Kmart® WEED and FEED

Covers 5000 Sq. Feet

KEEP OUT OF REACH OF CHILDREN

CAUTION

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention. See back panel for additional precautionary statements.

ACTIVE INGREDIENT:

*Dimethylamine salt of 2,4-dichlorophenoxyacetic acid 1.22%

INERT INGREDIENTS: 98.78%

TOTAL 100.00%

*Equivalent to 2,4-dichlorophenoxyacetic acid 1.01%

EPA Reg. No. 557-1706

EPA Est. No. 557-CA-1

NET WT. 20 LBS.



395

Kmart

WEED and FEED

10 - 6 - 4

GUARANTEED ANALYSIS

TOTAL NITROGEN (N) 10.0%
 10.0% Ammoniacal Nitrogen
 AVAILABLE PHOSPHORIC ACID (P₂O₅) 6.0%
 SOLUBLE POTASH (K₂O) 4.0%
 Derived from ammoniated superphosphate, ammonium sulfate and
 muriate of potash. Potential Acidity 200 lbs. Calcium Carbonate
 Equivalent per ton.

PRECAUTIONARY STATEMENTS
 HAZARD TO HUMANS & DOMESTIC ANIMALS
 CAUTION

Avoid contact with skin, eyes and clothing. Avoid breathing dust. Wash thoroughly after handling. Harmful if swallowed. Keep out of reach of children. When using this product, wear long-sleeved shirt, long pants, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment washwaters. Do not contaminate any body of water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from target area. Do not contaminate irrigational ditches or water used for domestic purposes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment areas during application. Do not enter treatment areas until spray has dried or dust has settled.

KILLS WEEDS AS IT FEEDS YOUR LAWN

WEED AND FEED YOUR LAWN IN ONE EASY APPLICATION. FAST, EASY WAY TO FEED YOUR LAWN AND KILL WEEDS TOO. SPECIALLY FORMULATED TO KILL DANDELIONS AND PLANTAIN.

THIS WEED KILLER ALSO KILLS:

BLACK MEDIC
 BUCKHORN PLANTAIN
 COMMON CHICKWEED
 DOCK
 GOLDENROD
 GROUND IVY

NETTLE
 PIGWEED
 RAGWEED
 SHEPHERDS PURSE
 THISTLES
 WILD CARROT

WILD GARLIC
 WILD MORNING GLORY
 WILD MUSTARD
 WILD ONION
 AND OTHER
 BROADLEAVED WEEDS

REAR PANEL

The plant food supplies the quick pick-up needed to fill out bare spots left by dead weeds and help prevent re-establishment of weeds by encouraging growth and formation of thick, weed resistant turf.

APPLY ONLY IN WARM WEATHER WHEN SOIL MOISTURE IS PLENTIFUL AND WEEDS ARE ACTIVELY GROWING.

Weeds are hard to kill when weather is cold in the early spring or during hot, dry summer weather. For best results, apply while lawn is wet with dew or after sprinkling lightly. Apply several days after mowing to allow for the presence of plenty of leaf surface to catch and retain Kmart WEED & FEED, then do not mow or water lawn for another day. Set your spreader as shown below, then cover your lawn evenly.

The maximum application rate for this product is 4.0 lbs. per 1,000 sq. ft. The maximum number of broadcast applications per treatment site is 2 per year.

Approximately 4 weeks after weed kill is evident, an application of 16% or higher nitrogen fertilizer should be made at a rate of 6 lbs. per 1,000 sq. ft. and watered in well.

Storage: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal. Keep bag tightly closed. Do not store near growing plants.

Product Disposal: Securely wrap original container in several layers of newspaper and discard in trash. Container Disposal: Do not re-use bag. Discard bag in trash.

SUGGESTED SPREADER SETTING

Note: Mechanical condition of spreader, walking speed, climatic conditions and terrain can cause wide variation in application rate. Calibrate spreader with each application. Close spreader openings when loading, turning or when stopped. Hand spreading is not recommended. One bag WEEDS AND FEEDS 5,000 sq. ft.

SPREADER	SUGGESTED SETTING
PRECISION 20C	4 1/2
CENTRAL 1662	6 1/2
SEARS CRAFTSMAN	5 1/4
Model 452-19203 (20")	
Model 452-19183 (18")	
Model 452-19194 (19")	
WARD'S GARDEN MARK	6 1/4
Model AX-109 3A	
LAWN BEAUTY	5
16", 18" or 24"	
CYCLONE	3 3/4
Model B & C	
SCOTTS 35-8	4

Clean spreader after each use.

595

REAR PANEL CONT.

CAUTION:

Avoid drift to or contact with vegetables, flowers, shrubs, ornamentals, or other plants that are sensitive to 2,4-D. Do not apply to areas underlaid by roots of desirable trees and shrubs.

Not recommended for use on bent grass, dichondra, centipede grass, St. Augustine grass or on lawns where clover is present and desired. DO NOT APPLY TO NEWLY SEEDED LAWNS.

MANUFACTURED IN U.S.A. FOR Kmart CORPORATION, TROY, MICHIGAN 48084

BY VIGORO INDUSTRIES, INC., FAIRVIEW HEIGHTS, IL 62208