

CORNER

FRONT

CORNER

SIDE

CORNER

BACK

29-1.1373

VIGORO

TREATS
4000
SQ. FT.

CHICKWEED, CLOVER AND WEED KILLER

KEEP FROM FREEZING

ACCEPTED

JUN 15 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 557-1752

SURE DEATH FOR MANY BROAD-LEAVED WEEDS

KILLS: DANDELIONS KNOTWEED HENBIT
CHICKWEED PLANTAINS SPURGE
AND MANY OTHER BROADLEAF WEEDS.

contains 2, 4-D, mcpp and dicamba

U. S. Patent No. 3,284,186

ACTIVE INGREDIENTS:

* Diethanolamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid 13.7%
** Diethanolamine salt of 2,4-dichlorophenoxyacetic acid 13.6%
*** Diethanolamine salt of Dicamba (3,6-dichloro-o-anisic acid) 1.3%

INERT INGREDIENTS

Total 71.4%
100.0%
*2-(2-methyl-4-chlorophenoxy) propionic acid equivalent 0.83 pounds per gallon
**2,4-dichlorophenoxyacetic acid equivalent 0.83 pounds per gallon
***3,6-dichloro-o-anisic acid equivalent 0.083 pounds per gallon

CAUTION: Keep Out of Reach of Children
See Back Panel For Additional Cautions.

NET 16 FL. OZ. (1 PT.)

DIRECTIONS FOR USE

WHERE TO USE: This product may be used on Bluegrass, Bermuda and Zoysia lawns, vacant lots, play ground and other similar areas where HARD-TO-KILL weeds are present. It is especially effective against these weeds such as SHEEP SORREL, KNOTWEED, SPOTTED SPURGE, CHICKWEED, OXALIS AND HENBIT. In addition to these, it kills a large number of weeds, some of which are listed below.

Do not allow spray to get on vegetable or ornamental plants since injury may result. Some injury can be expected to turf when used on Centipede - St. Augustine grass lawns. "Spot Spray" weeds only. Clover present will be damaged if sprayed.

Bedstraw	Clover	Henbit	Morning Glory	Poison Oak	Spurge
Black Medic	Dandelion	Knotweed	Peppergrass	Purslane	Wild Carrot
Buckhorn	Dock	Lambsquarter	Pigweed	Ragweed	Wild Garlic
Chicory	Ground Ivy	Lespedeza	Plantains	Shopards	Wild Lettuce
Chickweed	Heal-all	Mallow	Poison Ivy	Purse	Wild Onion
				Speedwell	Yarrow

WHEN TO USE: Spray at any time during the growing season when weeds are actively growing. Treat Chickweed, Henbit, Wild Onion and Garlic in the fall and again in the spring if needed. On new lawns wait until the turf has reached a height of 2 inches before making an overall spray treatment. "Spot spray" weed treatment, however, may be made at any time with a small pressure sprayer spraying only the weeds at ground level. Wait 3 weeks before putting new grass seed or stolens down in the bare spots. Poor weed control will result if spray is applied during drought or just before rain is expected. Do not water lawn within 24 hours after treatment.

HOW MUCH TO USE: TO SPRAY ENTIRE LAWN

1 Pint covers up to 4,000 sq. ft. of Lawn Area
Dilute with water as indicated below.

For Pressure Sprayers	Lawn Hose Sprayers**	Area to Spray
2 tblsp. in 1/2 gal.	2 tblsp. for 2 gals.	250 sq. ft.
4 oz. (8 tblsp.) in 1 1/2 gal.	8 tblsp. for 6 gals.	1000 sq. ft.
8 oz. in 3 gals.	16 tblsp. for 12 gals.	2000 sq. ft.
1 pint in 6 gals.	1 pint for 24 gals.	4000 sq. ft.

**For Lawn Hose Sprayers dilute the amount shown above with enough water to fill the sprayer bottle to the 2 gal. level for 250 sq. ft. to the 6 gal. level for 1000 sq. ft. or to the 12 gal. level for 2000 sq. ft.

TO SPOT SPRAY WEEDS

Lawn Hose Sprayers are not suited for this type of treatment.
Use a pressure sprayer.
Mix 8 oz. in 3 gals. water
and spray each weed to point of spray run-off.

CAUTION: Avoid contact with eyes or skin. In case of contact, wash with plenty of water. Avoid breathing a spray mist.

When container is empty, wash thoroughly with water. Never re-use. Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not contaminate domestic or irrigation water. Do not pour spray solutions near these plants. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Do not spray on carpet, grass, ditches, or lawns or turf where desirable bentgrass or fescues are present. Use only lawn sprayers for this use. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. This product may be injurious to St. Augustine grass.

DIST BY SWIFT AGRICULTURAL CHEMICALS CORPORATION, CHICAGO, ILLINOIS 60606
U.S.D.A. Reg. No. 557-1752 Form-2217433-69

CUSTOMER	GORDON CORP.	ENG.	29-1-1373	FACTY.	068	SIZE	313x111x514	
LAYOUT	068 #121	INDEX NO.	0-54	D.F.	5-71	STYLE	1PT. OBL.	
COLORS	RED - GREEN - BLACK ON WHITE COAT						ARTIST	9088+41

9104