

VIGORO.

ACCEPTED
JAN 25 1974
UNLESS OTHERWISE SPECIFIED
FOR PESTICIDE AND PESTICIDE ACT
FOR PESTICIDE AND PESTICIDE ACT

VIGORO



SNAIL
& SLUG
KILLER
MEAL

SNAIL & SLUG KILLER MEAL

ATTRACTS &
KILLS
SNAILS
& SLUGS

For use in Roses, Flow-
ers, Shrubs, Grass,
Dichondra, and other
outdoor areas around
homes

TRY THESE OTHER
FINE VIGORO
SPECIALTY
PRODUCTS

VIGORO Citrus and
Avocado Food

VIGORO Azalea, Rhododendron,
Camellia Food


VIGORO Rose Food

VIGORO Evergreen
Food

VIGORO Tomato Food

VIGORO Acid Food
(0-10-10)

ACTIVE INGREDIENTS
Metaldehyde 3.35%
INERT INGREDIENTS ... 96.65%
TOTAL 100.00%



NET WT. 40 OZS. (2.5 LBS.)

WARNING: Keep Out of Reach of Children. See Back Panel for Additional Cautions.

THIS PESTICIDE MAY BE FATAL TO CHILDREN, DOGS, OR OTHER PETS IF EATEN. KEEP CHILDREN AND PETS OUT OF THE TREATED AREA. SCATTER MATERIAL. DO NOT LEAVE IN PILES. FOLLOW DIRECTIONS AND USE WITH CARE.

E.P.A. Reg. No. 557-1753-AA

VIGORO.

SNAIL & SLUG KILLER

VIGORO

VIGORO



SNAIL
& SLUG
KILLER
MEAL

SNAIL & SLUG KILLER MEAL

ATTRACTS & KILLS SNAILS & SLUGS

Vigoro Snail & Slug Killer is a special preparation for the control of snails. Vigoro Snail & Slug Killer attracts and kills snails and slugs with Metaldehyde. The presence of snails and slugs can be detected by their shiny mucuous-like trail. They are night feeders and prefer damp, cool places in the ornamental garden, flowers, under boards or flower pots and around fences, hedges and bushes.

NOTE: This package was full when packed. Contents are sold by weight.

DIRECTIONS

Water area to be treated thoroughly. Broadcast Vigoro Snail & Slug Killer late in the day in the areas likely to be infested. Apply to the soil and not directly on plants. Scatter, do not leave in piles.

Continuous watering and high temperatures cause this bait to gradually lose some of its effectiveness. Repeat applications every week or ten days during spring and summer months. Treatment every two weeks during fall and winter months is usually satisfactory. Repeated treatments provide control of new broods emerging during this period. Three baitings ten days apart, usually gives control of new hatch.

Apply one pound of meal to each 250-300 square feet.

WARNING

MAY BE FATAL IF SWALLOWED. Harmful dust. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. Avoid food contamination. Wash hands with soap and water after using.

IF SWALLOWED, INDUCE VOMITING, CALL A PHYSICIAN IMMEDIATELY. Keep away from children, pets, and livestock. Keep animals and poultry off treated areas. This bait is attractive to dogs and chickens. Keep off plants. Do not reuse container. Destroy when empty. Birds feeding on treated areas may be killed.

MONEY-BACK GUARANTEE

YOU MUST BE PLEASED WITH RESULTS OR YOU ARE ENTITLED TO YOUR MONEY BACK. SIMPLY SEND SALES SLIP TO VIGORO, CHICAGO, ILL. 60604 FOR PROMPT REFUND.

MANUFACTURED BY
SWIFT CHEMICAL COMPANY • CHICAGO, ILLINOIS 60604 • HOUSTON, TEXAS 77007