


*PREVENTS
UGLY CRABGRASS
AS IT FEEDS
YOUR LAWN*



Deep Green VIGORO **LAWN FERTILIZER PLUS
CRABGRASS PREVENTER**

VIGORO
LAWN WEEDER & FEEDER

NET WT 35 LBS
Covers 2,300 sq ft

VIGORO

LAWN WEEDER & FEEDER

**FOR EXCLUSIVE USE ON CENTIPEDE,
ST AUGUSTINE & ZOYSIA GRASSES
12-5-7**

GUARANTEED ANALYSIS

TOTAL NITROGEN, not less than	12.00%
NITRATE NITROGEN, not less than	NONE
AMMONIACAL NITROGEN, not less than	2.00%
WATER SOLUBLE ORGANIC NITROGEN, not less than	6.50%
WATER SOLUBLE NITROGEN, not less than	3.50%
AVAILABLE PHOSPHORIC ACID, not less than	5.00%
SOLUBLE PHOSPHORIC ACID, not less than	0.50%
WATER SOLUBLE POTASH, not less than	7.00%
WATER SOLUBLE PRIMARY PLANT FOOD, not less than	24.00%
CHLORINE, not more than	1.00%

Control from Urea, Ammoniated Superphosphate, Ureafume, Sulfate of Potash (LA REG. NO. 732-735)
1964. Manufactured by Smith Agricultural Chemicals Corp., Chicago 60606, Western Division, U.S.A.

Controls Such Weeds as:

Controlled: Crabgrass, Barnyardgrass, Wiregrass, Paspalum Seedlings,
Barnyardgrass, Barnyardgrass, Sedges (such as Cyperus Compressus),
and others.

Apply between January 1 and March 31 (Florida to June 15th).

Established and Germinating: Betony, Blackmedic (yellow trefoil) Burclover,
Chickweed, Dandelion, Oxalis, Florida Pusley, Pennywort (Dollarweed) and
Creeping Charlie (Mintweed).

EASY TO USE

For best results apply before weeds come up (pre-emergent). Established weeds are more difficult to control and may require an application in Spring and Fall. On newly sprigged St. Augustine, Centipede, or Zoysia Lawns, apply immediately after planting to prevent weeds from retarding growth.

Water in well.

CAUTION: Keep out of the reach of children. Keep away from feed or food products. Do not take internally. Avoid inhalation or contact with eyes. Wash after handling or using.

Do not use north of North Carolina. Do not use on lawns of Bahiagrass, Bermudagrass, Dichondra, Bentgrass, or carpetgrass or where clover is present and desired.

Avoid application in the root zones of trees and shrubs. Avoid applications on garden plants, flowers, and shrubs. Delay spring planting for one month after application. Store in a dry place. Do not store near growing plants or fertilizers, seeds, insecticides or fungicides. Clean spreader after use.

Apply only in Winter, Spring and Fall. This product is not recommended for use during the months of July, August and September. Do not reuse empty bag. Destroy when empty.

HOW TO SET YOUR SPREADER

CYCLONE.....6	SCOTTS.....7½
K-MART.....9	SEARS.....7½
LAWN BEAUTY.....9	WARDS.....10

NOTE: Spreader rate varies with walking speed; check spreader over known area.

557-1763

ACCEPTED

Jun 8 1971

UNDER THE PROVISIONS OF THE
NATIONAL BUREAU OF STANDARDS ACT
FOR ECONOMIC PROTECTION OF
CONSUMERS

VIGORO LAWN WEEDER
& FEEDER

VIGORO
LAWN WEEDER & FEEDER


NET WT 35 LBS
TREATS
2300 SQ
FT

VIGORO
LAWN WEEDER & FEEDER

**FOR EXCLUSIVE USE ON
CENTIPEDE, ST. AUGUSTINE
& ZOYSIA GRASSES**

CAUTION: Keep out of reach of children.
See back panel for additional cautions.

**WEEDS
AS IT
FEEDS**



ACTIVE INGREDIENTS:
*Atrazine (2-chloro-4-ethylamino-
6-isopropylamino s-triazine)..... 0.27%
INERT INGREDIENTS..... 99.73%
Total..... 100.00%

By Weight

*Contains 5.4 lbs. of atrazine per ton

USDA Reg No 557-1763

VIGORO LAWN WEEDER
& FEEDER