

GOLDEN VIGORO


NET WT 35 LBS

1146

GOLDEN

**LAWN FERTILIZER
PLUS
CRABGRASS PREVENTER**

2-10-71
557-1782



GOLDEN VIGORO

GOLDEN VIGORO

NET WT 35 LBS

ACCEPTED
2-10-71
557-1782
FUNGICIDE
FOR
ED UN

GOLDEN VIGORO

LAWN FERTILIZER
PLUS
CRABGRASS PREVENTER

11-4-6
GUARANTEED ANALYSIS

TOTAL NITROGEN 11.0%
3.7% Water Soluble Nitrogen
6.4% Water Soluble Nitrogen
AVAILABLE PHOSPHORIC ACID (P₂O₅) 4.0%
SOLUBLE POTASH 4.0%
Potassium Sulfate
Agricultural Chemicals Corporation, Chicago, Illinois

You must be pleased with the results or you are entitled to your money back. Simply send sales slip to VIGORO, 2 North Riverside Plaza, Chicago, Illinois 60606 for prompt refund.

Apply in early spring before germination of crabgrass seed occurs.
Set your spreader by the chart below, then cover the area twice; the second time at right angles to the first. Apply to only 2500 sq. ft. - no more!
Water lightly after one week.
On newly seeded lawns, wait until after the third mowing.
Do not rake lawn or disturb soil surface after application.
Do not apply to dichondra lawns.

CAUTION: Harmful if swallowed. Do not take internally. Avoid contact with skin, eyes, and clothing. Wash after handling or using. Do not contaminate feed or foodstuffs. Store in a dry place.
Do not reuse this container. Destroy when empty.

CYCLONE	4½	SCOTTS	6
K-MART	5	SEARS	6
LAWN BEAUTY	8	WARDS	9

NOTE: Spread on lawns with walking spreaders.

GOLDEN VIGORO

LAWN FERTILIZER PLUS CRABGRASS PREVENTER

11-4-6

GUARANTEED ANALYSIS

TOTAL NITROGEN (N) 11.0%
 0.9% Ammoniacal Nitrogen (derived from ammoniated sulfates and ammoniated phosphates)
 3.7% Water Insoluble Nitrogen (derived from condensed urea-formaldehyde)
 6.4% Water Soluble Nitrogen (derived from urea-formaldehyde)
 AVAILABLE PHOSPHORIC ACID (P₂O₅) (derived from ammoniated phosphates) 4.0%
 SOLUBLE POTASH (K₂O) (derived from muriate of potash) 6.0%
 Potential acidity 700 lbs Calcium Carbonate Equivalent per ton. Manufactured by Swift Agricultural Chemicals Corporation, Chicago, Illinois 60606.

MONEY BACK PROMISE

You must be pleased with the results or you are entitled to your money back. Simply send sales slip to VIGORO, 2 North Riverside Plaza, Chicago, Illinois 60606 for prompt refund.

EASY TO USE

- Apply in early spring before germination of crabgrass seed occurs.
- Set your spreader by the chart below, then cover the area twice; the second time at right angles to the first. Apply to only 2500 sq. ft. — no more!
- Water in lightly after one week.
- On newly seeded lawns, wait until after the third mowing.
- Do not rake lawn or disturb soil surface after application.
- Do not apply to dichondra lawns.

CAUTION: Harmful if swallowed. Do not take internally. Avoid contact with skin, eyes, and clothing. Wash after handling or using. Do not contaminate feed or foodstuffs. Store in a dry place.
 Do not reuse this container. Destroy when empty.

HOW TO SET YOUR SPREADER

CYCLONE	4½	SCOTTS	6
K-MART	5	SEARS	6
LAWN BEAUTY	8	WARDS	9

NOTE: Spreader rate varies with walking speed, check spreader over known area.

ACTIVE INGREDIENTS:

Dimethyl ester of tetrachloroterephthalic acid* 5.00%

INERT INGREDIENTS.


95.00%

100.00%

* U.S. Pat. 2,810,000; also see patented U.S. Pat. 2,723,074

CAUTION

KEEP OUT OF REACH OF CHILDREN
SEE BACK OF BAG FOR ADDITIONAL CAUTION



ROCKLAND CHEMICAL CO., INC.

WEST CALDWELL, N. J.

ROCKLAND CRABGRASS PREVENTER



GENERAL INFORMATION

ROCKLAND CRABGRASS PREVENTER is a highly effective pre-emergence herbicide. When applied as recommended, ROCKLAND CRABGRASS PREVENTER will give excellent control of crabgrass and the following weeds in turf and ornamental plantings: Carpetweed, Common Chickweed, Fall Panicum, Florida Pursley, Green Foxtail, Hairy Crabgrass, Johnsongrass, Lambsquarters, Lovegrass, Purslane, Smooth Crabgrass, Texas Millet, Witchgrass, Yellow Foxtail.

DIRECTIONS

WHEN TO APPLY: ROCKLAND CRABGRASS PREVENTER kills crabgrass and other annual weeds as they germinate, but not after they have rooted. To be effective, ROCKLAND CRABGRASS PREVENTER must be applied before weed seed germination in early spring. This usually coincides with the time when Forsythia blossoms start to fall. A single spring application should suffice for most geographic locations. Make application on a calm day to avoid wind drift.

HOW MUCH TO APPLY: Apply contents of bag (12 lbs.) evenly to 1700 sq. ft. (an area 50 x 50 ft.). Use any lawn spreader in good condition. Do not spread over larger areas than recommended as such application would give poor results. To control annual bluegrass or other annual or late-germinating grasses, a late summer or fall application of 12 lbs. per 1700 sq. ft. should supplement the early spring application (contents of bag will treat 1600 sq. ft.).

After each application, lawn should be raked to remove extraneous ma-