

FRONT PANEL

Reg # 557-1812

PM-23

123

Contains 0.9% Iron

Treats 5,000 Sq.Ft.
17-3-4

Golden Vigoro

Weed Control plus Fertilizer

WEEDS
AS IT
FEEDS

Formulated with IBDU
a patented form of slow
release nitrogen

ACCEPTED
OCT 14 1987
Federal Insecticide
Fungicide and Rodenticide
Act
557-1812

KEEP OUT OF REACH OF CHILDREN

C A U T I O N

In case of contact, wash skin with soap and water;
for eyes, flush with water for 15 minutes and get
medical attention.

See back panel for additional precautionary statements.

ACTIVE INGREDIENT

*Dimethylamine salt of 2,4-dichloro-
phenoxyacetic acid 0.918%
INERT INGREDIENTS 99.082%

Total 100.00%

*Equivalent to 2,4-dichlorophenoxyacetic
acid 0.765%

*Isomer Specific by AOAC method 6D01-5

E.P.A. Est. No. 557-IL-1

EPA Reg. No. 557-1812

NET WT. 25 LBS.

NET WT 25 LBS.

GOLDEN VIGORO

WEED CONTROL PLUS FERTILIZER

17-3-4

GUARANTEED ANALYSIS

TOTAL NITROGEN (N) 17.0%
 0.9% Ammoniacal Nitrogen
 11.4% Urea Nitrogen
 4.7% Water Insoluble Nitrogen
 AVAILABLE PHOSPHORIC ACID (P_2O_5) 3.0%
 SOLUBLE POTASH (K_2O) 4.0%
 Iron (Fe) 0.9%
 Derived from ammoniated phosphate, urea, isobutylidene diurea, muriate
 of potash and iron sulphate. Potential Acidity 700 lbs. Calcium Carbonate
 Equivalent per ton. Manufactured by Vigoro Industries, Inc., Fairview
 Heights, IL 62208.

PRECAUTIONARY STATEMENTS
 HAZARD TO HUMANS & DOMESTIC ANIMALS

CAUTION

Avoid contact with skin, eyes or clothing. Avoid breathing dust.
 Wash thoroughly after handling. Harmful if swallowed.

ENVIRONMENTAL HAZARD

Do not apply directly to water. Do not apply when weather conditions
 favor drift from target area. Do not contaminate water intended for
 irrigation or domestic use.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner
 inconsistent with its labeling.

Apply when weeds are young and growing actively. Weather must be above
 70° and below 90°. Water lawn one day before application to ensure
 adequate soil moisture. Do not apply to wet grass. To get maximum weed
 killing results, do not rake, mow or water lawn until two days after
 applying. Repeat treatment, if necessary, to control hard to kill weeds.
 Controls or kills dandelions, plantain, chickweed, clover, oxalis, spurge,
 ground ivy, burr clover and many other similar broadleaf weeds. FOR USE ON
BLADE GRASS LAWNS ONLY.

Storage: Store in original con-
 tainer in safe, dry place. Do
 not contaminate other pesticides,
 fertilizer, seeds, food or feed.
 Return spillage to original con-
 tainer prior to disposal.

Product Disposal: Securely wrap
 original container in several
 layers of newspaper and discard
 in trash.
 Container Disposal: Do not re-use
 bag. Discard bag in trash.

SUGGESTED SPREADER SETTING ✓

SPREADER	SETTING	SPREADER	SETTING
Cyclone	4 1/2	Central Drop	10
Scotts PF-1	6 1/2	Central Broadcast	4 1/2
Wards	6 1/2	Precision 20C	4

NOTE: Spreader setting varies with terrain, walking speed and condition of equipment. To insure proper coverage, calibrate before use by checking over measured area.

CAUTION: Close spreader openings when loading, turning or when stopped.

PRECAUTIONS: Not recommended for use on St. Augustine grass, carpetgrass, bentgrass, dichondra, centipede grass, or on lawns where clover is present and desired. Do not apply to newly seeded lawns. Do not store near growing plants. Avoid drift or contact with vegetables, flowers, shrubs, ornamentals or other plants that are sensitive to 2,4-D. Do not apply to areas underlaid by roots of desirable trees and shrubs. Clean spreader after each use. ✓