



INDUSTRIES, INC.

CERTIFICATION OF COMPLIANCE

I, Judith K. Lucash, being an authorized representative of Vigoro Industries, Inc., certify that all containers of

Golden Vigoro 17-3-4 Weed Control Plus Fertilizer Weed as it Feeds

EPA Reg. No. 557-1819

produced by June 15, 1994 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing use products. I further certify that all containers of said product sold or distributed by this company by January 1, 1995 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products.

  
Judith K. Lucash  
Manager of Regulatory Affairs

December 21, 1992

274

FRONT PANEL

Contains 0.9% Iron

Treats 5,000 Sq. Ft.  
17-3-4

Golden Vigoro

Weed Control plus Fertilizer

WEEDS  
AS IT  
FEEDS

Formulated with IBDU®  
a patented form of slow  
release nitrogen

-----  
KEEP OUT OF REACH OF CHILDREN  
CAUTION

In case of contact, wash skin with plenty of soap  
and water; for eyes, flush with water for 15 minutes  
and get medical attention. See back panel for  
additional precautionary statements.  
-----

-----  
ACTIVE INGREDIENT:


+*Dimethylamine salt of 2,4-dichlorophenoxyacetic		
acid . . . . .		0.918%
INERT INGREDIENTS . . . . .		99.082%
TOTAL . . . . .		100.000%

\*Equivalent to 2,4-dichlorophenoxyacetic acid 0.765%  
+Isomer Specific by AOAC method 6.D01-5

EPA Reg. No. 557-1819

EPA Est. No. 557-IL-1

-----  
NET WT. 25 LBS.



NET WT. 25 LBS.

GOLDEN VIGORO®

WEED CONTROL PLUS FERTILIZER

17-3-4

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	17.0%
0.9% Ammoniacal Nitrogen	
11.4% Urea Nitrogen	
4.7% Water Insoluble Nitrogen	
AVAILABLE PHOSPHORIC ACID (P <sub>2</sub> O <sub>5</sub> )	3.0%
SOLUBLE POTASH (K <sub>2</sub> O)	4.0%
Iron (Fe)	0.9%

Derived from ammoniated phosphate, urea, isobutylidene diurea, muriate of potash and iron sulphate. Potential Acidity 700 lbs. Calcium Carbonate Equivalent per ton.

PRECAUTIONARY STATEMENTS  
HAZARD TO HUMANS & DOMESTIC ANIMALS  
CAUTION

Avoid contact with skin, eyes and clothing. Avoid breathing dust. Wash thoroughly after handling. Harmful if swallowed. Keep out of reach of children. When using this product, wear long-sleeved shirt, long pants, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

ENVIRONMENTAL HAZARD

This product is toxic to aquatic invertebrates. Drift or run-off may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate any body of water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area. Do not contaminate irrigational ditches or water used for domestic purposes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment areas during application. Do not enter treatment areas until spray has dried or dust has settled.

Apply when weeds are young and growing actively. Weather must be above 70° and below 90 . Water lawn one day before application to ensure adequate soil moisture. Do not apply to wet grass. To get maximum weed killing results, do not rake, mow or water lawn until two days after applying. Repeat treatment, if necessary, to control hard to kill weeds. Controls or kills dandelions, plantain, chickweed, clover, oxalis, spurge, ground ivy, burr clover and many other similar broadleaf weeds. FOR USE ON BLADE GRASS LAWNS ONLY.

The maximum application rate for this product is 5.0 lbs. per 1,000 sq. ft. The maximum number of broadcast applications per treatment site is 2 per year.

Approximately 4 weeks after weed kill is evident, an application of 16% or higher nitrogen fertilizer should be made at a rate of 6 lbs. per 1,000 sq. ft. and watered in well.

Storage: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal. Keep bag tightly closed. Do not store near growing plants.

Product Disposal: Securely wrap original container in several layers of newspaper and discard in trash. Container Disposal: Do not re-use bag. Discard bag in trash.

SUGGESTED SPREADER SETTING

SPREADER	SETTING	SPREADER	SETTING
Cyclone	4 1/2	Central Drop	10
Scotts PF-1	6 1/2	Central Broadcast	4 1/2
Wards	6 1/2	Precision 20C	4

NOTE: Spreader setting varies with terrain, walking speed and condition of equipment. To insure proper coverage, calibrate before use by checking over measured area.

CAUTION: Close spreader openings when loading, turning or when stopped.

PRECAUTIONS: Not recommended for use on St. Augustine grass, carpetgrass, bentgrass, dichondra, centipede grass or where clover is present and desired. Do not apply to newly seeded lawns. Avoid drift or contact with vegetables, flowers, shrubs, ornamentals or other plants that are sensitive to 2,4-D. Do not apply to areas underlaid by roots of desirable trees and shrubs. Clean spreader after each use.

Manufactured by Vigoro Industries, Inc., Fairview Heights, IL 62208. Vigoro® and IBDU® are registered trademarks of Vigoro Industries, Inc.