
PLEASE NOTE

This image contains more than one label approved for this product on this date.



Ġ



Water area to be treated thoroughly. Broadcast $Swift_{Snail \& Slug Killer}$ late in the day in the areas likely to be infested. Apply to the soil and not directly on plants.

Continuous watering and high temperatures cause this bait to gradually lose some of its effectiveness. Repeat applications every week or ten days during spring and summer months. Treatment every two weeks during fall and winter months is usually satisfactory. Repeated treatments provide control of new broods emerging during this period. Three baitings ten days apart, usually gives control of new hatch.

Apply one pound of meal to each 200-250 square feet.

WARNING

MAY BE FATAL IF SWALLOWED. Harmful dust. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. Avoid food contamination. Wash hands with soap and water after using.

IF SWALLOWED, INDUCE VOMITING, CALL A PHYSICIAN IMMEDIATELY. Keep away from children, pets, and livestock. Keep animals and poultry off treated areas. This bait is attractive to dogs and chickens. Keep off plants. Do not reuse container. Destroy when empty.

This product is toxic to wildlife. Birds feeding on treated areas may be killed.

Do not contaminate water by cleaning equipment, or disposal of wastes.

MANUFACTURED

BY SWIFT AGRICULTURAL CHEMICALS CORP., CHICAGO, ILL. 50606

U.S.D.A. Reg. No. 557-1824

TOP AND POTTON PATELS

;




SMAIL & SLUG KOLLER MEAL

LEFT CATO PALVE.

ţi.

RIGHT SIDE PAUL




NEXT

. مرد


LABEL.

•

<u>_</u>...



BACK PAPFL



U.S.D.A. Reg No. 557-1824

TOP AND BOTTOM PANFLS




Suisfaction Always

100°P (FTT ENTED

PIGHT STEE PARTS

PULL OUT PUSH HERE





n