

VIGORO®

BEST DOCUMENT AVAILABLE

ACCEPTED

8-26-71

557-1825

UNDER THE FEDERAL FOOD, DRUG, AND COSMETIC ACT, THIS PRODUCT IS NOT TO BE USED IN THE MANNER SHOWN ON THE LABEL.

WEED & GRASS KILLER

NON-SELECTIVE
KILLS MOST VEGETATION

FOR USE ON DRIVEWAYS
FENCE ROWS • FIREBREAKS
PARKING LOTS and WALKS

NET WT. 32 OZ. (2 LB.)

ACTIVE
INGREDIENTS:
Sodium Metaborate Tetrahydrate
($\text{Na}_2\text{B}_4\text{O}_7 \cdot 4\text{H}_2\text{O}$) 66.50%
(Boron Trioxide
(B_2O_3) Equivalent = 22.7%)
Sodium Chlorate
(NaClO_3) 30.00%
Diuron [3-(3,4-dichlorophenyl)-
1,1-dimethylurea] 1.25%
INERT
INGREDIENTS 2.25%
TOTAL 100.00%

• REGISTERED
TRADEMARK
U. S. BORAX &
CHEMICAL
COMPANY
U. S. PATENT
NO. 3,032,405
U. S. D. A. REG.
NO. 557-1825

CONTAINS
MONOBOR-
CHLORATE®
GRANULAR D

VIGORO WEED & GRASS KILLER

DIRECTIONS FOR USE

Apply uniformly with shaker top container to areas such as drive-ways, patios, walks, fence lines and posts, and other similar places where vegetation is undesirable. Vigoro Broad-Spectrum Weed and Grass Killer is specifically designed for quick knock-down and kill of growing weeds.

GENERAL VEGETATION CONTROL: Before applying, sprinkle vegetation or apply when vegetation is wet with dew. Apply at the rate of $\frac{1}{2}$ to 2 pounds per 100 square feet. The lighter rate should be used only at time of weed emergence and soil is moist or when precipitation is anticipated (artificial watering may be used) while weeds are still young.

PERSISTENT PERENNIAL WEED CONTROL: These types of weeds are most effectively treated when nearing maturity or during dormant stages in the fall. Apply 2 to 4 pounds per 100 square feet and water well so as to leach the herbicide into the soil if rain does not fall within 24 hours. On deep-rooted perennials such as Bindweed, Bermuda Grass, Canada Thistle, and Quack Grass the heavier rate may be required.

SOIL TREATMENT FOR WEED CONTROL: Apply 1-2 lbs. per 100 square feet in early spring before weeds start to grow. This treatment may provide season long control

of seedlings on treated areas. Artificial watering may be necessary to leach the herbicide into the soil. Soil type, rainfall, and length of growing season will influence the residual control of seedlings.

PRECAUTIONS

This product is not selective in action and may be toxic to all types of vegetation and may render the entire treated area totally or partially unproductive for one or more years. Care should be taken to confine the application to the particular area intended to be treated. Do not apply within the drip line of shrubs and trees or on lawns and other desirable plants which are not intended to be destroyed or injured. This includes areas which may be underlaid by roots of adjacent valuable growths. Avoid careless application of this material, or washing by water run-off, to areas where desirable plants are growing or which will be used for later planting as this may result in injury to desirable plants. Do not contaminate water supplies.

CAUTIONS

Harmful if swallowed. Avoid contact with skin or clothing. Do not leave container where children or animals may take internally.

Avoid contamination of feed and foodstuffs. Do not store in direct contact with ground or concrete floors.

Do not reuse container. Destroy when empty.

7

18