

VIGORA

ACCEPTED

MAR 27 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 537-1827

5%

MALATHION DUST

**For Control of Many Insects That
Attack Ornamentals and Vegetables**

**CAUTION: KEEP OUT OF REACH OF CHILDREN.
See Back/Side Panel for Required Precautions.**

ACTIVE INGREDIENT:

*Malathion	5.00%
INERT INGREDIENTS	95.00%
TOTAL	100.00%

*0, O-Dimethyl Dithiophosphate of Diethyl Maltionphosphate

NET CONTENTS 4 LBS.

MANUFACTURED BY

Swift Agricultural Chemicals Company

CHICAGO, ILLINOIS 60604

U.S.D.A. REG. NO. 537-1827

CAUTION

Do not swallow. Avoid breathing dust. Avoid splashing on eyes. Avoid contact with skin. Wash thoroughly after use with soap and water. Avoid contact with animal food and foodstuffs. Keep out of reach of children.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not get on clothes. Clean water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

DIRECTIONS FOR USE

VIGORO 5% MALATHION DUST is to be used without dilution. Apply uniformly to foliage as for best results dust regularly at 7-10 day intervals. Apply preferably in early morning or late evening when the air is quiet, and cover all exposed parts of the plant with a thin, even film of dust.

For control of millipedes, apply dust to soil in garden beds, around plants, and around home in various gardens. Apply at the rate of 1 lb. per 100 sq. ft. to the soil surface. Repeat at 7-10 day intervals as needed. **DO NOT APPLY TO VEGETABLE CROPS OR TO GREENHOUSES WHERE FOOD CROPS ARE GROWN.**

ORNAMENTALS

For control of Aphids, Bagworms, Spider Mites on Ornamentals. See directions for use.

VEGETABLE CROPS	INSECTS	INTERVAL (Last Dust to Harvest)
<u>Asparagus</u>	Asparagus Beetle, Thrips	3 days
<u>Beans</u>	Aphids, Cucumber Beetles, Japanese Beetles, Leafhoppers, American Bean Beetle, Spider Mites	1 day
<u>Beets, Carrots</u>	Aphids, Leafhoppers	7 days
<u>Broccoli, Brussels Sprouts</u>	Aphids	7 days
<u>Cabbage, Cauliflower, Kale,</u> <u>Kohlrabi, Mustard Greens,</u> <u>Spinach, Swiss Chard, Cornards</u>		
<u>Cantaloup, Watermelon,</u> <u>Cucumber</u>	Aphids, Cucumber Beetles, Spider Mites, Fleahorn	1 day
<u>Celery</u>	Aphids, Spider Mites	7 days
<u>Eggplant</u>	Eggplant Tree Bug, Spider Mites, Aphids	3 days

<u>Asparagus</u>	Asparagus beetle, Thrips	3 days
<u>Beans</u>	Aphids, Cucumber Beetles, Japanese Beetles, Leafhoppers, Mexican Bean Beetle, Spider Mites	1 day
<u>Beets, Carrots</u>	Aphids, Leafhoppers	7 days
<u>Broccoli, Brussels Sprouts</u>	Aphids	7 days
<u>Cabbage, Cauliflower, Kale, Kohlrabi, Mustard Greens, Spinach, Swiss Chard, Collards</u>		
<u>Cantaloup, Watermelon, Cucumber</u>	Aphids, Cucumber Beetles, Spider Mites, Pickleworms	1 day
<u>Celery</u>	Aphids, Spider Mites	7 days
<u>Eggplant</u>	Eggplant Leaf Bug, Spider Mites, Aphids	3 days
<u>Lettuce, Endive (Escarole)</u>	Aphids, Leafhoppers, Cabbage Loopers, Mites	14 days leaf 7 days head and Endive
<u>Okra</u>	Aphids, Japanese Beetle	Do not apply After pods start
<u>Onions</u>	Onion Thrips, Onion Maggot	3 days
<u>Peas</u>	Aphids, Pea Weevil	3 days
<u>Peppers</u>	Aphids, Pepper Maggot	3 days
<u>Potatoes</u>	Aphids, False Chinch Bugs, Leafhoppers, Thrips, Mealybugs	0 day
<u>Pumpkin, Squash</u>	Aphids, Squash Vine Borer, Mites	3 days pumpkin 1 day squash
<u>Parsnips, Salsify, Horse Radishes, Radishes</u>	Aphids, Leafhoppers	7 days
<u>Sweet Potatoes</u>	Leafhoppers	0 day
<u>Tomatoes</u>	Aphids, Drosophila, Spider Mites, Tomato Pinch Mites	5 days

Dogs and Cats
Do not repeat application for 7 days and only if infestation is present.

Animal Quarters (Dog and Cat)

Dust with any garden type duster, puff duster or a shaker type dispenser. Dust thoroughly.

Dust in kennels, pens, yards, lawns and other areas at the rate of 1 or 2 lbs. per 1000 sq. ft. using a hand duster. Mammals and birds should be removed before dusting.