

VIGORO

FRUIT TREE SPRAY

APPLE - PEACH - PEAR
PLUM - CHERRY

NET WT. 16 OZ. (1 LB.)

VIGORO FRUIT TREE SPRAY

GENERAL DIRECTIONS

ONE POUND MAKES 12 GALLONS OF DILUTE SPRAY

Apply thoroughly to fruit tree foliage and growing fruit or berries at frequent intervals, 7 to 10 days, for maximum spray protection. On small trees 5 to 6 feet high, apply approximately $\frac{1}{4}$ to 1 quart of dilute spray per tree. Larger trees will take $\frac{1}{4}$ to 1 gallon and possibly more for complete coverage. Heavy foliage trees such as peaches may require additional spray due to their growing habit. For mixing small amounts add 5 tablespoons to 1 gallon. Use a sprayer that will give a fine spray mist to penetrate the foliage. With hand sprayers occasionally stir or shake the mixture to keep the particles in suspension.

APPLES AND PEARS: Controls insects such as codling moth, plum curculio, two spotted mites, European red mites, aphids, Forbes scale, apple maggots and pear slugs. Diseases such as apple scab, black rot, sooty blotch, Brooks spot. Make first application when at pink stage and before blossoms open. Do not spray during bloom to protect bees. Make second application when $\frac{1}{4}$ of the petals have fallen. Apply 10 days later and at 10 day intervals. If insect or disease problems arise during the season with normal 10 days sprayings reduce time to 5-7 days but in any case not later than 7 days before harvest.

PEACHES: Controls two spotted mite, oriental fruit moth (injures new growing tips), Japanese beetle and rose chater; diseases such as brown rot and peach scab. Make first application when petals are falling for peach scab spray also at full pink blossom stage. Again when shucks around the small fruit begin to split and fall, then at 10 day intervals but not later than 21 days before anticipated picking. If peach brown rot becomes a problem apply wettable sulphur to trees every 7 days at manufacturer's recommended spray rates.

PLUMS: Controls two spotted mites, plum curculio and Japanese beetles. Apply when petals are falling again when most of the shucks have fallen from the new fruit, 7 to 10 days later and at 10 day intervals up until 14 days of harvest.

CHERRIES: Controls insects such as plum curculio, leaf slugs, aphids and two spotted mites as well as diseases such as cherry leaf spot and brown rot. Apply when petals are falling and again when shucks around the small fruit begin to fall, then at 10 day intervals but not later than 7 days before harvest or anticipated picking. For leaf spot apply spray right after harvest to protect against premature leaf drop from cherry leaf spot.

RASPBERRIES, BLACKBERRIES: For spider mites, thrips, leafhoppers, Japanese beetles, spur blight and anthracnose. For insects apply when first seen and repeat as needed for control. For diseases apply when blossoms are in bud (young canes are 8 to 10 inches long). Make a second application 2 weeks later. Apply in fall when old canes are removed. Do not apply within 14 days of harvest.

STRAWBERRIES: Make first application as growth starts after uncovering plants. Re-spray as follows for most effective results: 1. Just after bloom. 2. 10 to 14 days following bloom period. 3. 2 weeks after harvest or as needed and again in fall. (Oct. 1st and Nov. 1st in Wisconsin and northern states.) Controls aphids, cyclamen mites, strawberry weevil, strawberry leafrollers, spittlebugs, spider mites, leaf spot and botrytis fruit rot. Do not apply within 14 days of harvest.

GRAPES: Controls black rot, leafhoppers, grape berry moth, grape leaf skeletonizer, Japanese beetles, spider mites and mealybugs. For insects apply when they first appear. Use 2 or more sprays before or after harvest. For black rot apply 2 sprays before bloom and one just after. Repeat making up to 3 sprays before bunches "close." Do not apply within 14 days of harvest.

CAUTION: Harmful if swallowed. Do not inhale dust or spray mist. Avoid contact with skin. Wash thoroughly after using. If eyes get medical attention. Keep out of reach of children. Do not contaminate feed, food, or food equipment.

Do not apply within 7 days of harvest on apples and pears; 7 days on cherries and plums; 21 days on peaches. Remove any residue on fruit after harvest by washing, wiping or other effective means.

Keep out of any body of water. This product will kill fish.

Do not apply when weather conditions favor drift of spray to other areas treated. Apply the product only as directed on this label.

Reuse equipment and container, and dispose of wastes by burying in deep land away from water supply. Containers should be disposed by punching holes in them and burying with waste, or by burning. Keep out of fire.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

ACTIVE INGREDIENT

Methoxychlor Technical

Methoxychlor 0.0 Dimethylphosphate of methoxychlor 1.0
Methoxychlor 1.0 Dimethylphosphate of methoxychlor 2.0

INERT INGREDIENTS

Methoxychlor 1.0 Dimethylphosphate of methoxychlor 2.0
Methoxychlor 1.0 Dimethylphosphate of methoxychlor 2.0

VIGORO®

FRUIT TREE SPRAY

APPLE - PEACH - PEAR
PLUM - CHERRY

NET WT. 16 OZ. (1 LB.)

VIGORO. FRUIT TREE SPRAY

GENERAL DIRECTIONS

ONE POUND MAKES 12 GALLONS OF DILUTE SPRAY

Apply thoroughly to fruit tree foliage and growing fruit or berries. Frequent intervals (7 to 10 days) for maximum spray protection. On small trees 5 to 6 feet high, apply approximately 1/2 to 1 quart of dilute spray per tree. Larger trees will take 1 to 1 gallon and possibly more for complete coverage. Heavy foliage trees such as peaches may require additional spray due to their growing habit. For mixing small amounts add 5 tablespoons to 1 gallon. Use a sprayer that will give a fine spray mist to penetrate the foliage. With hand sprayers, occasionally stir or shake the mixture to keep the particles in suspension.

APPLES AND PEARS: Controls insects such as codling moth, plum curculio, two spotted mites, European red mites, aphids, Forbes scale, apple maggots and pear slugs; Diseases such as apple scab, black rot, sooty blotch, Brooks spot. Make first application when at pink stage and before blossoms open. Do not spray during bloom to protect bees. Make second application when 1/4 of the petals have fallen. Apply 10 days later and at 10 day intervals. If insect or disease problems arise during the season with normal 10 days sprayings, reduce time to 5-7 days but in any case not later than 7 days before harvest.

PEACHES: Controls two spotted mite, oriental fruit moth (injures new growing tips), Japanese beetle and rose chafer; diseases such as brown rot and peach scab. Make first application when petals are falling (for peach scab spray also at full pink blossom stage). Again when shucks around the small fruit begin to split and fall, then at 10 day intervals but not later than 21 days before anticipated picking. If peach brown rot becomes a problem apply wettable sulphur to trees every 7 days at manufacturer's recommended spray rates.

PLUMS: Controls two spotted mites, plum curculio and Japanese beetles. Apply when petals are falling, again when most of the shucks have fallen from the new fruit, 7 to 10 days later and at 10 day intervals up until 14 days of harvest.

CHERRIES: Controls insects such as plum curculio, leaf slugs, aphids and two spotted mites as well as diseases such as cherry leaf spot and brown rot. Apply when petals are falling and again when shucks around the small fruit begin to fall, then at 10 day intervals but not later than 7 days before harvest or anticipated picking. For leaf spot apply spray right after harvest to protect against premature leaf drop from cherry leaf spot.

RASPBERRIES, BLACKBERRIES: For spider mites, thrips, leafhoppers, Japanese beetles, spur blight and anthracnose. For insects apply when first seen and repeat as needed for control. For diseases apply when blossoms are in bud (young canes are 8 to 10 inches long). Make a second application 2 weeks later. Apply in fall when old canes are removed. Do not apply within 14 days of harvest.

STRAWBERRIES: Make first application as growth starts after uncovering plants. Re-spray as follows for most effective results: 1: Just after bloom. 2: 10 to 14 days following bloom period. 3: 2 weeks after harvest or as needed and again in fall. Oct. 1st and Nov. 1st in Wisconsin and northern states. Controls aphids, cyclamen mites, strawberry weevil, strawberry leafrollers, spittlebugs, spider mites, leaf spot and botrytis fruit rot. Do not apply within 14 days of harvest.

GRAPES: Controls black rot, leafhoppers, grape berry moth, grape leaf skeletonizer, Japanese beetles, spider mites and mealybugs. For insects apply when they first appear. Use 2 or more sprays before or after harvest. For black rot apply 2 sprays before bloom and one just after. Repeat making up to 3 sprays before bunches "close." Do not apply within 14 days of harvest.

CAUTION: Harmful if swallowed. Do not inhale dust or spray mist. Avoid contact with skin. Wash thoroughly after use. If swallowed get medical attention. Keep out of reach of children. Do not contaminate feed, food, or food containers.

Do not apply within 7 days of harvest on apples and pears, 7 days on cherries and plums, 14 days on peaches. Remove all visible fruit after harvest by washing, wiping or other effective means.

Keep out of any body of water. This product will kill fish.

Do not apply when weather conditions favor drift of spray, dust from areas treated. Apply this product only as specified on this label.

Rinse equipment and containers and dispose of wastes by burying in soil, or placing away from water, or by incineration. Do not burn. Do not puncture by punching holes in them and burying with waste, or by burning. Keep out of fire.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. No conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary. Seller makes no claims other than those stated on this label.

ACTIVE INGREDIENT

Malathion, Technical

Malathion, 0.0 D. Malathion, 0.0 D. Malathion, 0.0 D.

Malathion, 0.0 D. Malathion, 0.0 D. Malathion, 0.0 D.

INERT INGREDIENTS