


WEED CONTROL
PLUS
LAWN FERTILIZER



GOLDEN



net wt 20 lbs
TREATS
5000 sq ft
24-4-8


GOLDEN VIGORO WEED CONTROL PLUS LAWN FERTILIZER

GOLDEN VIGORO

WEED CONTROL PLUS LAWN FERTILIZER


WEED AND FEED

**Feeds lawn as
3-way action
kills chickweed,
dandelion, knotweed,
plantains, henbit,
spurge and many
other weeds**



ACTIVE INGREDIENTS	
2,4-D (2,4-dichlorophenoxy)acetic acid	0.73
2,4,5-T (2,4,5-trichlorophenoxy)acetic acid	0.01
Diazinon (O,O-dimethyl phosphorodithioic acid S-(1-methyl-2-pyridinyl)ethyl ester)	0.27
INERT INGREDIENTS	98.99
EPA REG. NO. 274	100.00

CAUTION: Keep out of reach of children. See back panel for additional cautions.



WEED CONTROL PLUS LAWN FERTILIZER

net wt 20 lbs

GOLDEN VIGORO

WEED CONTROL PLUS LAWN FERTILIZER

WEED & FEED

24-4-8

*GUARANTEED ANALYSIS

TOTAL NITROGEN (N) 24.0%
 10% Ammoniacal Nitrogen (derived from ammoniated phosphates)
 8.0% Water Insoluble Nitrogen (derived from ureaformaldehyde and Isobutylidene Diurea)
 15.0% Water Soluble Nitrogen (derived from urea, ureaformaldehyde and Isobutylidene Diurea)

AVAILABLE PHOSPHORIC ACID (P₂O₅) (derived from ammoniated phosphates) 4.0%
 SOLUBLE POTASH (K₂O) (derived from muriate of potash) 8.0%

Potential acidity 950 lbs calcium carbonate equivalent per ton. *Minimum guaranteed analysis in Michigan. Manufactured by Swift Agricultural Chemicals Corporation, Chicago, Illinois 60604

MONEY BACK PROMISE

You must be pleased with results or you are entitled to your money back. Simply send sales slip to Vigoro, Chicago, Illinois 60604 for prompt refund.

EASY TO USE

WHEN TO APPLY: For best results apply when weeds are actively growing in the spring or early fall after heavy rains. Apply while lawn is wet with dew or after sprinkling lightly. To get maximum weed killing results, do not rake, mow or water the lawn for at least 24 hours after applying.

WHERE TO APPLY: This product may be used on bluegrass, fescue, rye, Bermuda and zoysia lawns. If adverse low moisture conditions exist, weeds are in a poor state of growth or the lawn is heavily overrun with weeds, more than one application may be required. Apply the second application in 30 days.

HOW TO APPLY: Using the spreader settings provided below, apply this product to 5000 square feet of lawn area (100' x 50').

WEEDS CONTROLLED: The list below is a partial list of the common weeds controlled by this product.

Beetroot	Dandelion	Lespedeza	Poison Ivy	Spurge
Black Medick	Dock	Yarrow	Poison Oak	Thistle
Buckhorn	Ground Ivy	Morning Glory	Purslane	Wild Carrot
Burdock	Heavenly	Oxalis	Ragweed	Wild Garlic
Chickweed	Henbit	Peppercorn	Sheep Sorrel	Wild Lettuce
Chickweed	Knotweed	Pigweed	Shepherd's Purse	Wild Onion
Clover	Lambsquarters	Plantain	Speedwell	Yarrow

CAUTION

Harmful if swallowed

Wash after handling

Avoid breathing dust. Avoid contact with eyes. Avoid contact with skin. Avoid contact with clothing. Avoid contact with water. Do not use in areas where children or pets play. Do not use on bent grass, St. Augustine grass, or Dichondra. Lawn injury may result. Do not use this product where desirable clovers are present or allow the product to wind drift on flowers, shrubs, vegetables or ornamental shrubbery. Do not apply to newly seeded lawns. During application, close spreader opening when not in use. Apply when wind is calm. Do not use on lawns treated with insecticides or fungicides. Do not use on lawns treated with herbicides. Do not apply to areas underlaid by roots of desirable trees and shrubs. Do not use container. Destroy by burying in a safe place. Do not burn. Do not contaminate water used for domestic purposes.

HOW TO SET YOUR SPREADER

Cylinder	3%	Sears	5
K-Mat	2%	Sears	5
Lawn Broadcast	8%	Ward	5

NOTE: Use the above settings with a broadcast spreader on a lawn area

ED CONTROL
IN FERTILIZER

GOLDEN
VISCOSO
WEED CO

WEED
AND
FEE

Feeds lawn
3-way action
kills chickweed,
dandelion, l
plantains, h
spurge and
other weed

GOLDEN VISCOSO WEED CO
PLUS
LAWN FEE


GOLDEN VIGORO WEED CONTROL PLUS LAWN FERTILIZER

GOLDEN VIGORO

WEED CONTROL PLUS LAWN FERTILIZER

WEED AND FEED

Feeds lawn as 3-way action kills chickweed, dandelion, knotweed, plantains, henbit, spurge and many other weeds



ACTIVE INGREDIENTS:	
2,4-dichlorophenoxyacetic acid	0.73%
2-(2-methyl-4-chlorophenoxy) propionic acid	0.31%
Dicarb.a (3,6-dichloro-o-anisic acid)	0.07%
INERT INGREDIENTS:	
	98.89%
E.P.A. Reg. No. 557-1845	Total 100.00%

CAUTION: Keep out of reach of children. See back panel for additional cautions.

GOLDEN VIGORO

WEED CONTROL PLUS LAWN FERTILIZER

net wt 20 lbs

GOLDEN VIGORO

WEED CONTROL PLUS LAWN FERTILIZER

WEED & FEED

24-4-8

*GUARANTEED ANALYSIS

TOTAL NITROGEN (N) 24.0%

1.0% Ammoniacal Nitrogen (derived from ammoniated phosphates)

8.0% Water Insoluble Nitrogen (derived from ureaformaldehyde and Isobutylidene Diurea)

15.0% Water Soluble Nitrogen (derived from urea, ureaformaldehyde and Isobutylidene Diurea)

AVAILABLE PHOSPHORIC ACID (P₂O₅) (derived from ammoniated phosphates) 4.0%

SOLUBLE POTASH (K₂O) (derived from muriate of potash) 8.0%

Potential acidity 950 lbs calcium carbonate equivalent per ton. *Minimum guaranteed analysis in Michigan. Manufactured by Swift Agricultural Chemicals Corporation, Chicago, Illinois 60604.

MONEY BACK PROMISE

You must be pleased with results or you are entitled to your money back. Simply send sales slip to Vigoro, Chicago, Illinois 60604 for prompt refund.

EASY TO USE

WHEN TO APPLY—For best results apply when weeds are actively growing in the spring or early fall after heavy rains. Apply while lawn is wet with dew or after sprinkling lightly. To get maximum weed killing results, do not rake, mow or water the lawn for at least 24 hours after applying.

WHERE TO APPLY—This product may be used on bluegrass, fescue, rye, Bermuda, and zoysia lawns. If adverse low moisture conditions exist, weeds are in a poor state of growth, or the lawn is heavily overrun with weeds, more than one application may be required. Apply the second application in 30 days.

HOW TO APPLY—Using the spreader settings provided below, apply this product to 5000 square feet of lawn area (100 x 50).

WEEDS CONTROLLED—The list below is a partial list of the common weeds controlled by this product

Bedstraw	Dandelion	Lespedeza	Poison Ivy	Scourge
Black Medic	Dock	Mallow	Poison Oak	Thistle
Buckhorn	Ground Ivy	Morning Glory	Purslane	Wild Carrot
Burdock	Heal-all	Oxalis	Ragweed	Wild Garlic
Chicory	Henbit	Peppergrass	Sheep Sorrel	Wild Lettuce
Chickweed	Knotweed	Pigweed	Shepherd's Purse	Wild Onion
Clover	Lambsquarter	Plantains	Speedwell	Yarrow

CAUTION

*GUARANTEED ANALYSIS

TOTAL NITROGEN (N) 24.0%
 1.0% Ammoniacal Nitrogen (derived from ammoniated phosphates)
 8.0% Water Insoluble Nitrogen (derived from ureaformaldehyde and Isobutylidene Diurea)
 15.0% Water Soluble Nitrogen (derived from urea, ureaformaldehyde and Isobutylidene Diurea)

AVAILABLE PHOSPHORIC ACID (P₂O₅) (derived from ammoniated phosphates) 4.0%
 SOLUBLE POTASH (K₂O) (derived from muriate of potash) 8.0%

Potential acidity 950 lbs calcium carbonate equivalent per ton *Minimum guaranteed analysis in Michigan. Manufactured by Swift Agricultural Chemicals Corporation, Chicago, Illinois 60604.

MONEY BACK PROMISE

You must be pleased with results or you are entitled to your money back. Simply send sales slip to Vigoro, Chicago, Illinois 60604 for prompt refund.

EASY TO USE

WHEN TO APPLY - For best results apply when weeds are actively growing in the spring or early fall after heavy rains. Apply while lawn is wet with dew or after sprinkling lightly. To get maximum weed killing results, do not rake, mow or water the lawn for at least 24 hours after applying.

WHERE TO APPLY - This product may be used on bluegrass, fescue, rye, Bermuda, and zoysia lawns. If adverse low moisture conditions exist, weeds are in a poor state of growth, or the lawn is heavily overrun with weeds, more than one application may be required. Apply the second application in 30 days.

HOW TO APPLY - Using the spreader settings provided below, apply this product to 5000 square feet of lawn area (100 x 50).

WEEDS CONTROLLED - The list below is a partial list of the common weeds controlled by this product.

Bedstraw	Dandelion	Lespedeza	Poison Ivy	Scurge
Black Medic	Dock	Mallow	Poison Oak	Thistle
Buckhorn	Ground Ivy	Morning Glory	Purslane	Wild Carrot
Burdock	Heal-all	Oxalis	Ragweed	Wild Garlic
Chicory	Henbit	Peppergrass	Sheep Sorrel	Wild Lettuce
Chickweed	Knotweed	Pigweed	Shepherd's Purse	Wild Onion
Clover	Lambsquarter	Plantains	Speedwell	Yarrow

CAUTION

Harmful if swallowed.

Wash after handling.

Avoid inhaling dust or contact with skin, eyes and clothing.

In case of contact, flush with plenty of water. For eyes, get medical attention.

Not recommended for use on bent grass, St. Augustine grass, or Dichandra Lawns. Its injury may result. Do not use this product where desirable clovers are present or allow the product to "wind drift" on flowers, shrubs, vegetables or ornamental shrubbery. Do not apply to newly seeded lawns. During application, close spreader openings when turning. Apply when air is calm. Keep away from fertilizers, seeds, insecticides or fungicides. Do not store near feed or foodstuffs. Do not apply to areas underlaid by roots of desirable trees and shrubs. Do not reuse container. Destroy by burying in a safe place. Do not burn. Do not contaminate water used for domestic purposes.

HOW TO SET YOUR SPREADER

Cyclone	3 1/2	Scotts	5
K-Mart	2 1/2	Sears	5
Lawn Beauty	8 1/2	Wards	5

NOTE: Spreader rate varies with walking speed. Check spreader over known area.

WEED CONTROL
PLUS
LAWN FERTILIZER

GOLDEN VIGORO
557-1845

ACCEPTED
5/27/845
APR 10 1972
UNDER THE FEDERAL INSECT,
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PESTS REGISTERED
ED UNDER NO. 557-1845
SUBJECT TO ATTACHED CONDITIONS

net wt 20 lbs
TREATS
5000 sq ft
24-4-8

GOLDEN VIGORO
WEED CONTROL
PLUS
LAWN FERTILIZER

GOLDEN VIGORO

WEED CONTROL PLUS LAWN FERTILIZER

WEED AND FEED

Feeds lawn as
3-way action
kills chickweed,
dandelion, knotweed,
plantains, henbit,
spurge and many
other weeds



ACTIVE INGREDIENTS:
2,4-dichlorophenoxyacetic acid 0.73%
2-(2-methoxy-4-chlorophenoxy) propionic acid 0.31%
Dicamba (3,6-dichloro-o-anisic acid) 0.07%
INERT INGREDIENTS: 98.89%
E.P.A. Reg No. 557-1845 Total 100.00%

CAUTION: Keep out of reach of children. See back panel for additional cautions.

GOLDEN VIGORO
WEED CONTROL
PLUS
LAWN FERTILIZER

net wt 20 lbs

57-184

GOLDEN VIGORO

WEED CONTROL PLUS LAWN FERTILIZER

WEED & FEED

24-4-8

*GUARANTEED ANALYSIS

TOTAL NITROGEN (N) 24.0%
 1.0% Ammoniacal Nitrogen (derived from ammoniated phosphates)
 8.0% Water Insoluble Nitrogen (derived from ureaformaldehyde and isobutylidene Diurea)
 15.0% Water Soluble Nitrogen (derived from urea, ureaformaldehyde and isobutylidene Diurea)

AVAILABLE PHOSPHORIC ACID (P₂O₅) (derived from ammoniated phosphates) 4.0%
 SOLUBLE POTASH (K₂O) (derived from muriate of potash) 8.0%

Potential acidity 950 lbs calcium carbonate equivalent per ton. *Minimum guaranteed analysis in Michigan. Manufactured by Swift Agricultural Chemicals Corporation, Chicago, Illinois 60604.

MONEY BACK PROMISE

You must be pleased with results or you are entitled to your money back. Simply send sales slip to Vigoro, Chicago, Illinois 60604 for prompt refund.

EASY TO USE

WHEN TO APPLY—For best results apply when weeds are actively growing in the spring or early fall after heavy rains. Apply while lawn is wet with dew or after sprinkling lightly. To get maximum weed killing results, do not rake, mow or water the lawn for at least 24 hours after applying.

WHERE TO APPLY—This product may be used on bluegrass, fescue, rye, Bermuda, and zoysia lawns. If adverse low moisture conditions exist, weeds are in a poor state of growth, or the lawn is heavily overrun with weeds, more than one application may be required. Apply the second application in 30 days.

HOW TO APPLY—Using the spreader settings provided below, apply this product to 5000 square feet of lawn area (100 x 50).

WEEDS CONTROLLED—The list below is a partial list of the common weeds controlled by this product.

Bedstraw	Dandelion	Lespedeza	Poison Ivy	Spurge
Black Medic	Dock	Mallow	Poison Oak	Thistle
Buckhorn	Ground Ivy	Morning Glory	Purslane	Wild Carrot
Burdock	Heart-all	Oxalis	Ragweed	Wild Garlic
Chirory	Henbit	Peppergrass	Sheep Sorrel	Wild Lettuce
Chickweed	Knockweed	Pigweed	Shepherd's Purse	Wild Onion
Clover	Lambsquarter	Plantains	Speedwell	Yarrow

CAUTION

Harmful if swallowed.
 Wash after handling.
 Avoid inhaling dust or contact with skin, eyes and clothing.
 In case of contact, flush with plenty of water. For eyes, get medical attention.
 Not recommended for use on bent grass, St. Augustine grass, or Dichondra Lawns. Its injury may result. Do not use this product where desirable clovers are present or allow the product to "wind drift" on flowers, shrubs, vegetables or ornamental shrubbery. Do not apply to newly seeded lawns. During application, close spreader openings when turning. Apply when air is calm. Keep away from fertilizers, seeds, insecticides or fungicides. Do not store near feed or foodstuffs. Do not apply to areas underlaid by roots of desirable trees and shrubs. Do not reuse container. Destroy by burying in a safe place. Do not burn. Do not contaminate water used for domestic purposes.

HOW TO SET YOUR SPREADER

Cyclone.....	3 1/2	Scot's.....	5
K-Mart.....	2 1/2	Sea 6.....	5
Lawn Beauty.....	8 1/2	Wal.....	5

NOTE: Spreader rate varies with walking speed check spreader over known area.