

VIGORO

VIGORO GARDEN FUNGICIDE

*Apply before diseases occur or appear and repeat at 7 to 10-day intervals as needed.

Crop and Diseases	Amount to Use Per Gallon of Water	SPECIAL INSTRUCTIONS When to Apply
<u>Chrysanthemums</u> Powdery Mildew	1 1/2 Tbs.	Apply weekly during periods of rapid new growth and severe infection; twice weekly during continuous rains or high humidity.
<u>Roses</u> Black Spot, Powdery Mildew	1 1/2 Tbs.	Apply when leaves first appear. Repeat at 7 to 10-day intervals or as necessary throughout season.
<u>Zinnias</u> Powdery Mildew Leaf Spot	1 1/2 Tbs.	Apply weekly during periods of rapid growth and severe infection and twice weekly during continuous rains or high humidity.
<u>Aster, Phlox,</u> Sporangium Powdery Mildew	1 1/2 Tbs.	*Same as above.
<u>Citrus</u> Citrus Scab	8 Tbs. per 10 Gals. of Water.	Apply first application in dormant period before bloom. Make second application when 1/2 of the petals have fallen.
<u>Citrus</u> Melanose	8 Tbs. per 10 Gals. of Water.	Apply one week following bloom beginning in February and ending in April; complete as soon as possible. Do not apply to trees bearing mature fruit which will be harvested during the current season. Do not apply during off-season bloom period.
<u>Cucumbers,</u> <u>Cantaloupes,</u> <u>Honeydew Melons,</u> <u>Muskmelons,</u> <u>Pumpkins, Squash</u> (Summer & Winter) <u>Watermelons</u> <u>Garlic, Onions</u> Downy Mildew, Powdery Mildew	1 1/2 Tbs. per Gal. of Water.	Apply 2 quarts per 100 ft. of row. On garlic and onions, use 1 quart per 100 ft. of row.
<u>Tomatoes</u> Anthracnose	1 1/2 Tbs. per Gal. of Water	Apply 2 quarts per 100 ft. of row.

DIRECTIONS FOR USE

Mix the required amount of VIGORO GARDEN FUNGICIDE in a little water to make a smooth thin paste, then add required amount of water. Keep mixture agitated while spraying. Do not allow mixture to stand. Do not use a combination of wettable powders with emulsifiable liquids unless previous use of such mixtures has proved them to be physically compatible. Spray all parts of plants thoroughly.

CAUTION: Avoid inhalation of dust or spray mist. Avoid contact with skin. Wash hands after use. Avoid contamination of feed and food-stuffs. Dispose of empty container promptly. Keep out of reach of children.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Apply this product only as specified on this label.

ACTIVE INGREDIENT:
Folpet N - (trichloromethylthio) phthalamide 75.9%
INERT INGREDIENT 24.1%
Total 100.0%
U. S. Patent Nos. 2,553,770; 2,553,771; 2,553,776.

GARDEN FUNGICIDE

FOR FLOWERS
ROSES, CITRUS,
VEGETABLES

NET WT. 6 OZ.

ACCEPTED

MAR 28 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 657-1840

CAUTION: Keep out of reach of children
See other panel for additional caution

MANUFACTURED BY

EMMET ABBOTT LABORATORIES CHEMICALS CORPORATION
CHICAGO, ILLINOIS 60604
E. A. B. 557 1843 AA

