

net wt 25 lbs
TREATS
5000^{sq}_{ft}
18-4-8

GOLDEN **VIGORO**

WEED CONTROL PLUS LAWN FERTILIZER

**WEED
AND
FEED**

ACCEPTED
AUG 6 1979
Under the Federal Insecticide,
Fungicide and Rodenticide Act,
this product has been found to
be effective for the control of
weeds and is registered under
EPA Reg. No. 577-17-1000

Feeds lawn as
3-way action
kills chickweed,
dandelion, knotweed,
plantains, henbit,
spurge and many
other weeds

ACCEPTED
AUG 6 1979
Under the Federal Insecticide,
Fungicide and Rodenticide Act,
this product has been found to
be effective for the control of
weeds and is registered under
EPA Reg. No. 577-17-1000



CAUTION Keep out of reach of children. See back panel for additional cautions

VIGORO **WEED CONTROL
PLUS
LAWN FERTILIZER**

U

WEED CONTROL PLUS LAWN FERTILIZER

18-4.8

• **Aspirin** is a common analgesic and antipyretic.

- $$\begin{aligned} \text{a. } \frac{1}{2} \cdot \frac{1}{3} &= \frac{1}{6} \\ \text{b. } \frac{1}{2} \cdot \frac{1}{2} &= \frac{1}{4} \\ \text{c. } \frac{1}{2} \cdot \frac{1}{4} &= \frac{1}{8} \end{aligned}$$

[illegible]

S. J. Lieberman

Margaret Joseph, SWA Queensland member, presented the following paper:

[illegible]

WHERE TO APPLY: To get a copy of the application form, contact the American Society of Human Resources, 1000 17th St., N.W., Washington, D.C. 20036. The form is available for \$10.00. You will also receive a copy of the ASHR directory, which lists the names and addresses of 1,000 employers who are members of the society. You will also receive a copy of the ASHR directory, which lists the names and addresses of 1,000 employers who are members of the society.

TABLE 1. *Summary of the 1996-1997 season for the 1000-hour program. The number of hours of instruction for each of the 1000-hour program participants is shown. The number of participants who completed the program is also shown. The number of participants who did not complete the program is also shown.*

[illegible]

As a consequence, the following theorem holds:

Theorem 1. *Let $\mathcal{A}$ be a linear algebra and let $\mathcal{B}$ be a linear algebra over $\mathcal{A}$. Then, the following conditions are equivalent:*

- $\mathcal{B}$ is a linear algebra over $\mathcal{A}$.*
- $\mathcal{B}$ is a linear algebra over $\mathcal{A}$.*
- $\mathcal{B}$ is a linear algebra over $\mathcal{A}$.*
- $\mathcal{B}$ is a linear algebra over $\mathcal{A}$.*

very low (≤ 1) BDI of the seed bank in the soil samples collected in the degraded area. Although the BDI of the seed bank in the degraded area was low, the seeds present in the seed bank of the degraded area were not lost. The BDI of the seed bank in the degraded area was low because the seeds of the species that were present in the seed bank of the degraded area were not viable. Despite the low BDI of the seed bank in the degraded area, the seeds of the species that were present in the seed bank of the degraded area were not lost.

[illegible]

NOTE: This document contains information that is exempt from public release under the Freedom of Information Act, 5 U.S.C. 552, and is being furnished to you for your information only. It is not to be distributed outside your agency.

net wt 25 lbs
TREATS
5000^{sq}_{ft}
18-4-8

GOLDEN VIGORO

WEED CONTROL PLUS LAWN FERTILIZER

**WEED
AND
FEED**

ACCEPTED
AUG 6 1979
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the use of this
product under 557-1850
EPA Reg. No.

Feeds lawn as
3-way action
kills chickweed,
dandelion, knotweed,
plantains, henbit,
spurge and many
other weeds

ACTIVE INGREDIENTS

*Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	0.69%
**Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy)propionic acid	0.31
***Dimethylamine salt of Dicamba (3,3-dichloro-o-anisic acid)	0.07
	98.93

INERT INGREDIENTS

EPA Reg No 557-1850	Total	100.00%
*Equivalent to 2,4-dichlorophenoxyacetic acid		0.57%
**Equivalent to 2-(2-methyl-4-chlorophenoxy)propionic acid		0.26
***Equivalent to 3,6-dichloro-o-anisic acid		0.06%

EPA Est No 557-IL-1

CAUTION: Keep out of reach of children. See back panel for additional cautions.



GOLDEN

VIGORO **WEED CONTROL
PLUS
LAWN FERTILIZER**

net wt 25 lbs

GOLDEN VIGORO

WEED CONTROL PLUS LAWN FERTILIZER
WEED & FEED

18-4-8

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	18.0
0.6% Ammoniacal Nitrogen	
5.0% Water Insoluble Nitrogen	
12.4% Water Soluble Nitrogen	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)	4.0
SOLUBLE POTASH (K ₂ O)	8.0

Derived from ammoniated phosphates, ureaformaldehyde, isobutylidene diurea, urea and muriate of potash. Potential Acidity 600 lbs. Calcium Carbonate Equivalent per ton. Manufactured by Swift Agricultural Chemicals Corporation, Chicago, Illinois 60602.

EASY TO USE

WHEN TO APPLY—For best results apply when weeds are actively growing in the spring or early fall after heavy rains. Apply while lawn is wet with dew or after sprinkling lightly. To get maximum weed killing results, do not rake, mow or water the lawn for at least 24 hours after applying.

WHERE TO APPLY—This product may be used on bluegrass, fescue, rye, Bermuda and zoysia lawns. If adverse low moisture conditions exist, weeds are in a poor state of growth, or the lawn is heavily overrun with weeds, more than one application may be required. Apply the second application in 30 days.

HOW TO APPLY—Using the spreader settings provided below, apply this product to 5000 square feet of lawn area (100 x 50).

WEEDS CONTROLLED—The list below is a partial list of the common weeds controlled by this product.

Bedstraw	Dandelion	Lespedeza	Poison Ivy	Sage
Black Medic	Dock	Mallow	Poison Oak	Thistle
Buckhorn	Ground Ivy	Morning Glory	Purslane	Wild Carrot
Burdock	Heat-all	Oxalis	Ragweed	Wild Garlic
Chicory	Henbit	Peppergrass	Sheep Sorrel	Wild Lettuce
Chickweed	Knotweed	Pigweed	Shepherd's Purse	Wild Onion
Clover	Lambsquarter	Plantain	Smartweed	Yarrow

CAUTION

Harmful if swallowed.

Wash after handling.

Avoid inhaling dust or contact with skin, eyes and clothing.

In case of contact, flush with plenty of water. For eyes, get medical attention.

Not recommended for use on bent grass, St. Augustine grass or Dichondra lawns, as injury may result. Do not use this product where desirable clovers are present or allow the product to "wind drift" on flowers, shrubs, vegetables or ornamental shrubbery. Do not apply to newly seeded lawns. During application, close spreader openings when turning. Apply when air is calm. Keep away from fertilizers, seeds, insecticides or fungicides. Do not store near feed or foodstuffs. Do not apply to areas underlain by roots of desirable trees and shrubs. Do not reuse container. Destroy by burying in a safe place. Do not burn. Do not contaminate water used for domestic purposes.

SPREADER SETTINGS

SPREADER	WIDTH	SETTING
Scott's Professional R-7	9 ft.	G-H
Scott's Rotary C-9	6 ft.	C
Central Roto-Spread	7 ft.	3½
Cyclone	6 ft.	3½
Scott's Drop	—	3½
Lawn Beauty	—	7½
Sears	—	6½
Wards	—	6½
K-Mart	—	6½

NOTE: Spreader rate varies with walking speed. To insure proper coverage, check spreader over known area.