

NET WT 25 LBS

TREATS

5000^{SQ.} FT.

18-4-8

GOLDEN

VIGORO

WEED CONTROL PLUS LAWN FERTILIZER

WEED AND FEED

Feeds lawn as 3-way action
kills chickweed, dandelion,
knotweed, plantains,
henbit, spurge and many
other weeds

ACTIVE INGREDIENTS

INERT INGREDIENTS



USE AS DIRECTED
Is comparable with RR-Mobies 92-2.
See label for labeling dated

KEEP OUT OF REACH OF CHILDREN
CAUTION

GUARANTEED ANALYSIS

Formaldehyde monochloride spates urea ureaformaldehyde isobutylidene
urea formaldehyde isobutyl Penta-acidity 600 lbs Calcium Carbonate
Epoxy Resin 2 Methylolol, Est-on Inc Chicago Illinois 60602


DISPOSAL

Front Panel

Net Wt 25 lbs
TREATS
5,000 sq. ft.
18-4-8

GOLDEN
VIGORO
WEED CONTROL PLUS LAWN FERTILIZER
WEED
and
FEED

Feeds lawn as
3-way action
kills chickweed,
dandelion, knotweed,
plantains, henbit,
spurge and many
other weeds



KEEP OUT OF REACH OF CHILDREN

CAUTION

In case of contact, wash skin with plenty
of soap and water; for eyes, flush with
water for 15 minutes and get medical
attention.
See back panel for additional precautionary
statements.

ACTIVE INGREDIENTS:

+*Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	0.65%
**Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	0.31%
***Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	0.07%

INERT INGREDIENTS	98.97%
-------------------------	--------

E.P.A. Reg. No. 557-1850	Total	100.00%
--------------------------	-------------	---------

+*Equivalent to 2,4-dichlorophenoxyacetic acid	0.54%
**Equivalent to 2-(2-methyl-4-chlorophenoxy) propionic acid	0.26%
***Equivalent to 3,6-dichloro-o-anisic acid	0.06%
+Isomer Specific by AOAC Method 6.D01.5	

E.P.A. Est. No. 557-CA-1

Back Panel

Net Wt 25 lbs.

GOLDEN

VIGORO

WEED CONTROL PLUS LAWN FERTILIZER

WEED & FEED

18-4-8

GUARANTEED ANALYSIS

TOTAL NITROGEN (N) 18.0%

0.6% Ammoniacal Nitrogen

12.4% Water Soluble Urea Nitrogen

5.0% Water Insoluble Nitrogen

AVAILABLE PHOSPHORIC ACID (P_2O_5) 4.0%

SOLUBLE POTASH (K_2O) 8.0%

Derived from ammoniated phosphates, urea, ureaformaldehyde, isobutylidene diurea and muriate of potash. Potential acidity 600 lbs. Calcium Carbonate Equivalent per ton. Manufactured by Estech, Inc., Chicago, Illinois 60602.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly after handling. Harmful if swallowed.

ENVIRONMENTAL HAZARD

Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate any body of water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

~~It is a violation of Federal law to use this product in a manner inconsistent with its labeling.~~

WHEN TO APPLY - For best results apply when weeds are actively growing in the spring or early fall after heavy rains. Apply while lawn is wet with dew or after sprinkling lightly. To get maximum weed killing results, do not rake, mow or water the lawn for at least 24 hours after applying.

Back Panel (contd.)

WHERE TO APPLY - This product may be used on bluegrass, fescue, rye, Bermuda, and zoysia lawns. If adverse low moisture conditions exist, weeds are in a poor state of growth, or the lawn is heavily overrun with weeds, more than one application may be required. Apply the second application in 30 days.

HOW TO APPLY - Using the spreader settings provided below, apply this product to 5,000 square feet of lawn area (100 X 50).

WEEDS CONTROLLED - The list below is a partial list of the common weeds controlled by this product.

Bedstraw	Dandelion	Lespedeza	Poison Ivy	Spurge
Black Medic	Dock	Mallow	Poison Oak	Thistle
Buckhorn	Ground Ivy	Morning Glory	Purslane	Wild Carrot
Burdock	Heal-all	Oxalis	Ragweed	Wild Garlic
Chicory	Henbit	Peppergrass	Sheep Sorrel	Wild Lettuce
Chickweed	Knotweed	Pigweed	Shepherd's Purse	Wild Onion
Clover	Lambsquarter	Plantains	Speedwell	Yarrow

CAUTION: Not recommended for use on bent grass, St. Augustine grass or Dichondra lawns, as injury may result. Do not use this product where desirable clovers are present or allow the product to "wind drift" on flowers, shrubs, vegetables or ornamental shrubbery. Do not apply to newly seeded lawns. During application close spreader openings when turning. Apply when air is calm. Do not apply to areas underlaid by roots of desirable trees and shrubs.

SPREADER SETTINGS

SPREADER	WIDTH	SETTING
Scott's Professional R-7	9 ft.	G-H
Scott's Rotary C-9	6 ft.	C
Central "Roto-Spred"	7 ft.	3½
Cyclone	6 ft.	3-5/8
Scott's Drop	-	3½
Lawn Beauty	-	7½
Sears	-	6-3/4
Wards	-	6½
Kmart	-	6½

NOTE: Spreader rate varies with walking speed. To insure proper coverage, ~~check spreader over known area.~~

STORAGE AND DISPOSAL

STORAGE: Store in a dry place. Do not contaminate feed and foodstuffs, seeds, fertilizers or other pesticides.

DISPOSAL: Do not reuse bag. Discard bag in trash.