

GOLDEN
VIGORO
WEED CONTROL PLUS LAWN FERTILIZER
WEED AND FEED

**WEED CONTROL
PLUS
FERTILIZER**

五、

GOLDEN VIGORO

WEED & FEED

18-4-8

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	18.0%
0.6% Ammoniacal Nitrogen	
12.4% Water Soluble Urea Nitrogen	
5.0% Water Insoluble Nitrogen	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)	4.0%
SOLUBLE POTASH (K ₂ O)	8.0%

Derived from ammoniated phosphates, urea, ureaformaldehyde, isobutyridene diurea and muriate of potash. Potential acidity 600 lbs. Calcium Carbonate Equivalent per ton. Manufactured by Estech Inc., Chicago, Illinois 60602.

ENVIRONMENTAL HAZARD
Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate any body of water by cleaning of equipment or disposal of wastes.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

WHEN TO APPLY—For best results apply when weeds are actively growing in the soil or early

words are actively growing in the spring or early fall after heavy rains. Water lawn one day before

application to ensure adequate soil moisture for root growth. Do not apply to wet grass. To get maximum weed killing results, do not mow or water away again until two days after applying. Do not apply during excessively hot weather (85°F or above). Repeat treatment if necessary to control any regrowth.

in bluegrass, fescue, rye, Bermuda, and ryegrass. If adverse low moisture conditions exist and weeds are in a poor state of growth, the lawn is heavily overtopped with weeds, more than one application may be required. Apply the herbicide application in 30 days.

provided below. Apply this product to a 100 square foot lawn area. 20 x 50.

Home State	Home Telephone	Home Address
State of Maine	1-207-555-1234	123 Main St., Portland, ME 04101

[illegible]

beet grass, St. Augustine grass or Dichondra
varies as injury may result. Do not use in spring
and where desirable ciphers are present or allow
the product to wind to it. On flowers, shrubs,
vegetables or ornamental shrubbery. Do not
apply to newly seeded lawns. During application
use a spreader for lawns when turning. Apply
when air is calm. Do not apply to trees, vines,
and shrubs. Use on citrus trees and shrubs.

[illegible][illegible]

NOTE: These rates vary with the age of the child. For example, the rate for a 10-year-old child is 10%.

Storage: Store in original container in a safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, livestock feed. Return spillage to original container or to disposal.

Product Disposal: See above with regard to disposal of several years' product that is no longer in use.

knotweed, plantains,
henbit, spurge and many
other weeds

NOV 28 1984
EPA Reg. No. 557-1850

ACTIVE INGREDIENTS:

++Dimethylamine salt of 2,4-dichlorophenoxy- acetic acid	0.65%
**Dimethylamine salt of 2-(2-methyl-4-chloro- phenoxy) propionic acid	0.31%
***Dimethylamine salt of dicamba (3,6 dichloro-6- anisic acid)	0.07%
INERT INGREDIENTS	98.97%
Total	100.00%

E.P.A. Reg. No. 557-1850

++Equivalent to 2,4-dichlorophenoxyacetic acid	0.54%
**Equivalent to 2-(2-methyl-4-chlorophenoxy) propionic acid	0.26%
***Equivalent to 3,6-dichloro-o-anisic acid.....	0.06%
†Isomer Specific by AOAC Method 6 D01.5	

E.P.A. Est. No. 557-CA-1


KEEP OUT OF REACH OF CHILDREN

CAUTION

In case of contact, wash skin with plenty of soap and water;
for eyes, flush with water for 15 minutes and get medical
attention. See back panel for additional precautionary
statements.

GOLDEN
V I G O R O

WEED CONTROL
PLUS
FERTILIZER



GOLDEN

VIGORO®

WEED CONTROL PLUS LAWN FERTILIZER

WEED & FEED

18-4-8

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)18.0%

0.6% Ammoniacal Nitrogen
12.4% Water Soluble Urea Nitrogen
5.0% Water Insoluble Nitrogen

AVAILABLE PHOSPHORIC ACID (P₂O₅) 4.0%
SOLUBLE POTASH (K₂O)..... 8.0%

Derived from ammoniated phosphates, urea, ureaformaldehyde, isobutylidene diurea and muriate of potash. Potential acidity 600 lbs. Calcium Carbonate Equivalent per ton. Manufactured by Estech, Inc., Chicago, Illinois 60602.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
& DOMESTIC ANIMALS

Avoid contact with skin, eyes or clothing.
Avoid breathing dust. Wash thoroughly
after handling. Harmful if swallowed.

ENVIRONMENTAL HAZARD

Do not apply directly to water. Do
not apply when weather conditions favor
drift from target area. Do not
contaminate any body of water by clean-
ing of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use
this product in a manner inconsistent
with its labeling.

WHEN TO APPLY -- For best results apply
when weeds are actively growing in the
spring or early fall after heavy rains.
Water lawn one day before application
to ensure adequate soil moisture. Do
not apply to wet grass. To get maximum
weed killing results, do not rake,
mow or water again until two days after
applying. Do not apply during excessive-
ly hot weather (85° F or above). Repeat

WEEDS CONTROLLED--The list below is
a partial list of the common weeds
controlled by this product.

Bedstraw	Knotweed	Ragweed
Black Medic	Lambsquarter	Sheep Sorrel
Buckhorn	Lespedeza	Shepherd's
Burdock	Mallow	Purse
Chicory	Morning Glory	Speedwell
Chickweed	Oxalis	Spurge
Clover	Peppergrass	Thistle
Dandelion	Pigweed	Wild Carrot
Dock	Plantains	Wild Garlic
Ground Ivy	Poison Ivy	Wild Lettuce
Heal-all	Poison Oak	Wild Onion
Henbit	Purslane	Yarrow

PRECAUTIONS: Not recommended for use
on bent grass, St. Augustine grass or
Dichondra lawns, as injury may result.
Do not use this product where desirable
clovers are present or allow the product
to "wind drift" on flowers, shrubs,
vegetables or ornamental shrubbery. Do
not apply to newly seeded lawns. During
application close spreader openings when
turning. Apply when air is calm. Do not
apply to areas underlaid by roots of
desirable trees and shrubs.

SPREADER SETTINGS
SPREADER WIDTH SETTING

treatment if necessary to control hard to kill weeds,

WHERE TO APPLY--This product may be used on bluegrass, fescue, rye, Bermuda, and zoysia lawns. If adverse low moisture conditions exist, weeds are in a poor state of growth, or the lawn is heavily overrun with weeds, more than one application may be required. Apply the second application in 30 days.

HOW TO APPLY--Using the spreader settings provided below, apply this product to 5,000 square feet of lawn area (100 x 50).

Scott's Professional R-7..9 ft.	G-H
Scott's Rotary C-96 ft.	C
Central "Roto-Spred"..... 7 ft.	3 1/2
Cyclone..... 6 ft.	3 5/8
Scott's Drop	3 1/2
Lawn Beauty	7 1/2
Sears	6 3/4
Wards	6 1/2
Kmart	6 1/4

NOTE: Spreader rate varies with walking speed. To insure proper coverage, check spreader over known area.

Storage: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal.

Product Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not re-use bag. Discard bag in trash.

