

ACCEPTED

NET WT.  
22 LBS.  
20-3-6  
TREATS  
2500 sq ft  
\$5.95



SUPER WEEDER-FEEDER for  
st. augustine, centipede,  
and zoysia grasses  
20-3-6


# SUPER WEEDER-FEEDER

for st. augustine  
centipede & zoysia  
grasses

SUPER WEEDER & FEEDER  
for st. augustine centipede and  
zoysia grasses

weeds  
as it  
feeds

CAUTION: This product  
is a weed killer and  
should be used  
carefully. Do not  
use on lawns of  
St. Augustine, Centipede,  
or Zoysia grasses.



SUPER WEEDER & FEEDER  
for st. augustine centipede and  
zoysia grasses

### GUARANTEED ANALYSIS

TOTAL NITROGEN	20.00%
NITRATE NITROGEN	0.00%
AMMONIACAL NITROGEN	2.30%
WATER SOLUBLE NITROGEN (Organic)	11.70%
WATER INSOLUBLE NITROGEN	6.00%
AVAILABLE PHOSPHORIC ACID	3.00%
WATERSOLUBLE POTASH	6.00%
TOTAL AVAILABLE PRIMARY PLANT FOOD	29.00%
CHLORINE, not more than	1.00%

PRIMARY PLANT FOOD SOURCES: Urea, Ammoniated Superphosphate, Ureatom, Sulphate of Potash and Isobutylidene Diurea, Florida Reg. No. 99-3003

Controls Such Weeds As

GERMINATING: Crabgrass, Barnyardgrass, Wiregrass, Paspalum Seedlings (Dallisgrass, Vaseygrass, Bahiagrass), Sedges (such as Cyperus Compressus), and Spurge

Apply between January 1 and March 31 (Florida to June 15th)

ESTABLISHED AND GERMINATING: Betony, Blackmedic (yellow trefoil), Burclover, Chickweed, Dandelion, Oxalis, Florida Pusley, Pennywort (Dollarweed) and Creeping Charlie (Matchweed)

### EASY TO USE

For best results, apply before weeds come up (pre-emergent). Established weeds are more difficult to control and may require an application in Spring and Fall. On newly sprigged St. Augustine, Centipede or Zoysia lawns, apply immediately after planting to prevent weeds from retarding growth.

Water in well

### CAUTION

Keep out of reach of children. Keep away from feed or foodstuffs. Do not take internally. Avoid inhalation or contact with eyes. Wash after handling or using.

Do not use north of North Carolina. Do not use on lawns of Bahiagrass, Bermudagrass, Dichondra, Bentgrass, or carpetgrass or where clover is present and desired.

Avoid application to the root zones of trees and shrubs. Avoid applications on garden plants, flowers, and shrubs. Delay spring planting for one month after application. Store in a dry place. Do not store near growing plants or fertilizers, seeds, insecticides or fungicides. Clean spreader after each use.

Apply only in Winter, Spring and Fall. This product is not recommended for use during months of July, August and September. Do not reuse empty bag. Destroy when empty.

### HOW TO SET YOUR SPREADER

CYCLONE	5	SCOTT'S	4 1/2
SEARS			7

NOTE: Spreader rate varies with walking speed. Check spreader over known area.