

Reg no: 557-1863

PM-25

193

FRONT PANEL

TREATS

5,000 sq. ft.


24-4-8

S U P E R
WEEDER-FEEDER

FOR ST. AUGUSTINE
CENTIPEDE & ZOYSIA GRASSES

WEEDS AS IT FEEDS

KEEP OUT OF REACH OF CHILDREN



C A U T I O N

In case of contact: for skin, wash with plenty
of soap and water; for eyes, flush with water for
15 minutes and get medical attention if irritation
persists.

See back panel for additional precautionary
statements.

ACTIVE INGREDIENTS:


Atrazine (2-chloro-4-ethylamino- 6-isopropylamino-s-triazine)	1.17%
INERT INGREDIENTS	98.83%
Total	100.00%

Contains 23.4 lbs. of Atrazine per ton

E.P.A. Est. No. 557-FL-1

E.P.A. Reg. No. 557-1863

Net Wt. 18 Lbs. (8.16 Kg)



S U P E R

WEEDER-FEEDER

FOR ST. AUGUSTINE, CENTIPEDE & ZOYSIA GRASSES

24-4-8

GUARANTEED ANALYSIS

TOTAL NITROGEN	24.00%
Ammoniacal Nitrogen.....	2.35%
Water Soluble Nitrogen (Organic).....	21.65%
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅).....	4.00%
SOLUBLE POTASH (K ₂ O)	8.00%
CHLORINE, NOT MORE THAN	9.00%

Derived from Urea, Triple Superphosphate, Muriate of Potash and Ammonium Sulphate.

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS & DOMESTIC ANIMALS

CAUTION

Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. Keep out of reach of children. Do not take internally.


ENVIRONMENTAL HAZARD

Keep out of lakes, streams or ponds.

Do not use north of North Carolina. Do not use on lawns of Bahiagrass, Bermudagrass, Dichondra, Bentgrass, or carpetgrass or where clover is present and desired.

Avoid application to the root zones of trees and shrubs. Avoid applications on garden plants, flowers and shrubs. Delay spring planting for one month after application. Clean spreader after each use.

Apply only in Winter, Spring and Fall. This product is not recommended for use during months of July, August and September.



GENERAL CLASSIFICATION

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Apply between January 1 and March 31 (Florida and other southern states to June 15th) at the rate of 18 lbs. on 5,000 sq. ft.

CONTROLS SUCH WEEDS AS:

GERMINATING: Crabgrass, Barnyardgrass, Wiregrass, Paspalum Seedlings (Dallisgrass, Vaseygrass, Bahiagrass), Sedges (such as Cyperus Compressus), and Spurge.

ESTABLISHED AND GERMINATING: Betony, Blackmedic (yellow trefoil), Burclover, Chickweed, Dandelion, Oxalis, Florida Pusley, Pennywort (Dollarweed), and Creeping Charlie (Matchweed).

For best results, apply before weeds come up (pre-emergent). Established weeds are more difficult to control and may require an application in Spring and Fall. On newly sprigged St. Augustine, Centipede, or Zoysia lawns, apply immediately after planting to prevent weeds from retarding growth. Water in well.

<p>Storage: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal.</p>	<p>Product Disposal: Securely wrap original container in several layers of newspaper and discard in trash.</p> <p>Container Disposal: Do not re-use bag. Discard bag in trash.</p>
--	--

SUGGESTED SPREADER SETTINGS

SPREADER	SETTING	SPREADER	SETTING
Wards 1091, 1093, 1095.....	6 1/2	Precision 20C Drop	4
Wards 1080, 1081.....	4 3/4	SB 40 Broadcast..	12
Jackson & Sears	4 3/4	24F Drop.....	1
Scott	5	Central 40 Drop	5 1/4
Kmart.....	5	60 Broadcast	4 1/8
		Cyclone	4

NOTE: Spreader rate varies with walking speed
To insure proper coverage, check spreader over known area.

NET WT. 18 LBS. (8.16 Kg)

Manufactured by

Estech, Inc., 30 North LaSalle Street
Chicago, Il. 60602