

# Super Weeder-Feeder

NOT REVIEWED  
In Accordance with EPA Pesticide 82-2,  
Based on Draft Labeling, Dated 3-13-85  
4-17-85

Treats  
5,000 sq. ft.  
24-4-8

# Super Weeder-Feeder

- FOR ST. AUGUSTINE CENTIPEDE & ZOYSIA GRASSES
- WEEDS AS IT FEEDS

KEEP OUT OF REACH OF CHILDREN

### CAUTION

In case of contact with skin, wash with plenty of water and soap. If eyes are affected, flush with water for 15 minutes and get medical attention if irritation persists.

See back panel for additional precautionary statements.

### ACTIVE INGREDIENTS

Atrazine (3-chloro-4-ethylamino-6-isopropylamino-2-pyridinyl) 1.17%

INERT INGREDIENTS 98.83%

Contains 23.4% of Atrazine (1.17%)

EPA Reg. No. 105-1166



Net Wt. 18 Lbs. (8.16 Kg)

## Super Weeder-Feeder

Super Weeder-Feeder

# Super Weeder-Feeder

## Super Weeder-Feeder FOR ST. AUGUSTINE, CENTIPEDE & ZOYSIA GRASSES 24-4-8

### GUARANTEED ANALYSIS

TOTAL NITROGEN		24.00%
Ammoniacal Nitrogen	2.35%	
Water Soluble Nitrogen (Organic)	21.65%	
AVAILABLE PHOSPHORIC ACID (P <sub>2</sub> O <sub>5</sub> )		4.00%
SOLUBLE POTASH (K <sub>2</sub> O)		8.00%
CHLORINE NOT MORE THAN		8.00%

Derived from Urea, Triple Superphosphate, Muriate of Potash and Ammonium Sulphate

### PRECAUTIONARY STATEMENTS

#### HAZARD TO HUMANS & DOMESTIC ANIMALS

#### CAUTION

Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. Keep out of reach of children. Do not take internally.

#### ENVIRONMENTAL HAZARD

Keep out of lakes, streams or ponds.

Do not use north of North Carolina. Do not use on lawns of Bahiagrass, Bermudagrass, Dichondra, Bentgrass or carpetgrass or where clover is present and desired.

Avoid application to the root zones of trees and shrubs. Avoid applications on garden plants, flowers and shrubs. Delay spring planting for one month after application. Clean spreader after each use.

Apply only in Winter, Spring and Fall. This product is not recommended for use during months of July, August and September.

### DIRECTIONS FOR USE

#### General Classification

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Apply between January 1 and March 31 (Florida and other southern states to June 15th) at the rate of 18 lbs. on 5,000 sq. ft.

#### Controls Such Weeds As:

GERMINATING: Crabgrass, Barnyardgrass, Wiregrass, Paspalum Seedlings (Dallisgrass, Vasegrass, Bahiagrass), Sedges (such as Cyperus Compressus) and Spurge.

ESTABLISHED AND GERMINATING: Betony, Blackmedic (yellow trefoil), Burclover, Chickweed, Dandelion, Oaks, Florida Pusley, Pennywort (Dollarweed) and Creeping Charlie (Matchweed).

For best results apply before weeds come up (pre-emergent). Established weeds are more difficult to control and may require an application in Spring and Fall. On newly sprigged St. Augustine, Centipede or Zoysia lawns, apply immediately after planting to prevent weeds from retarding growth. Water in well.

#### STORAGE:

Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal.

#### PRODUCT DISPOSAL:

Securely wrap original container in several layers of newspaper and discard in trash.

#### CONTAINER DISPOSAL:

Do not re-use bag. Discard bag in trash.

### SUGGESTED SPREADER SETTINGS

SPREADER	SETTING	SPREADER	SETTING
Wards 1051, 1093, 1095	6	Precision 20C Drop	4
Wards 1080, 1081	4	SB 40 Broadcast	12
Jackson & Sears	4	24" Drop	1
Scott	5	Central 40 Drop	5
Kmart	5	60 Broadcast	4
		Cyclone	4

NOTE: Spreader rate varies with walking speed. To insure proper coverage, check spreader over known area.

NET WT. 18 LBS. (8.16 Kg)

Manufactured by  
Estech, Inc., 30 North LaSalle Street  
Chicago, Il. 60602

## Super Weeder-Feeder