

FRONT PANEL

TREATS

5,000 sq. ft.

24-4-8

S U P E R

WEEDER-FEEDER

FOR ST. AUGUSTINE
CENTIPEDE & ZOYSIA GRASSES

WEEDS AS IT FEEDS

KEEP OUT OF REACH OF CHILDREN

C A U T I O N

In case of contact: for skin, wash with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention if irritation persists.


See back panel for additional precautionary statements.

ACTIVE INGREDIENTS:

*Atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine)	1.11%
*Related Compounds	0.06%
INERT INGREDIENTS	98.83%
Total	100.00%

*Contains 23.4 lbs. of Atrazine and related compounds per ton.

E.P.A. Est. No. 557-FL-1
E.P.A. Reg. No. 557-1863



Net Wt. 18 Lbs. (8.16 Kg)

REAR PANEL

S U P E R
WEEDER-FEEDER
FOR ST. AUGUSTINE, CENTIPEDE & ZOYSIA GRASSES

24-4-8

GUARANTEED ANALYSIS

TOTAL NITROGEN (N).....	24.00Z
Ammoniacal Nitrogen	2.35Z
Water Soluble Nitrogen (Organic)	21.65Z
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅).....	4.00Z
SOLUBLE POTASH (K ₂ O).....	8.00Z
Chlorine, not more than	8.00Z

Derived from Urea, Triple Superphosphate, Muriate of Potash and Ammonium Sulphate.

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS & DOMESTIC ANIMALS

C A U T I O N

Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. Keep out of reach of children. Do not take internally.

ENVIRONMENTAL HAZARD

Atrazine can leach through soil and can enter groundwater which may be used as drinking water. Atrazine has been found in groundwater. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters. Keep out of lakes, streams or ponds.

Do not use north of North Carolina. Do not use on lawns of Bahia-grass, Bermudagrass, Dichondra, Bentgrass, or carpetgrass where clover is present and desired.

Avoid application to the root zones of trees and shrubs. Avoid applications on garden plants, flowers and shrubs. Delay spring planting for one month after application. Clean spreader after each use.

Apply only in Winter, Spring and Fall. This product is not recommended for use during months of July, August and September.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Apply between January 1 and March 31 (Florida and other southern states to June 15th) at the rate of 18 lbs. on 5,000 sq. ft.

CONTROLS SUCH WEEDS AS:

GERMINATING: Crabgrass, Barnyardgrass, Wiregrass, Paspalum Seedlings (Dallisgrass, Vaseygrass, Bahiagrass), Sedges (such as Cyperus Compressus), and Spurge.

ESTABLISHED AND GERMINATING: Betony, Blackmedic (yellow trefoil), Burclover, Chickweed, Dandelion, Oxalis, Florida Pusley, Pennywort (Dollarweed), and Creeping Charlie (Matchweed).

For best results, apply before weeds come up (pre-emergent). Established weeds are more difficult to control and may require an application in Spring and Fall. On newly sprigged St. Augustine, Centipede, or Zoysia lawns, apply immediately after planting to prevent weeds from retarding growth. Water in well.

Approximately 4 weeks after initial treatment with product, an application of 16% or higher nitrogen fertilizer should be made at a rate of 6 lbs. per 1,000 sq. ft. and watered in well.

Storage: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal.	Product Disposal: Securely wrap original container in several layers of newspaper and discard in trash.
	Container Disposal: Do not re-use bag. Discard bag in trash.

SUGGESTED SPREADER SETTINGS

SPREADER	SETTING	SPREADER	SETTING
Wards 1091, 1093, 1095	6 1/4	Precision 20C Drop	4
Wards 1080, 1081	4 3/4	SB 40 Broadcast	12
Jackson & Sears	4 3/4	24F Drop	1
Scott	5	Central 40 Drop	5 1/4
Kmart	5	60 Broadcast	4 1/8
		Cyclone	4

NOTE: Spreader rate varies with walking speed. To insure proper coverage, check spreader over known area.

CAUTION: Close spreader openings when loading, turning or when stopped.

NET WT. 18 LBS. (8.16 Kg)

Manufactured by Vigoro Industries, Inc., Fairview Heights, IL 62203. VIGORO is a registered trademark of Vigoro Industries, Inc.