

557-1866

DIRECTIONS FOR USE

VIGORO ST. AUGUSTINE GRASS LAWN WEEDER AND FEEDER destroys growing broadleaf weeds as it feeds St. Augustine, Centipede and all other turf grasses.

CONTROLS SUCH WEEDS AS: Florida Pulsey, Black Medic, Chickweed, Clover, Cocklebur, Dandelion, Creeping Charlie, Dock, Henbit, Knotweed, Lambsquarter, Miard, Oxalis, Peppergrass, Pigweed, Plantains, Poison Ivy, Purslane, Ragweed, Sida, Spurge, Wild Onion, Wild Lettuce, and most other broadleaf weeds.

TIME TO APPLY: Apply any time broadleaf weeds are present, except avoid treating St. Augustine grass in summer months (June through September) when it is growing rapidly, because temporary foliage discoloration may occur. Best results are obtained when weeds are young and growing vigorously. If weeds are growing slowly, control will be slow and more than one application may be needed. Moisture from dew or watering on weed foliage at time of application aids control. Do not water for 24 to 48 hours after application.

HOW MUCH TO APPLY: Use 35 lbs. on 2,300 sq. ft. for St. Augustine Grass and all other grasses. Over-application may cause temporary discoloration.

SUGGESTED SPREADER SETTINGS

May vary with age, model and condition of spreader and walking speed. Check your spreader by trial on a known area.

Make of Spreader	SPREADER SETTINGS
Cyclone	6
K-Mart	9
Lawn Beauty	9
Scott	7½
Sears	7½
Ward	10

CAUTION: Keep out of reach of children. Keep away from feed and food products. Avoid inhalation or contact with eyes, skin or clothing. Wash after use. Do not re-use empty bag. Destroy when empty.

Avoid application on garden plants, flowers and shrubs. Do not use on newly seeded lawns. Do not contaminate irrigational ditches or water used for irrigational purposes. Avoid drift to desirable plants. Do not use on dichondra or lippla ground cover. Do not apply in areas underlaid by roots of desirable trees and shrubs. Store in a dry place, away from plants, seeds, fertilizers, insecticides or fungicides.

VIGORO

St. Augustine Grass
Lawn Weeder
& Feeder