

Use herbicides properly
Do not overdose
Read DIRECTIONS FOR USE
carefully

Net Wt 35 Lbs.
TREATS
2300 Sq. Ft.

ACC
July 21, 1976
TO ATT

VIGORO
St. Augustine Grass
Lawn Weeder & Feeder
Contains Trimec® Herbicide * Concentrate
For Use on St. Augustine Grass
And Other Grasses For
Control of Growing Broadleaf
Weeds

CAUTION: Keep Out of Reach of Children
See Back Panel for Additional Cautions

WEEDS
AS IT
FEEDS

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2,4-dichloro- phenoxyacetic acid	0.042%
**Dimethylamine Salt of 2- (2-methyl-4- chlorophenoxy) propionic acid	0.130%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	0.015%
INERT INGREDIENTS:	99.813%
TOTAL	100.000%

Used at 35 lbs. on 2,300 sq. ft. equivalent to:

- *2,4-D Acid - 0.035% - 0.23 lbs. per acre or 0.70 lbs. per ton.
- **MCPA Acid - 0.110% - 0.73 lbs. per acre or 2.29 lbs. per ton.
- ***Dicamba Acid - 0.013% - 0.08 lbs. per acre or 0.27 lbs. per ton.

Reg. T.M. PBI
U.S. Patent No. 3,284,186

VIGORO

E.P.A. Reg. No. 557-1866
St. Augustine Grass
Lawn Weeder &
Feeder

Net Wt. 35 Lbs.
Covers 2,300 Sq. Ft.
VIGORO

St. Augustine Grass
Lawn Weeder & Feeder
For Control of Growing Broadleaf Weed In
St. Augustine and All Other Grasses
12-5-7

12

GUARANTEED ANALYSIS

TOTAL NITROGEN,	12.00%
NITRATE NITROGEN,	None
AMMONIACAL NITROGEN,	2.00%
WATER SOLUBLE ORGANIC NITROGEN,	6.50%
WATER INSOLUBLE NITROGEN,	3.50%
AVAILABLE PHOSPHORIC ACID,	5.00%
WATER SOLUBLE POTASH,	7.00%
TOTAL AVAILABLE PRIMARY PLANT FOOD,	24.00%

St. Augustine Grass
Lawn Weeder & Feeder
Contains Trimec® Herbicide * Concentrate
For Use on St. Augustine Grass
And Other Grasses For
Control of Growing Broadleaf
Weeds

CAUTION: Keep Out of Reach of Children
See Back Panel for Additional Cautions

WEEDS
AS IT
FEEDS

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2,4-dichloro- phenoxyacetic acid	0.042%
**Dimethylamine Salt of 2- (2-methyl-4- chlorophenoxy) propionic acid	0.130%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	0.015%
INERT INGREDIENTS:	99.813%
TOTAL	100.000%

Used at 35 lbs. on 2,300 sq. ft. equivalent to:

- *2,4-D Acid - 0.035% - 0.23 lbs. per acre or 0.70 lbs. per ton.
- **MCPA Acid - 0.110% - 0.73 lbs. per acre or 2.29 lbs. per ton.
- ***Dicamba Acid - 0.013% - 0.08 lbs. per acre or 0.27 lbs. per ton.

Reg. T.M. PBI

*U.S. Patent No. 3,284,186

VIGORO

E.P.A. Reg. No. 557-1866

St. Augustine Grass
Lawn Weeder &
Feeder

Net Wt. 35 Lbs.
Covers 2,300 Sq. Ft.
VIGORO

St. Augustine Grass
Lawn Weeder & Feeder

For Control of Growing Broadleaf Weeds In
St. Augustine and All Other Grasses
12-5-7

GUARANTEED ANALYSIS

TOTAL NITROGEN,	12.00%
NITRATE NITROGEN,	None
AMMONIACAL NITROGEN,	2.00%
WATER SOLUBLE ORGANIC NITROGEN,	6.50%
WATER INSOLUBLE NITROGEN,	3.50%
AVAILABLE PHOSPHORIC ACID,	5.00%
WATER SOLUBLE POTASH,	7.00%
TOTAL AVAILABLE PRIMARY PLANT FOOD,	24.00%
CHLORINE, not more than	1.00%

Derived from Urea, Ammoniated Superphosphate,
Ureaform, Sulfate of Potash.
732-735-5058

Manufactured by Swift Agricultural Chemicals Corporation, Chicago, Illinois 60604;
(An Estech Company)

Use herbicides properly
Do not overdose
Read DIRECTIONS FOR USE
carefully.

Net Wt 35 Lbs.
TREATS
2300 Sq. Ft.

AC

VIGORO
St. Augustine Grass
Lawn Weeder & Feeder
Contains 2,4-D + Herbicide + Concentrate
Formulated on St. Augustine Grass
and Other Grasses For
Control of Growing Broadleaf
Weeds

CAUTION: Keep Out of Reach of Children
See Back Panel for Additional Cautions

WEEDS
AS IT
FEEDS

ACTIVE INGREDIENTS:

*Dimethyl 2,4-D	0.042%
**MCPA	0.130%
***Dicamba Acid	0.015%
INERT INGREDIENTS	99.813%
Total	100.000%

Used at Rate of:

- *2,4-D Acid 0.23 lb. per acre or 0.70 lbs. per ton.
- **MCPA 0.73 lb. per acre or 2.29 lbs. per ton.
- ***Dicamba Acid 0.27 lb. per acre or 0.27 lbs. per ton.

Reg. E.M.P.P.

*U.S. Patent No. 2,841,441

E.A. Reg. No. 557-1866

St. Augustine Grass

and Other Grasses

and

WEEDS
AS IT
FEEDS

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2,4-dichloro-phenoxyacetic acid	0.042%
**Dimethylamine Salt of 2- (2-methyl-4-chlorophenoxy) propionic acid	0.130%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	0.015%
INERT INGREDIENTS:	99.813%
TOTAL	100.000%

Used at 35 lbs. on 2,300 sq. ft. equivalent to:

*2,4-D Acid - 0.035% - 0.23 lbs. per acre or 0.70 lbs. per ton.	
**MCPA Acid - 0.110% - 0.73 lbs. per acre or 2.29 lbs. per ton.	
***Dicamba Acid - 0.013% - 0.08 lbs. per acre or 0.27 lbs. per ton.	
Reg. T.M. PBI	
*U.S. Patent No. 3,284,186	E.P.A. Reg. No. 557-1866
VIGORO	St. Augustine Grass

Lawn Weeder &
Feeder

Net Wt. 35 Lbs.
Covers 2,300 Sq. Ft.
VIGORO

St. Augustine Grass
Lawn Weeder & Feeder

For Control of Growing Broadleaf Weeds In
St. Augustine and All Other Grasses
12-5-7

GUARANTEED ANALYSIS

TOTAL NITROGEN	12.00%
NITRATE NITROGEN	None
AMMONIACAL NITROGEN	2.00%
WATER SOLUBLE ORGANIC NITROGEN	6.50%
WATER INSOLUBLE NITROGEN	3.50%
AVAILABLE PHOSPHORIC ACID	5.00%
WATER SOLUBLE POTASH	7.00%
TOTAL AVAILABLE PRIMARY PLANT FOOD	24.00%
CHLORINE not more than	1.00%
Derived from Urea, Ammoniated Superphosphate, Greatest Sulfate of Potash	
REGISTERED	

Manufactured by V.I. Agricultural Chemicals Corporation, Chicago, Illinois 60604
U.S. Pat. 3,284,186

TIME TO APPLY: Apply any time broadleaf weeds are present, except avoid treating St. Augustine grass in summer months (June through September) when it is growing rapidly, because temporary foliage discoloration may occur. Best results are obtained when weeds are young and growing vigorously. If weeds are growing slowly, control will be slow and more than one application may be needed. Moisture from dew or watering on weed foliage at time of application aids control. Do not water for 24 to 48 hours after application.

HOW MUCH TO APPLY: Use 35 lbs. on 2,300 sq. ft. for St. Augustine Grass and all other grasses. Over-application may cause temporary discoloration.

SUGGESTED SPREADER SETTINGS

May vary with age, model and condition of spreader and walking speed. Check your spreader by trial on a known area.

Make of Spreader	SPREADER SETTINGS
Cyclone	6
K-Mart	9
Lawn Beauty	9
Scott	7½
Sears	7½
Ward	10

CAUTION: Keep out of reach of children. Keep away from feed and food products. Avoid inhalation or contact with eyes, skin or clothing. Wash after use. Do not re-use empty bag. Destroy when empty.

Avoid application on garden plants, flowers and shrubs. Do not use on newly seeded lawns. Do not contaminate irrigational ditches or water used for irrigational purposes. Avoid drift to desirable plants. Do not use on dichondra or lippia ground cover. Do not apply in areas underlaid by roots of desirable trees and shrubs. Store in a dry place, away from plants, seeds, fertilizers, insecticides or fungicides.

VIGORO

St. Augustine Grass
Lawn Weeder
& Feeder