

Reg no: 557-1866

PM-25

194

Do not overdose. Read
DIRECTIONS FOR USE
carefully

NET WT. 40 LBS. (18.14 Kg.)
12-5-7
TREATS 5,000 SQ. FT.

V I G O R O

ST. AUGUSTINE GRASS
LAWN WEEDER & FEEDER

WEEDS AS IT FEEDS

For Use on St. Augustine Grass,
Bermuda Grass, Bluegrass, Fescue
And other Grasses for
Control of Growing Broadleaf Weeds

Contains Trimec® Herbicide* Concentrate

ACCEPTED

AUG - 1 1985
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 557-1866

KEEP OUT OF REACH OF CHILDREN

C A U T I O N

In case of contact, wash skin with
plenty of soap and water; for eyes,
flush with water for 15 minutes
and get medical attention. See
back panel for additional pre-
cautionary statements.

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2,4- dichlorophenoxyacetic acid	0.073%
**Dimethylamine Salt of 2-(2-methyl-4- chlorophenoxy) propionic acid	0.255%
***Dimethylamine Salt of Dicamba (3,6- dichloro-o-anisic acid	0.031%
INERT INGREDIENTS	99.641%
TOTAL	100.000%

Used at 40 lbs. on 5,000 sq. ft. equivalent to:

- *2,4-D Acid-0.061%-0.21 lbs. per acre or
1.22 lbs. per ton
- **MCPP Acid-0.212%-0.73 lbs. per acre or
4.24 lbs. per ton
- ***Dicamba Acid-0.025%-0.08 lbs. per acre or
0.50 lbs. per ton
- *Isomer Specific by AOAC Method 6 D01-5
-Reg T M PBI

U.S. Patent No. 3,284,186

E.P.A. Reg. No: 557-1866

E.P.A. Est. No: 557-11-1

234

REAR PANEL

NET WT. 40 LBS. (18.14 Kg.)

TREATS 5,000 SQ. FT.

V I G O R O

ST. AUGUSTINE GRASS
LAWN WEEDER & FEEDER

For Control of Growing Broadleaf Weeds in
St. Augustine, Bermudagrass, Bluegrass, Fescue
and All Other Grasses

12-5-7

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	12.0%
12.0% Ammoniacal Nitrogen	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)	5.0%
SOLUBLE POTASH (K ₂ O)	7.0%

Derived from ammoniated phosphates, ammonium sulphate and muriate of potash.

Statement of Secondary Nutrients

Sulphur, combined (S)	11.0%
-----------------------------	-------

Derived from ammonium sulphate

Chlorine, not more than	5.8%
-------------------------------	------

Potential Acidity 900 lbs. Calcium Carbonate Equivalent per ton. Manufactured by Estech, Inc., Chicago, Illinois 60602. Vigoro® is a registered trademark of Estech, Inc.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

C A U T I O N

Avoid contact with skin, eyes, or clothing. Avoid breathing dust.
Keep out of reach of children. Wash thoroughly after handling.
Harmful if swallowed.

BEST AVAILABLE COPY

ENVIRONMENTAL HAZARD

Do not apply directly to lakes, streams or ponds. Do not apply when weather conditions favor drift from target area. Avoid application on garden plants, flowers and shrubs. Do not use on newly seeded lawns. Do not contaminate irrigational ditches or water used for irrigational purposes. Avoid drift to desirable plants. Do not use on Dichondra, Lippia or other ground cover. Do not apply in areas underlaid by roots of desirable trees and shrubs.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

VIGORO ST. AUGUSTINE GRASS LAWN WEEDER AND FEEDER destroys growing broadleaf weeds as it feeds St. Augustine, Bermuda, Bluegrass, Fescue, Centipede and all other turf grasses.

CONTROLS SUCH WEEDS AS: Florida Pusley, Black Medic, Chickweed, Clover, Cocklebur, Dandelion, Creeping Charlie, Dock, Henbit, Knotweed, Lambsquarter, Mustard, Oxalis, Peppergrass, Pigweed, Plantains, Poison Ivy, Purslane, Ragweed, Sida, Spurge, Wild Onion, Wild Lettuce, and most other broadleaf weeds.

TIME TO APPLY: Apply any time broadleaf weeds are present, except avoid treating St. Augustine grass in summer months (June through September) when it is growing rapidly, because temporary foliage discoloration may occur. Best results are obtained when weeds are young and growing vigorously. If weeds are growing slowly, control will be slow and more than one application may be needed. Moisture from dew or watering on weed foliage at time of application aids control. Do not water 24 to 48 hours after application.

HOW MUCH TO APPLY: Use 40 lbs. on 5,000 sq. ft. for St. Augustine Grass and all other grasses. Over-application may cause temporary discoloration.

SUGGESTED SPREADER SETTINGS

May vary with age, model and condition of spreader and walking speed. Check your spreader by trial on a known area.

SPREADER	SETTING	SPREADER	SETTING
Vigoro Rotary....	4 3/4	Wards	4 3/4
Vigoro Drop	6 1/2	Scotts PF-1 ...	7 1/4
Cyclone	4 3/4	K-mart	4 1/2
Sears Mark II ...	8		



Storage: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal.

Product Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not re-use bag. Discard bag in trash.

