

PM-25

Reg no: 557-1866

1 2

EPA 82-2

In Accordance with IR Notice 82-2.

Based on Draft Labeling Dated 8-1-85

10-1-85

VIGORO
ST. AUGUSTINE GRASS
LAWN WEEDER AND
FEEDER

NET WT. 40 LBS. (18.14 kg.)

12-5-7
TREATS 5,000 SQ. FT.

Do not overdose. Read
DIRECTIONS FOR USE
carefully.

VIGORO

ST. AUGUSTINE GRASS LAWN WEEDER & FEEDER

WEEDS AS IT FEEDS

For Use on St. Augustine Grass,
Bermuda Grass, Bluegrass, Fescue,
And other Grasses for
Control of Growing Broadleaf Weeds

Contains Trimec® Herbicide® Concentrate

KEEP OUT
OF REACH OF CHILDREN

CAUTION

In case of contact, wash skin with plenty of
soap and water. For eyes, flush with water
for 15 minutes and get medical attention.
See this label for additional precautionary
statements.

ACTIVE INGREDIENTS

Trimec® Herbicide® Concentrate	100%
Other Ingredients	0%
TOTAL	100%

Net Wt. 40 LBS. (18.14 kg.)
Gross Wt. 40 LBS. (18.14 kg.)
Net Wt. 40 LBS. (18.14 kg.)
Gross Wt. 40 LBS. (18.14 kg.)
Net Wt. 40 LBS. (18.14 kg.)
Gross Wt. 40 LBS. (18.14 kg.)

VIGORO

ST. AUGUSTINE GRASS
LAWN WEEDER AND
FEEDER

VICPRO

TREATS 5,000 SQ. FT.

VIGORO

ST. AUGUSTINE GRASS LAWN WEEDER & FEEDER

**For Control of Growing Broadleaf Weeds in
St. Augustine, Bermudagrass, Bluegrass,
Fescue, and All Other Grasses**

12-5-7

GUARANTEED ANALYSIS

[illegible]

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

As mentioned in your report, clothing is being worn by the
subject in the form of a shirt, trousers, shoes, socks, and underwear.

ENVIRONMENTAL HAZARD

[illegible]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner not consistent with its labeling.

VIGORO ST AUGUSTINE GRASS LAWN WEEDER AND FEEDER destroys growing broadleaf weeds as it feeds St Augustine, Bermuda, Bluegrass, Fescue, Centipede and all other turf grasses.

CONTROLS SUCH WEEDS AS: Kruse Pursley Black Margy (Pheasant) Chick Cornish
Dandelion Creeping Charlie Dock Smartweed Lamb's-ear Mustard Irony Pigweed Grass
Pigweed Plants Portulaca Aca and Ragweed Soda Spurge Red Clover Stink Lemon and
most other broadleaf weeds

TIME TO APPLY: Apply any time broadleaf weeds are present, except during fruiting. Start applying grass in summer months (June to August/September) when it is growing rapidly, but after temperatures have begun to rise. In some areas, application may occur later. Best results are obtained when weeds are young and growing vigorously. If weeds are growing slowly or late in the season, increase the dosage and/or apply more frequently. Additional applications may be needed. Most lawn grasses will tolerate 6-8 treatments per year. However, if there is significant damage to the turf, reduce the frequency of applications.

HOW MUCH TO APPLY USA & IN THE ISLANDS

SUGGESTED SPREADER SETTINGS

Only vary with age (model and results in 1 separate article by myself, and a co-author, 1994)

SPREADER	SETTING	SPREADER	SETTING
1. 1/2" 10/16" 1/2"	4%	1. 1/2" 10/16" 1/2"	4%
2. 1/2" 10/16" 1/2"	4%	2. 1/2" 10/16" 1/2"	4%
3. 1/2" 10/16" 1/2"	4%	3. 1/2" 10/16" 1/2"	4%
4. 1/2" 10/16" 1/2"	4%	4. 1/2" 10/16" 1/2"	4%

STORAGE: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal.

PRODUCT DISPOSAL: Securely wrap original container in brown paper or newspaper and discard in trash.

CONTAINER DISPOSAL: Do not re-use bag. Discard bag in trash.

VIGORO