

Reg No: 557-1866

PM-25

18 U

FRONT PANEL

Do not overdose. Read DIRECTIONS FOR USE carefully.

TREATS 5000 SQ. FT. 12-5-7


VIGORO®

ST. AUGUSTINE GRASS LAWN WEEDER & FEEDER

WEEDS AS IT FEEDS

For use on St. Augustine Grass, Bermuda Grass, Bluegrass, Fescue And other Grasses for Control of Growing Broadleaf Weeds

Contains Trimec® Herbicide* Concentrate



KEEP OUT OF REACH OF CHILDREN CAUTION

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention. See back panel for additional precautionary statements.

ACTIVE INGREDIENTS:

++Dimethylamine Salt of 2,4-dichlorophenoxyacetic acid	0.073%
**Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	0.255%
***Dimethylamine Salt of dicamba (3,6-dichloro-o-anisic acid)	0.031%
INERT INGREDIENTS:	99.641%
TOTAL	100.000%

Used at 40 lbs. on 5,000 sq. ft. equivalent to:

- +2,4-D Acid - 0.061% - 0.21 lbs. per acre or 1.22 lbs. per ton.
- **MCPP Acid - 0.212% - 0.73 lbs. per acre or 4.24 lbs. per ton.
- ***Dicamba Acid - 0.025% - 0.08 lbs. per acre or 0.50 lbs. per ton.
- +Isomer Specific by AOAC Method 6.D01.5
- Reg T M PBI

U.S. Patent No. 3,284,186
EPA Reg. No. 557-1866

EPA Est. No. 557-IL-1

NET WT. 40 LBS.



REAR PANEL

NET WT. 40 LBS. (18.14 Kg.)

TREATS 5,000 SQ. FT.

V I G O R O®

ST. AUGUSTINE GRASS

LAWN WEEDER & FEEDER

For Control of Growing Broadleaf Weeds
in St. Augustine, Bermudagrass, Bluegrass, Fescue
and All Other Grasses

12-5-7

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	12.0%
12.0% Ammoniacal Nitrogen	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)	5.0%
SOLUBLE POTASH (K ₂ O)	7.0%

Derived from ammoniated phosphates, ammonium sulphate and muriate of potash.

Statement of Secondary Nutrients

Sulphur, combined (S)	11.0%
---------------------------------	-------

Derived from ammonium sulphate.

Chlorine, not more than	5.8%
-----------------------------------	------

Potential Acidity 900 lbs. Calcium Carbonate Equivalent per ton.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. Keep out of reach of children. Do not re-enter or permit others to re-enter treated areas until dusts have settled.

Protective Clothing: Applicators must wear long pants, long-sleeved shirts, shoes and gloves.

BEST AVAILABLE COPY

REAR PANEL Cont.

ENVIRONMENTAL HAZARD

This product is toxic to aquatic invertebrates and non-target plants. Drift or run-off may adversely affect non-target plants. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate any body of water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area. Do not contaminate irrigational ditches or water used for irrigational purposes.

PRECAUTIONS: Avoid application on garden plants, flowers and shrubs. Do not use on newly seeded lawns. Avoid drift to desirable plants. Do not use on Dichondra, Lippia or other ground cover. Do not apply in areas underlaid by roots of desirable trees and shrubs.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

VIGORO® ST. AUGUSTINE GRASS LAWN WEEDER AND FEEDER destroys growing broadleaf weeds as it feeds St. Augustine, Bermuda, Bluegrass, Fescue, Centipede and all other turf grasses.

CONTROLS SUCH WEEDS AS: Florida Pusley, Black Medic, Chickweed, Clover, Cocklebur, Dandelion, Creeping Charlie, Dock, Henbit, Knotweed, Lambsquarter, Mustard, Oxalis, Peppergrass, Pigweed, Plantains, Poison Ivy, Purslane, Ragweed, Sida, Spurge, Wild Onion, Wild Lettuce, and most other broadleaf weeds.

TIME TO APPLY: Apply any time broadleaf weeds are present, except avoid treating St. Augustine grass in summer months (June through September) when it is growing rapidly, because temporary foliage discoloration may occur. Best results are obtained when weeds are young and growing vigorously. If weeds are growing slowly, control will be slow and more than one application may be needed. Moisture from dew or watering on weed foliage at time of application aids control. Do not water 24 to 48 hours after application.

HOW MUCH TO APPLY: Use 40 lbs. on 5,000 sq. ft. for St. Augustine Grass and all other grasses. Over-application may cause temporary discoloration. Approximately 4 weeks after initial treatment, an application of 16% or higher nitrogen fertilizer should be made at a rate of 6 lbs. per 1,000 sq. ft. and watered in well.

BEST AVAILABLE COPY

REAR PANEL Cont.

 STORAGE: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal.

PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

SUGGESTED SPREADER SETTINGS

May vary with age, model and condition of spreader and walking speed. Check your spreader by trial on a known area.

READER	SETTING	SI READER	SETTING
Vigoro Rotary	4 3/4	Wards	4 3/4
Vigoro Drop	6 1/2	Scotts PF-1	7 1/4
Cyclone	4 3/4	Kmart	4 1/2
Sears Mark II	8		

CAUTION: Close spreader openings when loading, turning or when stopped.

Manufactured by Vigoro Industries, Inc., Fairview Heights, IL 62208
 Vigoro ® is a registered trademark of Vigoro Industries, Inc.