




CERTIFICATION OF COMPLIANCE

I, Judith K. Lucash, being an authorized representative of Vigoro Industries, Inc., certify that all containers of

Vigoro St. Augustine Grass Lawn Weeder And Feeder

EPA Reg. No. 557-1866

produced by June 15, 1994 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing use products. I further certify that all containers of said product sold or distributed by this company by January 1, 1995 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products.

  
\_\_\_\_\_  
Judith K. Lucash  
Manager of Regulatory Affairs

December 21, 1992

2075

FRONT PANEL

Do not overdose. Read DIRECTIONS FOR USE carefully.

TREATS 5000 SQ. FT. 12-5-7

V I G O R O ®

ST. AUGUSTINE GRASS LAWN WEEDER & FEEDER

WEEDS AS IT FEEDS

For use on St. Augustine Grass, Bermuda Grass, Bluegrass, Fescue And other Grasses for Control of Growing Broadleaf Weeds

Contains Trimec® Herbicide\* Concentrate

KEEP OUT OF REACH OF CHILDREN CAUTION

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention. See back panel for additional precautionary statements.

ACTIVE INGREDIENTS:

+*Dimethylamine Salt of 2,4-dichlorophenoxyacetic acid . . . . .	0.073%
**Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid . . . . .	0.255%
***Dimethylamine Salt of dicamba (3,6-dichloro-o-anisic acid) . . . . .	0.031%
INERT INGREDIENTS: . . . . .	99.641%
TOTAL . . . . .	100.000%

Used at 40 lbs. on 5,000 sq. ft. equivalent to:

- +\*2,4-D Acid - 0.061% - 0.21 lbs. per acre or 1.22 lbs. per ton.
- \*\*MCPA Acid - 0.212% - 0.73 lbs. per acre or 4.24 lbs. per ton.
- \*\*\*Dicamba Acid - 0.025% - 0.08 lbs. per acre or 0.50 lbs. per ton.
- +Isomer Specific by AOAC Method 6.D01.5
- Reg T M PBI

U.S. Patent No. 3,284,186  
EPA Reg. No. 557-1866

EPA Est. No. 557-IL-1

NET WT. 40 LBS.

3075

NET WT. 40 LBS. (18.14 Kg.)

TREATS 5,000 SQ. FT.

V I G O R O®

ST. AUGUSTINE GRASS

LAWN WEEDER & FEEDER

For Control of Growing Broadleaf Weeds  
in St. Augustine, Bermudagrass, Bluegrass, Fescue  
and All Other Grasses

12-5-7

GUARANTEED ANALYSIS

12.0% Ammoniacal Nitrogen	12.0%
AVAILABLE PHOSPHORIC ACID (P <sub>2</sub> O <sub>5</sub> )	5.0%
SOLUBLE POTASH (K <sub>2</sub> O)	7.0%

Derived from ammoniated phosphates, ammonium sulphate and muriate of potash.

Statement of Secondary Nutrients

Sulphur, combined (S)	11.0%
-----------------------	-------

Derived from ammonium sulphate.

Chlorine, not more than	5.8%
-------------------------	------

Potential Acidity 900 lbs. Calcium Carbonate Equivalent per ton.

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION

Avoid contact with skin, eyes and clothing. Avoid breathing dust. Wash thoroughly after handling. Harmful if swallowed. Keep out of reach of children.

Protective Clothing: When using this product, wear long-sleeved shirt, long pants, socks, shoes and gloves. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

ENVIRONMENTAL HAZARD

This product is toxic to aquatic invertebrates. Drift or run-off may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate any body of water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area. Do not contaminate irrigational ditches or water used for irrigational purposes.

PRECAUTIONS: Avoid application on garden plants, flowers and shrubs. Do not use on newly seeded lawns. Avoid drift to desirable plants. Do not use on Dichondra, Lippia or other ground cover. Do not apply in areas underlaid by roots of desirable trees and shrubs.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment areas during application. Do not enter treatment areas until spray has dried or dust has settled.

VIGORO® ST. AUGUSTINE GRASS LAWN WEEDER AND FEEDER destroys growing broadleaf weeds as it feeds St. Augustine, Bermuda, Bluegrass, Fescue, Centipede and all other turf grasses.

) CONTROLS SUCH WEEDS AS: Florida Pusley, Black Medic, Chickweed, Clover, Cocklebur, Dandelion, Creeping Charlie, Dock, Henbit, Knotweed, Lambsquarter, Mustard, Oxalis, Peppergrass, Pigweed, Plantains, Poison Ivy, Purslane, Ragweed, Sida, Spurge, Wild Onion, Wild Lettuce, and most other broadleaf weeds.

TIME TO APPLY: Apply any time broadleaf weeds are present, except avoid treating St. Augustine grass in summer months (June through September) when it is growing rapidly, because temporary foliage discoloration may occur. Best results are obtained when weeds are young and growing vigorously. If weeds are growing slowly, control will be slow and more than one application may be needed. Moisture from dew or watering on weed foliage at time of application aids control. Do not water 24 to 48 hours after application.

) HOW MUCH TO APPLY: Use 40 lbs. on 5,000 sq. ft. for St. Augustine Grass and all other grasses. Over-application may cause temporary discoloration.

The maximum application rate for this product is 8.0 lbs. per 1,000 sq. ft. The maximum number of broadcast applications per treatment site is 2 per year.

Approximately 4 weeks after initial treatment, an application of 16% or higher nitrogen fertilizer should be made at a rate of 6 lbs. per 1,000 sq. ft. and watered in well.

REAR PANEL Cont.

-----  
 STORAGE: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal.

PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.  
 CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.  
 -----

SUGGESTED SPREADER SETTINGS

May vary with age, model and condition of spreader and walking speed. Check your spreader by trial on a known area.

SPREADER	SETTING	SPREADER	SETTING
Vigoro Rotary . . . . .	4 3/4	Wards . . . . .	4 3/4
Vigoro Drop . . . . .	6 1/2	Scotts PF-1 . . . . .	7 1/4
Cyclone . . . . .	4 3/4	Kmart . . . . .	4 1/2
Sears Mark II . . . . .	8		

CAUTION: Close spreader openings when loading, turning or when stopped.

Manufactured by Vigoro Industries, Inc., Fairview Heights, IL 62208  
Vigoro ® is a registered trademark of Vigoro Industries, Inc.