

Net Wt 35 Lbs.  
Covers 2,300 Sq. Ft.

VIGORO  
BAHIA and BERMUDA  
Lawn Weeder & Feeder

For Control of Growing Broadleaf Weeds  
in Bahia, Bermuda, Fescue, Zoysia  
Lawn Grasses

12-5-7

GUARANTEED ANALYSIS

TOTAL NITROGEN.....	12.00
NITRATE NITROGEN.....	None
AMMONIACAL NITROGEN.....	2.00%
WATER SOLUBLE ORGANIC NITROGEN.....	6.50%
WATER INSOLUBLE NITROGEN.....	3.50%
AVAILABLE PHOSPHORIC ACID.....	5.00%
WATER SOLUBLE POTASH.....	7.00%
TOTAL AVAILABLE PRIMARY PLANT FOOD.....	24.00%
CHLORINE, not more than.....	1.00%

Derived from Urea, Ammoniated Superphosphate,  
Ureaform, Sulfate of Potash  
732-735-5053

Manufactured by Swift Agricultural Chemicals Corporation,  
Chicago, Illinois 60604: (An Estech Company)

DIRECTIONS FOR USE

VIGORO BAHIA AND BERMUDA GRASS LAWN WEEDER AND FEEDER destroys  
growing broadleaf weeds as it feeds Bahia, Bermuda, Fescue, Zoysia turf grasses.

CONTROLS SUCH WEEDS AS: Florida Pusley, Black Medic, Chickweed, Clover,  
Cocklebur, Dandelion, Creeping Charlie, Dock, Henbit, Knotweed, Lambsquarter,  
Mustard, Oxalis, Peppergrass, Pigweed, Plantains, Poison Ivy, Purslane, Ragweed,  
Sida Spurge, Wild Onion, Wild Lettuce, and most other broadleaf weeds

HOW TO APPLY:

Apply when weeds are young and actively growing and the average temperature is  
over 70°F. Apply while lawn is wet with dew or after sprinkling lightly. Do not mow  
or water your lawn for two days after applying.

Use spreader settings below. Covers 2,300 sq. ft. (46' x 50') per bag

SUGGESTED SPREADER SETTINGS

May vary with age, model and condition of spreader and walking speed. Check your  
spreader by trial on a known area.

Make of Spreader	SPREADER SETTINGS
Cyclone	6
K Mart	9
Lawn Beauty	

Use herbicides properly  
Do not overdose  
Read DIRECTIONS FOR USE  
carefully

Net Wt 35 Lbs.  
TREATS  
2300 Sq. Ft.

VIGORO  
BAHIA AND BERMUDA  
Lawn Weeder & Feeder

Contains Trimec Herbicide \* Concentrate  
For Use on Bahia, Bermuda, Fescue, and  
Zoysia, for Control of Growing Broadleaf Weeds

CAUTION: Keep Out of Reach of Children  
See Back Panel for Additional Cautions

WEEDS  
AS IT  
FEEDS

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2,4-dichloro- phenoxyacetic acid.....	0.084%
**Dimethylamine Salt of 2- (2-methyl-4- chlorophenoxy) propionic acid.....	0.260%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid).....	0.032%
INERT INGREDIENTS.....	99.624%
TOTAL.....	100.000%

Used at 35 lbs. on 2,300 sq. ft. equivalent to:

- \*2,4-D Acid - 0.070% - 0.46 lbs. per acre or 1.41 lbs. per ton.
- \*\*MCPP Acid - 0.220% - 1.46 lbs. per acre or 4.59 lbs. per ton.
- \*\*\*Dicamba Acid - 0.027% - 0.17 lbs. per acre or 0.55 lbs. per ton.

**VIGORO**  
**BAHIA AND BERMUDA**  
 Lawn Weeder & Feeder  
 Contains Trimec - Herbicide \* Concentrate  
 For Use on Bahia, Bermuda, Fescue, and  
 Zoysia, for Control of Growing Broadleaf Weeds  
**CAUTION:** Keep Out of Reach of Children  
 See Back Panel for Additional Cautions

WEEDS  
 AS IT  
 FEEDS

**ACTIVE INGREDIENTS:**

*Dimethylamine Salt of 2,4-dichloro- phenoxyacetic acid .....	0.084%
**Dimethylamine Salt of 2- (2-methyl-4- chlorophenoxy) propionic acid .....	0.260%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid) .....	0.032%
INERT INGREDIENTS: .....	99.624%
<b>TOTAL .....</b>	<b>100.000%</b>

Used at 35 lbs. on 2,300 sq. ft. equivalent to:

- \*2,4-D Acid - 0.070% - 0.46 lbs. per acre or 1.41 lbs. per ton.
- \*\*MCPP Acid - 0.220% - 1.46 lbs. per acre or 4.59 lbs. per ton.
- \*\*\*Dicamba Acid - 0.027% - 0.17 lbs. per acre or 0.55 lbs. per ton.

Reg. TM PBI - KCK  
 \*U.S. Patent No. 3,284,186  
**VIGORO**

E.P.A. Reg. No. 557-1867  
 Bahia and Bermuda  
 Lawn Weeder & Feeder

**TOTAL AVAILABLE NITROGEN PLANT FOOD, .....** 1.00%  
 CHLORINE, not more than .....

Derived from Urea, Ammoniated Superphosphate,  
 Ureaform, Sulfate of Potash.  
 732-735-5053

Manufactured by Swift Agricultural Chemicals Corporation,  
 Chicago, Illinois 60604: (An Estech Company)

**DIRECTIONS FOR USE**

**VIGORO BAHIA AND BERMUDA GRASS LAWN WEEDER AND FEEDER** destroys growing broadleaf weeds as it feeds Bahia, Bermuda, Fescue, Zoysia turf grasses.

**CONTROLS SUCH WEEDS AS:** Florida Pusley, Black Medic, Chickweed, Clover, Cocklebur, Dandelion, Creeping Charlie, Dock, Henbit, Knotweed, Lambsquarter, Mustard, Oxalis, Peppergrass, Pigweed, Plantains, Poison Ivy, Purslane, Ragweed, Sida Spurge, Wild Onion, Wild Lettuce, and most other broadleaf weeds

**HOW TO APPLY:**

Apply when weeds are young and actively growing and the average temperature is over 70°F. Apply while lawn is wet with dew or after sprinkling lightly. Do not mow or water your lawn for two days after applying.

Use spreader settings below. Covers 2,300 sq. ft. (46' x 50') per bag.

**SUGGESTED SPREADER SETTINGS**

May vary with age, model and condition of spreader and walking speed. Check your spreader by trial on a known area.

Make of Spreader	SPREADER SETTINGS
Cyclone	6
K-Mart	9
Lawn Beauty	9
Scott	7 1/2
Chars	7 1/2
Ward	10

**CAUTION:** Keep out of reach of children. Keep away from feed and food products. Avoid inhalation or contact with eyes, skin or clothing. Wash after use. Do not re-use empty bag. Destroy when empty.

Avoid application on garden plants, flowers and shrubs. Do not use on newly seeded lawns. Do not contaminate irrigational ditches or water used for irrigational purposes. Avoid drift to desirable plants. Do not use on Centipede, St. Augustine, dichondra, lippia or other ground cover. Do not apply in areas underlaid by roots of desirable trees and shrubs.

Store in a dry place, away from plants, seeds, fertilizers, insecticides or fungicides.

**VIGORO**

Bahia and Bermuda  
 Lawn Weeder  
 and Feeder