

lbs.  
TS  
0  
VNS  
-6  
TION  
back

NEW!  
VIGORO  
DICHONDRA  
TREATMENT

net wt 25 lbs

VIGORO®

SIX IN ONE  
DICHONDRA  
TREATMENT

SIX  
IN  
ONE

11-20-73  
557-1868  
comments

DICHONDRA TREATMENT

12-6-6

SWF



VIGOR

# DICHONDRA TREATMENT

## 12-6-6

### GUARANTEED ANALYSIS

TOTAL NITROGEN (N) ..... 12.0%  
 1.0% Ammoniacal Nitrogen (derived from ammoniated phosphates and ammonium sulfate)  
 4.0% Water Insoluble Nitrogen (derived from Condensed Urea Formaldehyde)  
 7.0% Water Soluble Nitrogen (derived from urea and urea formaldehyde)

AVAILABLE PHOSPHORIC ACID (P<sub>2</sub>O<sub>5</sub>)  
 (derived from ammoniated phosphates) ..... 6.0%

SOLUBLE POTASH (K<sub>2</sub>O) (derived from muriate of potash) ..... 6.0%

IRON EXPRESSED AS ELEMENTAL Fe (derived from ferrous sulfate) ..... 1.0%

Potential acidity 400 lbs Calcium Carbonate equivalent per ton.  
 Manufactured by Swift Chemical Company, Chicago, Illinois 60604

### MONEY BACK PROMISE

You must be pleased with the results or you are entitled to your money back. Simply send your sales slip to VIGORIO, Chicago, Illinois 60604 for prompt refund.

### EASY TO USE

Apply at least twice a year—but not more than 3 times a year at 3 months interval. Do not use in heavily shaded areas, or near flowers and shrubs, or over the root zone of trees. Do not reseed for 60 days after application. Always shut off spreader when making sharp turns and do not over-lap spreader widths.

Dichondra foliage must be dry—but do hose off lightly after application. Resume normal watering practices after 12 hours, but do not flood or puddle.

Apply once during the period November thru March—and again 3 months later to prevent weeds:

- |                  |                |          |
|------------------|----------------|----------|
| crabgrass        | chickweed      | pigweed  |
| annual bluegrass | knotweed       | purslane |
| ryegrass         | spotted spurge |          |
| sorrel           | yellow oxalis  |          |

may need follow-up spot treatment for long entrenched invaders


To subdue bermuda, apply before bermuda goes dormant in the fall and again just before it greens up in the spring

To kill lawnmoth larvae, cutworms, armyworms, sowbugs, ants, earwigs, and fleas, use any time they are a problem

For crabgrass control, apply April 1 thru June 20.

Established Lawns: Dichondra is a heavy feeder and should be fertilized with Vigoro Dichondra Fertilizer every 6 weeks to 2 months to maintain good growth and color. Well fed, healthy, properly maintained dichondra lawns are less apt to become weed infested than unhealthy lawns.

CAUTION





not over-lap spreader widths.

**Dichondra foliage must be dry** – but do hose off lightly after application. Resume normal watering practices after 12 hours. but do not flood or puddle.

Apply once during the period November thru March - and again 3 months later to prevent weeds:

- |                  |                |          |
|------------------|----------------|----------|
| crabgrass        | chickweed      | pigweed  |
| annual bluegrass | knotweed       | purslane |
| ryegrass         | spotted spurge |          |
| sorrel           | yellow oxalis  |          |

may need follow-up spot treatment for long entrenched invaders

**To subdue bermuda**, apply before bermuda goes dormant in the fall and again just before it greens up in the spring.

**To kill lawnmoth larvae, cutworms, armyworms, sowbugs, ants, earwigs, and fleas**, use any time they are a problem.

**For crabgrass control**, apply April 1 thru June 20.

**Established Lawns:** Dichondra is a heavy feeder and should be fertilized with Vigoro Dichondra Fertilizer every 6 weeks to 2 months to maintain good growth and color. Well fed, healthy, properly maintained dichondra lawns are less apt to become weed infested than unhealthy lawns.

### CAUTION

*(This section contains very faint, mostly illegible text, likely a standard warning or disclaimer for the product.)*

3-VD6/1-25

### HOW TO GET YOUR SHEARER

CYCLONE	5%	SCOTTS	7%
K-MART	9%	SEARS	8%
LAWN BEAUTY	9%	VARDS	8%

NOTE: See your local dealer for information on where to purchase this product.

CAUTION: This product is a restricted use pesticide. See label for restrictions.



SIX IN ONE

WIGORO

NET WT 25 LBS  
TREATS  
2000  
SQUARE FEET  
12-6-6  
SATISFACTION

557-1868

ACCEPTED

NOV 20 1973

U.S. DEPARTMENT OF AGRICULTURE  
FURNISHED UNDER THE PESTICIDE ACT  
FOR INFORMATION ONLY. NOT REGISTERED  
UNDER FEDERAL LAW. SUBJECT  
ATTACHED COMMENTS.

CONTAINS 1%  
**IRON**

# WIGORO

WEED & INSECT CONTROL PLUS FERTILIZER

# SIX IN

SIX IN ONE  
DICHONDRA  
TREATMENT

O

IN  
ONE

# DICHONDRA TREATMENT

FOR ESTABLISHED DICHONDRA LAWNS ONLY

1. Preparation of spray solution...
2. Application of spray solution...
3. Precautions...
4. Storage...

CAUTION: KEEP OUT OF REACH  
OF CHILDREN.

OK  
OK  
OK  
OK

# FOR ESTABLISHED DICHONDRA LAWNNS ONLY

- 1) Preparation of lawn: ...
- 2) ...
- 3) ...
- 4) ...
- 5) ...
- 6) ...

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**

SEE BACK PANEL FOR ADDITIONAL CAUTIONS.

**ACTIVE INGREDIENTS:**


Diphenamid (N,N-Dimethyl-2,2-d-phenylacetamide) .....	1.83%
Monuron (3-(p-chlorophenyl)-1,1-dimethylurea) .....	0.15%
Chlordane, technical .....	0.56%

**INERT INGREDIENTS:**

.....	97.43%
Total .....	100.00%

\*Equivalent to 0.34% of trichloro-4,7-Methane tetrahydroindane and 0.20% related compounds.

EPA Reg. No. 507-1508-AA



## DIRECTIONS

ANTS, MILLIPEDES, CENTIPEDES, CRICKETS, SOW BUGS, GRASSHOPPERS — apply with powder duster to the entire ground area about 4 to 6 feet out and around outside house foundation, in severe cases extend the powder treatment 8 to 10 feet out from the building. Be sure to apply to window wells, under porches and steps, along lower part of foundation, in cracks in cement walks and patios, around garbage containers, and to other areas where insects can hide and breed. Repeat as necessary.

ANTS IN LAWNS — apply with powder duster heavily around nests and ant hills and lightly over entire lawn area at the rate of 1/2 lb. to about 350 sq. ft. of area. Keep children and pets off treated areas until the area has been well watered and had dried completely to get PEST-X DUST into nests and soil. Repeat as necessary.

**CAUTION:** This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Apply this product only as specified on this label.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.



# PEST-X DUST

### ACTIVE INGREDIENTS:

Technical Chlordane 10%  
(equivalent to 6% octachloro-4, 7 metha-  
notetrahydroindane and 4% related com-  
pounds)

INERT INGREDIENTS 90%

**KILLS AND CONTROLS ANTS,  
AND CERTAIN OTHER INSECTS**

**CAUTION: KEEP OUT OF REACH OF  
OF CHILDREN.**

See back panel for additional cautions.

NET CONTENTS:

PRICE:

## DIRECTIONS

THRIPS, STRAWBERRY ROACHES,  
CHINCH BUGS, STINK BUGS,  
WORMS, EARWIGS, CUT-  
WHITE GRUBS—

apply with powder duster to  
upper and lower leaf surfaces  
and ornamentals and general  
ground around plants, working  
4 inches of soil around base  
Repeat as necessary.

**CAUTION:** Avoid inhalation,  
contact, wash with soap and  
hands after use. Avoid contact  
feed and foodstuffs. Do not  
drinking water or use near fish

*Terminix International*

1125 N. Third St., Milwaukee,

EPA Reg. No. 561-19

## DIRECTIONS

CENTIPEDES, CRICKETS, SOW BUGS, apply with powder duster to the entire ground out and around outside house foundation, in a powder treatment 8 to 10 feet out from to apply to window wells, under porches and of foundation, in cracks in cement walks and containers, and to other areas where insects meet as necessary.

Apply with powder duster heavily around nests over entire lawn area at the rate of 1/2 lb. area. Keep children and pets off treated areas well watered and had dried completely to get and soil. Repeat as necessary.

Product is toxic to fish and wildlife. Birds areas may be killed. Keep out of any use of this product only as specified on this

CONTAINER. DESTROY WHEN EMPTY.



# PEST-X DUST

### ACTIVE INGREDIENTS:

Technical Chlordane 10%  
(equivalent to 6% octachloro-4, 7 metha-  
notetrahydroindane and 4% related com-  
pounds)

INERT INGREDIENTS 90%

**KILLS AND CONTROLS ANTS,  
AND CERTAIN OTHER INSECTS**

**CAUTION: KEEP OUT OF REACH OF  
OF CHILDREN.**

See back panel for additional cautions.

NET CONTENTS:

PRICE:

## DIRECTIONS

THRIPS, STRAWBERRY ROOT WEEVIL,  
CHINCH BUGS, STINK BUGS, ARMY-  
WORMS, EARWIGS, CUTWORMS,  
WHITE GRUBS—

Apply with powder duster directly on upper and lower leaf surfaces of flowers and ornamentals and generously on the ground around plants, working into top 4 inches of soil around base of plants. Repeat as necessary.

**CAUTION:** Avoid inhalation. Avoid skin contact, wash with soap and water. Wash hands after use. Avoid contamination of feed and foodstuffs. Do not contaminate drinking water or use near fish ponds.

***Terminix International, Inc.***

1125 N. Third St., Milwaukee, Wis. 53203

EPA Reg. No. 561-19



CODE 0755

SOLD TO VETERINARIANS ONLY

# MANGEOL

## Insecticide

(Emulsifying)

A non-staining, greaseless acaricide and insecticide for dogs.

**ACTIVE INGREDIENTS:** Benzyl benzoate 30.00%; Triethanolamine 2.51%; Oleic acid 4.86%; Chlordane, technical\* 0.25%.

**INERT INGREDIENTS:** 62.38%.

**TOTAL INGREDIENTS:** 100.00%.

\*Equivalent to 0.15% Octachloro-4, 7-methanotetrahydroindane and 0.1% related compounds.

**CAUTION:** Keep Out of Reach of Children.

See side panel for additional precautions.

**1 U. S. GALLON = 0.83 IMP. GAL.**

**DIRECTIONS:** Apply to skin with swab or brush following soap bath and rinse well. Either allow animal to dry or towel it dry. Limit application to less than 1/3 of body surface at one treatment. In sarcoptic mange, repeat after a week, if necessary; in demodectic mange, twice weekly applications are recommended. Demodectic mange is often difficult to diagnose accurately and treatment may involve factors such as diet. If no improvement is noted after 10 days, reconsult a veterinarian. Mangeol is more effective if parasitized areas are clipped before treatment is applied.

**WARNING:** Flammable: Keep Away From Heat or Open Flame! Harmful if swallowed. Do not contaminate feed or drinking water. Avoid contact with eyes or mouths of animals. Avoid ingestion, inhalation or skin contact, and wash hands with soap and water after using. Do not use on cats. On young puppies, use as a spot treatment only. Rinse empty container thoroughly with water and discard it. Treatment: Call a physician. Remove material from skin or alimentary tract. Avoid oil laxatives. Nervous symptoms may be treated with barbiturates.  
EPA Reg. No. 569-16-AA

Lot No.

**Haver-Lockhart**  
LABORATORIES

