

VIGORO®

REGISTERED TRADEMARK
© 1970
VIGORO CORPORATION
1000 W. 10th St.
Oklahoma City, Oklahoma 73101
EPA REG. NO. 557-1870

10% SEVIN* DUST
FOR USE ON
VEGETABLES
LAWNS SHRUBS

10% SEVIN* DUST FOR USE ON VEGETABLES LAWNS SHRUBS

ACTIVE INGREDIENT:

| | |
|---|--------------|
| Carbaryl (l-Naphthyl N-methylcarbamate) | 10.0% |
| INERT INGREDIENTS: | <u>90.0%</u> |
| | 100.0% |

E.P.A. Reg. No. 557-1870

*Registered Trademark of Union Carbide Corporation for Carbaryl Insecticide

NET WT 4 LBS (64 OZ)

CAUTION Keep Out Of Reach Of Children
See Back Panel For Additional Cautions

VIGORO 10% SEVIN DUST

DIRECTIONS FOR USE

Vigoro 10% Sevin Dust is to be used without dilution. Apply uniformly to foliage with a conventional duster. Cover all exposed parts (both upper and lower leaf surfaces) with a thin, even film of dust using any garden type duster. Apply preferably in early morning or late evening when the air is quiet. Apply when insects or their damage appear. Best results are secured by dusting regularly at 7-10 day intervals throughout the growing season. Foliage should be dry before dust is applied.

| Crops & Insects Controlled | Last Application Allowed Before Harvest |
|--|---|
| <u>Cotton</u> - Thrips, Flea Beetles, Leafworm, Striped Blister Beetle, Boll Weevil, Bollworm, Fall Armyworms, Leaf Perforator, Leaf Rollers, Leafhoppers, Tarnished Plant Bug, Lygus & Other Plant Bugs, Grasshoppers, Pink Bollworm, Salt-marsh Caterpillar, Stink Bugs. Do not pasture dairy animals or animals being finished for slaughter in cotton fields or feed gin waste. | 0 Days |
| <u>Corn Field and Sweet Corn</u> - Corn Earworm, Corn Rootworm Adults, Fall Armyworms, Sap Beetles, Japanese Beetle, Flea Beetles, Leafhoppers. Apply when first silks appear and repeat at 2 to 3 day intervals during silking period. Direct dusters to ear zone to insure complete coverage of each silk. On sweet corn for food it can be applied up to and including the day of harvest. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population. | 7 Days For Forage |
| <u>Sugar Beets</u> - Leafhoppers, Armyworms, Webworms, Grasshoppers. | 14 Days |
| <u>Peanuts, Alfalfa, Clovers, Cowpeas, Soybeans</u> - Blister Beetles, Mexican Bean Beetle, Thrips, Leafhoppers, Three-cornered Alfalfa Hopper, Bean Leaf Beetle, Japanese Beetle, Alfalfa Caterpillar, Velvetbean Caterpillar, Green Cloverworm, Armyworms, Webworms, Corn Earworm, Stink Bugs, Grasshoppers. | 10 Days |
| <u>Tobacco (Field)</u> - Hornworms, Budworms, Flea Beetles, June Beetles, Japanese Beetles. <u>Tobacco (Plant Beds)</u> - Tobacco Flea Beetle (Apply 1 lb. per 100 sq. yds. 3 days before transplanting). | 3 Days Before Priming Or Cutting |
| <u>Rice, Forage Grass, Pasture</u> - Thrips, Stink Bugs, Armyworms, Grasshoppers. | 0 Days 14 Days On Rice |
| <u>Sorghum</u> - Armyworms, Webworms, Corn Earworm, Stink Bug, Grasshoppers. | 21 Days On Grain |
| <u>Sorghum Midge</u> - Apply 3 or 4 days after heads have emerged from boot. | 0 Days On Forage |
| <u>Asparagus (On Spears)</u> - Asparagus Beetle. Do not repeat applications within 3 days. <u>Asparagus (On Ferns and Brush)</u> - Asparagus Beetle, Apache Cicado. Post-Harvest Application. | 1 Day |
| <u>Beans</u> - Lygus Bugs, Stink Bugs, Corn Earworm, Leafhoppers, Fleabeetles, Tarnished Plant Bug, Bean Leaf Roller, Mexican Bean Beetle, Bean Leaf Beetle, Western Bean Cutworm, Japanese Beetle, Velvetbean Caterpillar, Armyworms. Do not feed forage to dairy animals or animals being finished for slaughter. | 0 Days |
| <u>Cucumbers, Squash, Cantaloupes, Watermelons</u> - Striped Cucumber Beetle, Spotted Cucumber Beetle, Melonworm, Pickleworm, Squash Bug, Leafhoppers, Fleabeetles. | 0 Days |
| <u>Potatoes, Tomatoes, Eggplants, Peppers</u> - Fleabeetles, Armyworms (Fall), Tomato Fruit Worm, Corn Earworm, Tomato Hornworms, Leafhoppers, European Corn Borer, Stink Bugs, Colorado Potato Beetle, Tomato Fruitworm. | 0 Days |

| | |
|---|---------------------------------|
| transplanting. | Cutting |
| <u>Rice, Forage Grass, Pasture</u> – Thrips, Stink Bugs, Armyworms, Grasshoppers. | 0 Days 14 Days On Rice |
| <u>Sorghum</u> – Armyworms, Webworms, Corn Earworm, Stink Bug, Grasshoppers. | 21 Days On Grain |
| <u>Sorghum Midge</u> – Apply 3 or 4 days after heads have emerged from boot. | 0 Days On Forage |
| <u>Asparagus (On Spears)</u> – Asparagus Beetle. Do not repeat applications within 3 days. | 1 Day |
| <u>Asparagus (On Ferns and Brush)</u> – Asparagus Beetle, Apache Cicado. Post-Harvest Application. | |
| <u>Beans</u> – Lygus Bugs, Stink Bugs, Corn Earworm, Leafhoppers, Fleabeetles, Tarnished Plant Bug, Bean Leaf Roller, Mexican Bean Bee'le, Bean Leaf Beetle, Western Bean Cutworm, Japanese Beetle, Velvetbean Caterpillar, Armyworms. Do not feed forage to dairy animals or animals being finished for slaughter. | 0 Days |
| <u>Cucumbers, Squash, Cantaloupes, Watermelons</u> – Striped Cucumber Beetle, Spotted Cucumber Beetle, Melonworm, Pickleworm, Squash Bug, Leafhoppers, Fleabeetles. | 0 Days |
| <u>Potatoes, Tomatoes, Eggplants, Peppers</u> – Fleabeetles, Armyworms (Fall), Tomato Fruit Worm, Corn Earworm, Tomato Hornworms, Leafhoppers, European Corn Borer, Stink Bugs, Colorado Potato Beetle, Tomato Fruitworm, Lace Bug, Lygus Bugs. | 0 Days |
| <u>Lettuce</u> – Fleabeetles, Stink Bugs, Corn Earworm, Leafhoppers, Armyworms, Harlequin Bug, Spittlebugs, Lygus Bugs. | 3 Days (Head) 14 Days (Leaf) |
| <u>Carrots</u> – Flea Beetles, Leafhoppers, Six-spotted Leafhopper, (Aster Yellows Vector), Armyworms, Spittlebugs, Lygus & Stink Bugs. | 0 Days |
| <u>Cabbage, Broccoli, Brussels Sprouts, Cauliflower, Kohlrabi</u> – Fleabeetles, Harlequin Bug, Corn Earworm, Armyworms. | 3 Days |
| <u>Okra</u> – Corn Earworm, Stink Bugs. | 0 Days |
| <u>Cranberries</u> – Leafhoppers, Fruitworms, Fireworms, Cutworms, Japanese Beetle. | 1 Day |
| <u>Strawberries</u> – Spittlebugs, Strawberry Leaf Roller, Strawberry Weevil. | 1 Day |
| <u>Blueberry</u> – Blueberry Maggot, Cherry Fruitworm, Cranberry Fruitworm, European Fruit Lecanium, Japanese Beetle. | 0 Day |
| <u>Grapes</u> – Grape Leafhoppers, Grape Leaf Folder, Grape Berry Moth, European Fruit Lecanium, Grape Leaf Skeletonizer, Japanese Beetle, June Beetles, Red-Banded Leaf Roller. | 0 Day |
| <u>Ornamentals</u> – Arbor Vitae, Azalea, Aster, Birch, Boxwood, Carnation, Camellia, Chrysanthemum, Dahlia, Dogwood, Elm, Gladiolus, Holly, Juniper, Lilac, Marigold, Petunia, Rhododendron, Rose, Rose of Sharon, Yew, Zinnia. Controls Aphids, Bagworm, Blister Beetle, Boxelder Bug, Leaf Beetles, Leafhoppers, Leaf Rollers, Orange Tortrix, Periodical Cicada, Plant Bug, Tent Caterpillar, Thrips (Exposed). Apply to completely cover foliage and repeat as necessary. | 0 Day |

CAUTION: Harmful if swallowed. Do not take internally. Avoid breathing of dust. Skin contact may be harmful. Avoid contact. Wash hands before eating. Wash thoroughly after using. Keep out of reach of children and animals.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Further protective information may be obtained from your cooperative State Agricultural Extension Service.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do not reuse container. Destroy when empty.

WILSON CHEMICAL COMPANY
 Estech Company
 10000 Wilshire Blvd, Los Angeles, CA 90024

