

VIGORO®

SEVIN® brand

Carbaryl Insecticide

10% DUST

**FOR USE
ON
VEGETABLES
LAWNS
SHRUBS**

ACCEPTED

MAR 26 1991

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended for the pesticide registered under EPA Reg. No. 557-1870

**KEEP OUT OF
REACH OF CHILDREN**

CAUTION

See additional
precautionary statements
on back panel

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl
N-methylcarbamate) . . . 10%

INERT INGREDIENT . . . 90%

TOTAL 100%

E.P.A. Est. No. 769-GA-1

E.P.A. Reg. No. 557-1870

SEVIN® is a registered trademark of Union Carbide Corporation for Carbaryl Insecticide

NET WEIGHT 4 LBS.

BEST AVAILABLE COPY

VIGORO 10% SEVIN DUST

DIRECTIONS FOR USE. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Vigoro 10% Sevin Dust is to be used without dilution. Apply uniformly to foliage with a conventional duster. Cover all exposed parts (both upper and lower leaf surfaces) with a thin, even film of dust using any garden type duster. Apply preferably in early morning or late evening

when the air is quiet. Apply when insects or their damage appear. Best results are secured by dusting regularly at 7-10 day intervals throughout the growing season. Foliage should be dry before dust is applied. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

CROPS	& INSECTS CONTROLLED	LAST APPLICATION ALLOWED BEFORE HARVEST
Sweet Corn	Corn Earworm, Corn Rootworm Adults, Fall Armyworms, Sap Beetles, Japanese Beetle, Flea Beetles, Leafhoppers. On sweet corn for food it can be applied up to and including the day of harvest. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.	0 Days
Asparagus (On Spears)	Asparagus Beetle. Do not repeat applications within 3 days.	1 Day
Asparagus (On Ferns and Brush)	Asparagus Beetle, Apache Cicada, Post-Harvest Application.	
Beans	Lycus Bugs, Stink Bugs, Corn Earworm, Leafhoppers, Fleabeetles, Tarnished Plant Bug, Bean Leaf Roller, Mexican Bean Beetle, Bean Leaf Beetle, Western Bean Cutworm, Japanese Beetle, Velvetbean Caterpillar, Armyworms. Do not feed forage to dairy animals or animals being finished for slaughter.	0 Days
Cucumbers, Squash, Cantaloupes, Watermelons	Striped Cucumber Beetle, Spotted Cucumber Beetle, Melonworm, Pickleworm, Squash Bug, Leafhoppers, Fleabeetles.	0 Days
Potatoes, Tomatoes, Eggplants, Peppers	Fleabeetles, Armyworms (Fall), Tomato Fruit Worm, Corn Earworm, Tomato Hornworms, Leafhoppers, European Corn Borer, Stink Bugs, Colorado Potato Beetle, Tomato Fruitworm, Lace Bug, Lycus Bugs	0 Days
Lettuce	Fleabeetles, Stink Bugs, Corn Earworm, Leafhoppers, Armyworms, Harlequin Bug, Spittlebugs, Lycus Bugs.	3 Days (Head) 14 Days (Leaf)
Carrots	Fleabeetles, Leafhoppers, Six-spotted Leafhopper, (Aster Yellows Vector), Armyworms, Spittlebugs, Lycus & Stink Bugs.	0 Days
Cabbage, Broccoli, Brussels Sprouts, Cauliflower, Kohlrabi	Fleabeetles, Harlequin Bug, Corn Earworm, Armyworms	3 Days
Okra	Corn Earworm, Stink Bugs	0 Days
Strawberries	Spittlebugs, Strawberry Leaf Roller, Strawberry Weevil	1 Day
Grapes	Grape Leafhoppers, Grape Leaf Folder, Grape Berry Moth, European Fruit Lecanium, Grape Leaf Skeletonizer, Japanese Beetle, June Beetles, Red-Banded Leaf Roller	0 Day
LAWNS & ORNAMENTAL TURF	Leafhoppers, Fall Armyworms, Sod Webworms (Lawn Moths) Ants, Fleas, Earwigs	Apply at the rate of 2 lbs. per 1000 sq. ft. For best results apply after rain or watering and do not water for at least 2 days after application. Repeat in 2 to 3 weeks if necessary.

SEVIN[®] brand Carbaryl Insecticide 10% DUST

ORNAMENTALS: Arbor Vitae, Azalea, Aster, Birch, Boxwood, Carnation, Camellia, Chrysanthemum, Dahlia, Dogwood, Elm, Gladiolus, Holly, Juniper, Lilac, Marigold, Petunia, Rhododendron, Rose, Rose of Sharon, Yew, Zinnia.

Controls Aphids, Bagworm, Blister Beetle, Boxelder Bug, Leaf Beetles, Leafhoppers, Leaf Rollers, Orange Tortrix, Periodical Cicada, Plant Bug, Tent Caterpillar, Thrips, (Exposed). Apply to completely cover foliage and repeat as necessary. 0 Day

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS:

CAUTION: May be harmful if swallowed. Avoid breathing vapors (dust or spray mist). Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Keep out of reach of children and animals.

ENVIRONMENTAL HAZARDS:

PRECAUTIONS: This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands, except for rice uses. Discharge from rice fields may kill aquatic and estuarine invertebrates. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment areas.

Further protective information may be obtained from your cooperative State Agricultural Extension Service.

STATEMENT OF PRACTICAL TREATMENT:

IF SWALLOWED, induce vomiting and seek medical attention immediately. **IF IN EYES OR ON SKIN,** flush eyes thoroughly with water, wash skin thoroughly with soap and water.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do not reuse container. Destroy when empty.

STORAGE AND DISPOSAL

STORAGE: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal.

PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

DISCLAIMER OF WARRANTIES

NOTE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, seller makes no claims other than those stated on this label.

BEST AVAILABLE COPY

Distributed by
Vigoro Industries, Inc.
Fairview Heights, IL 62208
Vigoro[®] is a registered trademark of Vigoro Industries, Inc.

VIGORO®

10% SEVIN DUST

10% SEVIN DUST

In Accordance with EPA Pesticide
Based on the Pesticide Act of 1972
Mar. 26. 91

ACTIVE INGREDIENT:

Carbaryl (1-Naphthyl N-methylcarbamate)..... 10.0%

INERT INGREDIENTS..... 90.0%

100.0%

®Registered Trademark of Union Carbide Corporation for Carbaryl Insecticide

E.P.A. Reg. No. 557-1870
E.P.A. Est. No. 34874-TX-1

**FOR USE ON
VEGETABLES
LAWNS • SHRUBS**

NET WEIGHT 4 LBS. (64 OZ.)

Keep Out Of Reach Of Children.

CAUTION

See Back Panel For Additional Cautions.

VIGORO®

VIGORO 10% SEVIN DUST

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Vigoro 10% Sevin Dust is to be used with dilution. Apply uniformly to foliage with a conventional duster. Cover all exposed parts (both upper and lower leaf surfaces) with a thin, even film of dust using any garden type duster. Apply preferably in

early morning or late evening when the air is quiet. Apply when insects or their damage appear. Best results are secured by dusting regularly at 7-10 day intervals throughout the growing season. Foliage should be dry before dust is applied. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s)

ORNAMENTALS Arbor Vitae, Azalea, Aster, Birch, Boxwood, Carnation, Camellia, Chrysanthemum, Dahlia, Dogwood, Elm, Gardenia, Holly, Juniper, Lilac, Marigold, Petunia, Rhododendron, Rose, Rose of Sharon, Yew, Zinnia

Controls Aphids, Bagworm, Blister Beetle, Boxelder Bug, Leaf Beetles, Leafhoppers, Leaf Rollers, Orange Tortrix, Periodical Cicada, Plant Bug, Tent Caterpillar, Thrips. (Exposed) Apply to completely cover foliage, and repeat as necessary

LAST APPLICATION ALLOWED BEFORE HARVEST

CROPS	& INSECTS CONTROLLED	LAST APPLICATION ALLOWED BEFORE HARVEST
Cotton	Thrips, Flea Beetles, Leafworm, Striped Blister Beetle, Boll Weevil, Bollworm, Fall Armyworms, Leaf Perforator, Leaf Rollers, Leafhoppers, Tarnished Plant Bug, Lygus & Other Plant Bugs, Grasshoppers, Pink Bollworm, Salt-marsh Caterpillar, Stink Bugs Do not pasture dairy animals or animals being finished for slaughter in cotton fields or feed gin waste	7 days
Field and Sweet Corn	Corn Earworm, Corn Rootworm Adults, Fall Armyworms, Sap Beetles, Japanese Beetle, Flea Beetles, Leafhoppers. Apply when first silks appear and repeat at 2 to 3 day intervals during silking period. Direct dusters to ear zone to insure complete coverage of each silk. On sweet corn for food it can be applied up to and including the day of harvest. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population	7 Days For Forage
Sugar Beets	Leafhoppers, Armyworms, Webworms, Grasshoppers	14 Days
Peanuts, Alfalfa, Clovers, Cowpeas, Soybeans	Blister Beetles, Mexican Bean Beetle, Thrips, Leafhoppers, Three-cornered Alfalfa Hopper, Bean Leaf Beetle, Japanese Beetle, Alfalfa Caterpillar, Velvetbean Caterpillar, Green Cloverworm, Armyworms, Webworms, Corn Earworm, Stink Bugs, Grasshoppers	10 Days
Tobacco (Field)	Hornworms, Budworms, Flea Beetles, June Beetles, Japanese Beetles	3 Days Before Priming Or Cutting
Tobacco (Plant Beds)	Tobacco Flea Beetle (Apply 1 lb per 100 sq yds 3 days before transplanting)	0 Days
Rice, Forage Grass, Pasture	Thrips, Stink Bugs, Armyworms, Grasshoppers. Do not use on rice fields in which crayfish and/or catfish farming is included in the cultural practice	14 Days On Rice
Sorghum	Armyworms, Webworms, Corn Earworm, Stink Bug, Grasshoppers	21 Days On Grain
Sorghum Midge	Apply 3 or 4 days after heads have emerged from boot	0 Days On Forage
Asparagus (On Spears)	Asparagus Beetle. Do not repeat applications within 3 days	1 Day
Asparagus (On Ferns and Brush)	Asparagus Beetle, Apache Cicada. Post-Harvest Application	
Beans	Lygus Bugs, Stink Bugs, Corn Earworm, Leafhoppers, Fleabeetles, Tarnished Plant Bug, Bean Leaf Roller, Mexican Bean Beetle, Bean Leaf Beetle, Western Bean Cutworm, Japanese Beetle, Velvetbean Caterpillar, Armyworms. Do not feed forage to dairy animals or animals being finished for slaughter	0 Days
Cucumbers, Squash, Cantaloupes, Watermelons	Striped Cucumber Beetle, Spotted Cucumber Beetle, Melonworm, Pickleworm, Squash Bug, Leafhoppers, Fleabeetles	0 Days
Potatoes, Tomatoes, Eggplants, Peppers	Fleabeetles, Armyworms (Fall), Tomato Fruit Worm, Corn Earworm, Tomato Hornworms, Leafhoppers, European Corn Borer, Stink Bugs, Colorado Potato Beetle, Tomato Fruitworm, Lace Bug, Lygus Bugs	0 Days
Lettuce	Fleabeetles, Stink Bugs, Corn Earworm, Leafhoppers, Armyworms, Harlequin Bug, Spittlebugs, Lygus Bugs	3 Days (Head) 14 Days (Leaf)
Carrots	Fleabeetles, Leafhoppers, Six-spotted Leafhopper, (Aster Yellow's Vector), Armyworms, Spittlebugs, Lygus & Stink Bugs	0 Days
Cabbage, Broccoli, Brussels Sprouts, Cauliflower, Kohlrabi	Fleabeetles, Harlequin Bug, Corn Earworm, Armyworms	3 Days
Okra	Corn Earworm, Stink Bugs	0 Days
Strawberries	Spittlebugs, Strawberry Leaf Roller, Strawberry Weevil	1 Day
Grapes	Grape Leafhoppers, Grape Leaf Folder, Grape Berry Moth, European Fruit Lecanium, Grape Leaf Skeletonizer, Japanese Beetle, June Beetles, Red-Banded Leaf Roller	0 Day

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS:

CAUTION May be harmful if swallowed. Avoid breathing vapour (spray mist). Avoid contact with eyes, skin or clothing. Wash after handling. Keep out of reach of children and animals.

ENVIRONMENTAL HAZARDS:

PRECAUTIONS This pesticide is extremely toxic to aquatic invertebrates. Do not apply directly to water or wetlands, except uses. Discharge from rice fields may kill aquatic invertebrates. Do not contaminate water by cleaning of or disposal of wastes. This product is highly toxic to bees exposed to treatment or residues on blooming crops or weeds. Do not product or allow it to drift to blooming crops or weeds if bees are the treatment areas.

Further protective information may be obtained from your State Agricultural Extension Service.

STATEMENT OF PRACTICAL TREATMENT:

IF SWALLOWED, induce vomiting and seek medical attention. **IF IN EYES OR ON SKIN**, flush eyes thoroughly with water, thoroughly with soap and water. **NOTE FOR PHYSICIAN** Carbaryl is a moderate, reversible, cholinergic inhibitor. Atropine is antidotal. Do not reuse container. Be empty.

STORAGE AND DISPOSAL

STORAGE Store in original container in safe dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal.

PRODUCT DISPOSAL wrap original container in several layers of newspaper, discard in trash. **CONTAINER DISPOSAL** use bag. Discard bag.

DISCLAIMER OF WARRANTIES

NOTE Seller warrants that this product conforms to the ingredient statement. Since conditions of use such as weather, compatibility with other conditions of application equipment, will vary, seller makes no claims other than stated on this label.

Distributed by
Estech, Inc., 30 North LaSalle Str
Chicago, Illinois 60602

Vigoro® is a registered trademark of Estech, Inc.

100%
SEVIN DUST

BEST AVAILABLE COPY