

Cameraman's Note

Poor Copy

DIRECTIONS FOR USE

Swift GSE Malathion contains 5 lbs. Malathion per gallon and is formulated to enable the user to prepare dilutions of varying concentrations. The amount of water depends upon the type of spray equipment used, the amount of vegetation and pest to be controlled. The amount of Swift GSE Malathion used should conform to the recommendations given, regardless of the amount of water necessary to obtain proper and even coverage of the area or plants to be treated.

CROPS TO BE TREATED	INSECTS CONTROLLED	RATE PER ACRE	INTERVAL TO HARVEST
ALFALFA & CLOVER	Aphids, Potato Leafhopper, Spider Mites, Minute Weevil, Armyworms, Spittlebugs, Grasshoppers, Downy Mildew, Acanthosoma, Leafhopper, Armyworms. Do not apply during full bloom. Use higher rates for taller alfalfa and clover.	1 1/2 Pts.	7 days
BARLEY, OATS, RYE, WHEAT (Other Forms)	English Grain Aphid, Green Bug, Green Leaf Beetle. Apply as needed and use sufficient water for good coverage. Armyworms.	1 1/2 Pts. 2 Pts.	7 days 7 days
RICE (Other Form)	Rice Leaf Miner. Apply soon after rice plants appear on surface of the water. Rice Stink Bug. Apply by airplane during early milk or dough stage and repeat as necessary.	2 1/2 Pts. 1-1 1/2 Pts.	7 days 7 days
SUGAR BEETS	Aphids, Spider Mites. Apply as needed and repeat if necessary. Do not use within 7 days of harvest if tops are to be used for feed.	1 1/2 Pts.	-
TOBACCO	Aphids (Field) Aphids (Plant Beds). Do not apply within 3 days of transplanting. Apply at rate of 5 gallons of finished spray per 100 sq. ft.	1-1 1/2 Pts. 2 Pts./100 gals. of water	7 days 7 days
ASPARAGUS	Asparagus Beetles Thrips Apply on foliage as needed	2 Pts. 1 1/2 Pts.	1 day 1 day
BEANS	Mexican Bean Beetle, Leafhopper, Spider Mites, Japanese Beetle, Aphid, Cucumber Beetle.	1 1/2 Pts.	7 days

BEETS	Aphids	1 1/2 Pts.	7 days
BRUSSEL SPROUTS, CAULIFLOWER, BROCCOLI, KALE, FENNEL, TOMATO, SPINACH, CUCUMBERS, CABBAGES	Aphids, Imported Cabbage Worm, Cabbage Looper, Leafhoppers	1-2 Pts.	7 days
CELERY	Aphids, Spider Mites	1 1/2 Pts.	7 days
COLLARDS, SWISS CHARD, PARSLEY	Aphids	1 1/2 Pts.	21 days
CUCUMBERS, SQUASH, MELONS, PUMPKINS	Aphids, Spider Mites, Pickleworm, Cucumber Beetles. Do not apply to cucurbits unless plants are dry. Squash Vine Borer.	1 1/2 Pts. 3 Pts.	7 days 7 days
EGGPLANT	Aphids, Spider Mites, Lace Bug.	1 Pt.	7 days
LETTUCE, SPINACH	Aphids, Mites, Cabbage Looper, Leafhoppers. Do not apply on leaf lettuce within 14 days of harvest.	2 Pts.	7 days
ONIONS, GARLIC	Thrips Onion Maggots	1 1/2 Pts. 2 1/2 Pts.	7 days 7 days
PEAS	Pea Aphid, Pea Weevil. Apply in early bloom period before eggs are laid and repeat as necessary.	1 1/2 Pts.	7 days
OKRA	Aphids Japanese Beetles Do not apply after pods have started to form.	1 1/2 Pts. 2 Pts.	-
POTATOES	Aphids, Leafhoppers, Mealybugs False Chinch Bugs	1-1 1/2 Pts. 1 1/2 Pts.	-
TOMATOES	Spider Mites, Aphids, Tomato Pinworm, Drosophila.	2 1/2 Pts.	7 days
STRAWBERRIES	Aphids, Spider Mites, Lygus Bug, Spittlebug, Field Crickets Potato Leafhopper, Strawberry Leafhopper, Whiteflies Strawberry Root Weevil Work in the top 6 to 8 inches of soil before planting.	1 1/2-3 Pts. 1 1/2-2 1/2 Pts.	7 days 7 days

		Rate per 100 Gallons of Water Unless Otherwise Indicated	
	Woolly Apple Aphid, Mealy Plum Scale, Cherry Fruit Fly, Forbes Scale, Hull Moth, San Jose Scale.	1 Pt.	3 days
	Green Apple Aphid, Rose Apple Aphid, Black Cherry Aphid, Cherry Fruit Tree Leaf Miner, Tent Caterpillar.	1 1/2 Pts.	3 days
	Mealybug, European Red Mite, Clover Mite, Wax Scale, Pear Tree Scale, Plum Curculionid, Red-banded Leafhopper, Pear Psylla, Pear Fruit Tree Leaf Miner.	2 Pts.	3 days
APPLES, PEARS, CHERRIES, PLUMS & PRUNES	Cover trees completely. Malathion in summer sprays may cause injury to McIntosh and Cortland varieties of apples. Bosc Pears and certain varieties of sweet cherries. Apply as needed and repeat at 10 day intervals until insects are controlled.		
	Codling Moth, Orange Tortrix, European Lecanium Scale, Soft Brown Scale, Aphids.	1 1/2 Pts.	7 days
APRICOTS, PEACHES	Oriental Fruit Moth, Terrapin Scale, Cotton Peach Scale.	2 Pts.	7 days
	Green Peach Aphid, Black Cherry Aphid, Black Peach Aphid, Rusty Plum Aphid.	1 Pt.	7 days
PECANS	Spider Mites, Aphids.	1-2 Pts.	7 days
	California Red Scale, Yellow Scale, Purple Scale, Black Scale (Small Branch), Soft Scale, Citricola Scale.	1-1 1/2 Pts.	7 days
GRAPEFRUIT, LEMONS, LIME, ORANGES, TANGERINES, YANGELOS, KUMQUATS	Florida Red Scale and Purple Scales (light and medium infestations).	2 Pts.	7 days
	Thrips Do not apply when trees are in bloom.	2 1/2 Pts./200 Gal./Acre	7 days

*Apply as needed and repeat as necessary.

5571873

		Rate per 100 Gallons of Water Unless Otherwise Indicated	
APPLES, PEARS, CHERRIES, PLUMS & PRAWNS	Woolly Apple Aphid, Mealy Plum Aphid, Pear Frost Fly, Forbes Scales, Bull Moth, San Jose Scale.	1 Pt.	3 days
	Green Apple Aphid, Pear Apple Aphid, Black Cherry Aphid, Cherry Fruit Tree Leaf Roller, Tent Caterpillar.	1 1/2 Pts.	3 days
	Mealybug, European Red Mite, Clover Mite, Walnuttite Mite, Two Spotted Mite, Codling Moth, Plum Curculio, Red banded Leaf Roller, Pear Psyllid, Pear Fruit Tree Leaf Roller.	2 Pts.	3 days
	Cover trees completely. Malathion in summer sprays may cause injury to McIntosh and Cortland varieties of apples, Bosc Pears and certain varieties of sweet cherries. Apply as needed and repeat at 10 day intervals until insects are controlled.		
APRICOTS, PEACHES	Codling Moth, Orange Tortrix, European Lecanium Scale, Soft Brown Scale, Aphids.	1 1/2-2 Pts.	*7 days
	Oriental Fruit Moth, Terrestrial Scale, Cotton Peach Scale.	2 Pts.	*7 days
	Green Peach Aphid, Black Cherry Aphid, Black Peach Aphid, Rusty Plum Aphid.	1 Pt.	*7 days
PECANS	Spider Mites, Aphids.	1-2 Pts.	*0 days
GRAPEFRUIT, LIME, ORANGES, TANGORINES, YACHTS, KUMQUATS	California Red Scale, Yellow Scale, Purple Scale, Black Scale (Single branched), Soft Scale, Citricola Scale.	1-1 1/2 Pts.	*7 days
	Florida Red Scale and Purple Scales (light and medium infestations).	2 Pts.	*7 days
	Thrips Do not apply when trees are in bloom.	2 1/2 Pts./ 200 Gal./ Acre	*7 days

*Apply as needed and repeat as necessary.

NET CONTENTS GALLONS

SWIFT

G5E MALATHION

CONTAINS 5 LBS. MALATHION PER GALLON

WARNING—FLAMMABLE!
KEEP AWAY FROM HEAT OR OPEN FLAME

ACTIVE INGREDIENTS:
Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate) 58.2%
Heavy Aromatic Naphtha 35.5%
INERT INGREDIENTS: 9.3%
E.P.A. Reg. No. 557-1873 Total 100.0%
E.P.A. Est. No. 557-FL-1

CAUTION: Keep Out Of Reach Of Children.
See Side Panels For Additional Cautions.

Manufactured By
SWIFT CHEMICAL COMPANY
(An Essoch Company)
CHICAGO, ILLINOIS 60604

CROPS TO BE TREATED	INSECTS CONTROLLED	RATE PER ACRE	INTERVAL TO HARVEST
ORNAMENTALS	Oyster Shell Scale, Lace Bug, Fungus Scale, Apple, Wax Moth, Spider Mites, Whitefly, Pear Leafhopper, Potato Leafhopper, Tarnished Plant Bug, Thrips, Glasswinged Fly, European Pear Shot Moth, Scaly Scale, Birch Tree Bark Beetle, Elm Wood Leaf Miner, Bagworms, Tent Caterpillars, Armyworms, Pear Nemes, Pine Tree Scale, Magnolia Scale, Fleeter Scale, Pear Scale, Juniper Scale, Black Scale, Crawlers, Monterey Pine Scale, Soft Scale, Pine Needle Scale.	2 1/2 Pts. or 2 teaspontfuls per gallon of finished spray	0 days
PASTURE AND RANGE GRASS	Grasshoppers	1-2 Pts.	7 days
CORN (Grain or Forage)	Aphids, Sap Beetles, Thrips, Grasshoppers (Young), Corn Rootworm Adults, Corn Leaf Beetle. Repeat as necessary.	1 1/2 Pts.	5 days

FLY CONTROL VII OUTDOORS

Use 1 Quart in 12 gallons of water (2 gallons in 100 gallons of water for spraying larger areas) and apply the spray at the rate of 1 gallon per 1000 square feet (Area 1/2 x 20' on painted surfaces and 7 gallons per 1000 square feet on concrete surfaces where flies alight or congregate. Addition of sugar at the rate of 2 1/2 pounds per 100 gallons or 20 pounds per 1000 gallons of finished spray prolongs the activity of Malathion. Does not serve as an attractant. Do not use in milk rooms, where edible products are handled or where lactating animals or young calves are present. Avoid contamination of feed, food products, drinking fountains and feed troughs.

CAUTION

Harmful if swallowed. Avoid breathing vapors or spray mist. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep away from children and domestic animals. Store above 32° F.
This product is highly toxic to bees exposed to it in treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.
This product is toxic to fish, shrimp, crab and other wildlife. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.
Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

ANTIDOTE

ANTIDOTE IF SWALLOWED: CALL A PHYSICIAN IMMEDIATELY.
NOTE TO PHYSICIAN: Malathion may cause cholinesterase inhibition. Atropine is antidotal.